

MULTIPLE SUMMING OPERATORS ON $C(K)$ SPACES

DAVID PÉREZ-GARCÍA AND IGNACIO VILLANUEVA

ABSTRACT. In this paper, we characterize, for $1 \leq p < \infty$, the multiple $(p, 1)$ -summing multilinear operators on the product of $C(K)$ spaces in terms of their representing polymeasures. As consequences, we obtain a new characterization of $(p, 1)$ -summing linear operators on $C(K)$ in terms of their representing measures and a new multilinear characterization of \mathcal{L}_∞ spaces. We also solve a problem stated by M.S. Ramanujan and E. Schock, improve a result of H.P. Rosenthal and S.J. Szarek and give new results about polymeasures.

1. INTRODUCTION AND NOTATION

Motivated by the importance of the theory of absolutely summing linear operators, there have been some attempts to generalize this concept and the related results and tools to the multilinear setting. Most of the previous efforts in this direction use the following definition of multilinear $(q; p_1, \dots, p_n)$ -summing operator, for certain choices of q, p_i :

A multilinear operator $T : X_1 \times \dots \times X_n \longrightarrow Y$ is called $(q; p_1, \dots, p_n)$ -summing if there exists a constant $K > 0$ such that

$$(1) \quad \left(\sum_{i=1}^m \|T(x_i^1, \dots, x_i^n)\|^q \right)^{\frac{1}{q}} \leq K \prod_{j=1}^n \|(x_i^j)_{i=1}^m\|_{p_j}^\omega$$

for all choices of $m \in \mathbb{N}$ and $x_1^j, \dots, x_m^j \in X_j$.

The interested reader can consult [9], [20] or [22] and the references therein to know more about this class of operators.

Recently, F. Bombal and both authors in [5] and [25], and M.C. Matos in [19] have defined and studied the class of *multiple summing multilinear operators*, see Definition 2.1 (although the origin of this class goes back to [27]). This class extends the notion of p -summing operator to the multilinear setting in a different way, it behaves better in many ways than the previous definitions of p -summing multilinear operators, and seems to be the “right” generalization of the linear case for many applications.

In particular, we prove in [5], [23], [24] and [25] several multilinear generalizations of Grothendieck’s theorem and relations with nuclear and Hilbert-Schmidt multilinear operators that extend and generalize classical linear

Both authors were partially supported by DGICYT grant BMF2001-1284.

results. It is easy to see that this “good behavior” is not shared by the $(q; p_1, \dots, p_n)$ -summing operators defined as above.

In this paper we continue studying the multiple summing multilinear operators. We give a simple characterization of the multiple 1-summing operators and the multiple $(p, 1)$ -summing operators on the product of $C(K)$ spaces in terms of their representing polymeasure. As a particular case, we obtain a new characterization of $(p, 1)$ -summing operators defined on $C(K)$ spaces in terms of their representing measure. As an application we can prove the rather surprising Corollary 3.2. This corollary will be the main tool used in Proposition 3.4, where we improve a result of H.P. Rosenthal and S.J. Szarek. Another application of our results is Proposition 3.6, which gives a multilinear characterization of \mathcal{L}_∞ spaces related to the main result of [9].

Several results in this paper (particularly Theorem 2.2 or Proposition 3.1) show that the class of multiple p -summing multilinear operators is relatively “small”. Thus, these results are specially surprising when compared with the Grothendieck type theorems given in [5] which show that every multilinear operator from the product of \mathcal{L}_∞ spaces to an \mathcal{L}_1 space is multiple 2-summing, or that every multilinear operator from the product of \mathcal{L}_1 spaces to a Hilbert space is multiple 1-summing.

In addition, we use some results of [5] to establish Example 3.13, which solves a problem stated in [27], and also to give non trivial new results about polymeasures (Corollaries 3.18 and 3.21).

The notations and terminology used along the paper are standard in Banach space theory, as for instance in [12]. This book is also our main reference for basic facts, definitions and unexplained notation all along the paper. However, before going any further, we shall establish some terminology: \mathbb{K} will be the scalar field, which can be considered to be either the real or complex numbers. X_i, Y will always be Banach spaces. $\mathcal{L}(X, Y)$ will note the Banach space of bounded linear mappings from X to Y . For $n \geq 2$, $\mathcal{L}^n(X_1, \dots, X_n; Y)$ will be the Banach space of all the continuous n -linear mappings from $X_1 \times \dots \times X_n$ into Y . When $Y = \mathbb{K}$ we will omit it and, from now on, ‘operator’ will mean linear or multilinear ‘continuous mapping’. As usual, $X_1 \hat{\otimes}_\epsilon \dots \hat{\otimes}_\epsilon X_n$ stands for the (completion of the) injective tensor product of the Banach spaces X_1, \dots, X_n and $X_1 \hat{\otimes}_\pi \dots \hat{\otimes}_\pi X_n$ will note (the completion of) their projective tensor product. Given a Banach space X , B_X denotes its unit ball, X^* stands for its topological dual and ω^* for the weak-star topology in X^* .

Given X , $1 \leq p < \infty$ and a finite sequence $(x_i)_{i=1}^m \subset X$, we note

$$\|(x_i)_{i=1}^m\|_p^\omega = \sup \left\{ \left(\sum_{i=1}^m |\langle x^*, x_i \rangle|^p \right)^{\frac{1}{p}} : x^* \in B_{X^*} \right\}.$$

For $1 \leq p \leq q < \infty$, we write $\Pi_{(q,p)}(X, Y)$ for the Banach space of (q, p) -summing operators from X into Y , and $\pi_{(q,p)}(T)$ stands for the (q, p) -summing norm of $T \in \Pi_{(q,p)}(X, Y)$. When $q = p$ we have the p -summing operators, and notation then will be $\Pi_p(X, Y)$ and $\pi_p(T)$.

Let $1 \leq p \leq \infty$ and $\lambda > 1$. A Banach space X is said to be an $\mathcal{L}_{p,\lambda}$ space if, for every finite dimensional subspace $E \subset X$ there exists another finite dimensional subspace F , with $E \subset F \subset X$ and such that there exists an isomorphism $v : F \rightarrow \ell_p^{\dim F}$ with $\|v\| \|v^{-1}\| < \lambda$. We say that X is an \mathcal{L}_p space if it is an $\mathcal{L}_{p,\lambda}$ space for some $\lambda > 1$. Clearly, $L_p(\mu)$ is the basic example of an \mathcal{L}_p -space.

Given $n, m_1, \dots, m_n \in \mathbb{N}$, $(x_{i_1, \dots, i_n})_{i_1, \dots, i_n=1}^{m_1, \dots, m_n}$ denotes a multiindex sequence with the index i_j varying from 1 to m_j ($1 \leq j \leq n$). $\sum_{i_1, \dots, i_n=1}^{m_1, \dots, m_n} x_{i_1, \dots, i_n}$ will be the notation for $\sum_{i_1=1}^{m_1} \cdots \sum_{i_n=1}^{m_n} x_{i_1, \dots, i_n}$.

If $T : X_1 \times \cdots \times X_n \rightarrow Y$ is a multilinear operator, we write $AB(T) : X_1^{**} \times \cdots \times X_n^{**} \rightarrow Y^{**}$ for its so-called Aron-Berner extension, which in general is not unique (see [3], or [8] and the references therein, for basic facts and different equivalent formulations of the Aron-Berner extension).

Let Σ_j be the Borel σ -algebra of a compact space K_j , $1 \leq j \leq n$ (or, in general, a σ -algebra defined on a set Ω_j). A function $\gamma : \Sigma_1 \times \cdots \times \Sigma_n \rightarrow Y$ is a (countably additive) *polymeasure* if it is separately (countably) additive.

Given a polymeasure $\gamma : \Sigma_1 \times \cdots \times \Sigma_n \rightarrow Y$, as in the case $n = 1$, its *semivariation* is defined as the set function

$$\|\gamma\| : \Sigma_1 \times \cdots \times \Sigma_n \rightarrow [0, +\infty]$$

given by

$$\|\gamma\|(A_1, \dots, A_n) = \sup \left\{ \left\| \sum_{k_1=1}^{r_1} \cdots \sum_{k_n=1}^{r_n} a_{k_1}^1 \cdots a_{k_n}^n \gamma(A_{k_1}^1, \dots, A_{k_n}^n) \right\| \right\}$$

where the supremum is taken over all the finite Σ_j -partitions $(A_{k_j}^j)_{k_j=1}^{r_j}$ of A_j ($1 \leq j \leq n$) and all the collections $(a_{k_j}^j)_{k_j=1}^{r_j}$ in the unit ball of the scalar field.

Let us also recall that its *variation* is defined as the set function

$$v(\gamma) : \Sigma_1 \times \cdots \times \Sigma_n \rightarrow [0, +\infty]$$

given by

$$v(\gamma)(A_1, \dots, A_n) = \sup \left\{ \sum_{k_1=1}^{r_1} \cdots \sum_{k_n=1}^{r_n} \|\gamma(A_{k_1}^1, \dots, A_{k_n}^n)\| \right\}$$

where the supremum is taken over all the finite Σ_j -partitions $(A_{k_j}^j)_{k_j=1}^{r_j}$ of A_j ($1 \leq j \leq n$).

In general, given $1 \leq p < \infty$, we can define its p -variation as the set function

$$v_p(\gamma) : \Sigma_1 \times \cdots \times \Sigma_n \longrightarrow [0, +\infty]$$

given by

$$v_p(\gamma)(A_1, \dots, A_n) = \sup \left\{ \left(\sum_{k_1=1}^{r_1} \cdots \sum_{k_n=1}^{r_n} \|\gamma(A_{k_1}^1, \dots, A_{k_n}^n)\|^p \right)^{\frac{1}{p}} \right\}$$

where the supremum is again taken over all the finite Σ_j -partitions $(A_{k_j}^j)_{k_j=1}^{r_j}$ of A_j ($1 \leq j \leq n$).

If γ has finite semivariation, an elementary integral $\int (f_1, f_2, \dots, f_n) d\gamma$ can be defined, where f_j are bounded, Σ_j -measurable scalar functions, just taking the limit of the integrals of n -uples of simple functions (with the obvious definition) uniformly converging to the f_j 's.

If K_1, \dots, K_n are compact Hausdorff spaces, then every multilinear operator $T \in \mathcal{L}^n(C(K_1), \dots, C(K_n); Y)$ has a unique *representing polymasure* $\gamma : \Sigma_1 \times \cdots \times \Sigma_n \rightarrow Y^{**}$ with finite semivariation, in such a way that

$$T(f_1, \dots, f_n) = \int (f_1, \dots, f_n) d\gamma \quad \text{for } f_j \in C(K_j),$$

and such that for every $y^* \in Y^*$, $y^* \circ \gamma$ is a separately regular, countably additive scalar polymasure. The idea behind this representation theorem can be easily described:

Given a compact Hausdorff space and its Borel σ -algebra Σ , we write $B(\Sigma)$ for the completion under the supremum norm of the space $S(\Sigma)$ of the Σ -simple scalar valued functions. It is well known that $C(K) \xrightarrow{1} B(\Sigma) \xrightarrow{1} C(K)^{**}$, where $\xrightarrow{1}$ denotes isometric embedding. So, for the operator T we consider its Aron-Berner extension to the product of the biduals $AB(T)$ (which is unique in this case) and restrict it to $\bar{T} : B(\Sigma_1) \times \cdots \times B(\Sigma_n) \rightarrow Y^{**}$. Now we define $\gamma(A_1, \dots, A_n) = \bar{T}(\chi_{A_1}, \dots, \chi_{A_n})$. In fact, as for the case of $C(K)$ spaces, easier reasonings yield an isometric isomorphism between $\mathcal{L}^n(B(\Sigma_1), \dots, B(\Sigma_n); Y)$ and $bpm(\Sigma_1, \dots, \Sigma_n; Y)$, the Banach space of the polymasures with bounded semivariation defined on $\Sigma_1 \times \cdots \times \Sigma_n$ with values in Y , endowed with the semivariation norm (see [6] and the references therein for more information about polymasures and the representation theorem).

2. DEFINITION AND FIRST RESULTS

We start recalling our definition.

Definition 2.1. Let $1 \leq p_1, \dots, p_n \leq q < +\infty$. A multilinear operator $T : X_1 \times \cdots \times X_n \rightarrow Y$ is *multiple $(q; p_1, \dots, p_n)$ -summing* if there exists a

constant $K > 0$ such that, for every choice of sequences $(x_{i_j}^j)_{i_j=1}^{m_j} \subset X_j$ the following relation holds

$$(2) \quad \left(\sum_{i_1, \dots, i_n=1}^{m_1, \dots, m_n} \|T(x_{i_1}^1, \dots, x_{i_n}^n)\|^q \right)^{\frac{1}{q}} \leq K \prod_{j=1}^n \|(x_{i_j}^j)_{i_j=1}^{m_j}\|_{p_j}^\omega.$$

In that case, we define the *multiple $(q; p_1, \dots, p_n)$ -summing norm* of T by

$$\pi_{(q; p_1, \dots, p_n)}(T) = \min\{K : K \text{ verifies (2)}\}.$$

A multiple $(q; p, \dots, p)$ -summing operator will be called *multiple (q, p) -summing*, and we write $\pi_{(q, p)}$ for the associated norm. Moreover, a multiple (p, p) -summing operator will be called *multiple p -summing* and we write π_p for the associated norm. The class $\Pi_{(q; p_1, \dots, p_n)}^n(X_1, \dots, X_n; Y)$ of multiple $(q; p_1, \dots, p_n)$ -summing multilinear operators is easily seen to be a Banach space with its norm $\pi_{(q; p_1, \dots, p_n)}$.

As in the linear case, if there exists $1 \leq j \leq n$ such that $p_j > p$, only the zero operator can satisfy (2). This is the reason to introduce the hypothesis $1 \leq p_1, \dots, p_n \leq q < +\infty$. Let us start showing the most basic example of this class of operators. Let $T : X_1 \times \dots \times X_n \rightarrow Y$ be a multilinear operator. Suppose that T is continuous for the ϵ topology and that its linearization $\hat{T} : X_1 \hat{\otimes}_\epsilon \dots \hat{\otimes}_\epsilon X_n \rightarrow Y$ is (q, p) -summing. Then, it follows easily from the definitions that T is multiple (q, p) -summing. In particular, for any $x_j^* \in X_j^*$, the multilinear form $(x_1^* \otimes \dots \otimes x_n^*)$ defined by $(x_1^* \otimes \dots \otimes x_n^*)(x_1, \dots, x_n) = x_1^*(x_1) \cdot \dots \cdot x_n^*(x_n)$ is multiple (q, p) -summing for any $1 \leq p \leq q \leq \infty$. It is probably worth mentioning that, in general, multilinear *forms* need not be multiple p -summing, as follows from Propositions 3.1 and [19].

Note that in this definition we require the sum

$$\left(\sum_{i_1=1}^{m_1} \dots \sum_{i_n=1}^{m_n} \|T(x_{i_1}^1, \dots, x_{i_n}^n)\|^q \right)^{\frac{1}{q}}$$

to be controlled by the product $\prod_{j=1}^n \|(x_{i_j}^j)_{i_j=1}^{m_j}\|_{p_j}^\omega$, whereas in the definition of $(q; p_1, \dots, p_n)$ -summing operators mentioned in the introduction and used previously by other authors, it is the “diagonal” sum

$$\left(\sum_{i=1}^m \|T(x_i^1, \dots, x_i^n)\|^q \right)^{\frac{1}{q}}$$

that must be controlled by the same product.

We show first the good behavior with respect to the extensions to the bidual that our operators share with the (q, p) -summing linear operators. Recall that the Aron-Berner extension of a multilinear operator is, in many ways, the natural generalization of the bitranspose of a linear operator. In that sense, the notion of weakly compact linear operator extends to the notion of multilinear operator whose Aron-Berner extension remains in the

image space. Following exactly the steps given in the proof of [14, Theorem 2.2] we obtain.

Theorem 2.2. *Let $T : X_1 \times \cdots \times X_n \longrightarrow Y$ be a multiple p -summing multilinear operator. Then, its Aron-Berner extension $AB(T)$ belongs to $\mathcal{L}(X_1^{**}, \dots, X_n^{**}; Y)$.*

We also have the following result which we will later need.

Theorem 2.3. *Let $1 \leq p_1, \dots, p_n \leq q < \infty$. A multilinear operator $T : X_1 \times \cdots \times X_n \longrightarrow Y$ is multiple $(q; p_1, \dots, p_n)$ -summing if and only if its Aron-Berner extension is multiple $(q; p_1, \dots, p_n)$ -summing.*

Moreover, in that case

$$\pi_{(q; p_1, \dots, p_n)}(T) = \pi_{(q; p_1, \dots, p_n)}(AB(T)).$$

The proof is obvious once we prove the following

Lemma 2.4. *Let X be a Banach space, $n \in \mathbb{N}$ and $1 \leq p < \infty$. Let $(z_i)_{i=1}^m \subset X^{**}$. Then, there exist a directed set Ω and nets $(x_\alpha^i)_{\alpha \in \Omega} \subset X$ such that*

$$x_\alpha^i \xrightarrow{\omega^*} z_i \quad \text{for every } 1 \leq i \leq n$$

and such that

$$\|(x_\alpha^i)_{i=1}^m\|_p^\omega \leq \|(z_i)_{i=1}^m\|_p^\omega \quad \text{for every } \alpha \in \Omega.$$

Proof. According to [11, Proposition 8.1], we know that the mapping given by $(y_i)_{i=1}^m \mapsto \sum_{i=1}^m e_i \otimes y_i$ establishes, for every Banach space Y , an isometric isomorphism between the Banach space of sequences of m vectors of Y , endowed with the norm $\|\cdot\|_p^\omega$, and $\ell_p^m \otimes_\epsilon Y$. Moreover the following isometric embeddings hold:

$$\ell_p^m \otimes_\epsilon X \hookrightarrow \ell_p^m \otimes_\epsilon X^{**} \hookrightarrow (\ell_p^m \otimes_\epsilon X)^{**}.$$

Since $(z_i)_{i=1}^m \subset \ell_p^m \otimes_\epsilon X^{**} \subset (\ell_p^m \otimes_\epsilon X)^{**}$, there exist a directed set Ω and a net $(w_\alpha)_{\alpha \in \Omega} \subset \ell_p^m \otimes_\epsilon X$ such that

$$w_\alpha \xrightarrow{\omega^*} (z_i)_{i=1}^m \quad \text{and} \quad \|w_\alpha\| \leq \|(z_i)_{i=1}^m\|_p^\omega.$$

Let x_α^i be such that $w_\alpha = \sum_{i=1}^m e_i \otimes x_\alpha^i$. We have that

$$\|(x_\alpha^i)_{i=1}^m\|_p^\omega = \|w_\alpha\| \leq \|(z_i)_{i=1}^m\|_p^\omega$$

and that, for every $x^* \in X^*$,

$$\langle x^*, x_\alpha^i \rangle = \langle e_i^* \otimes x^*, w_\alpha \rangle \xrightarrow{\alpha \in \Omega} \langle e_i^* \otimes x^*, \sum_{k=1}^m e_k \otimes z_k \rangle = z_i.$$

□

The following proposition can be easily proved as [19, Proposition 2.5].

Proposition 2.5. *Let $T : X_1 \times \cdots \times X_n \longrightarrow Y$ be a multilinear operator, let $1 \leq k \leq n - 1$ and let $T_k : X_1 \times \cdots \times X_k \longrightarrow \mathcal{L}^{n-k}(X_{k+1}, \dots, X_n; Y)$ be the associated k -linear operator.*

If $T_k \in \Pi_{(q;p_1, \dots, p_k)}^k(X_1, \dots, X_k; \Pi_{(q;p_{k+1}, \dots, p_n)}^{n-k}(X_{k+1}, \dots, X_n; Y))$, then $T \in \Pi_{(q;p_1, \dots, p_n)}^n(X_1, \dots, X_n; Y)$ and $\pi_{(q;p_1, \dots, p_n)}(T) \leq \pi_{(q;p_1, \dots, p_k)}(T_k)$.

We will see in Example 3.13 that, in general, the converse implication is not true. Nevertheless, it follows from Proposition 3.1 and [19] that the converse is true when $q = p_1 = \cdots = p_n = 1$ and all the X_j are $C(K)$ spaces (or in general \mathcal{L}_∞ spaces), or when $q = p_1 = \cdots = p_n = 2$ and all the X_j and Y are Hilbert spaces.

We state the following composition theorem for reference purposes, its proof, which can be seen in [5], follows along the lines of [12, 2.22].

Theorem 2.6. *Let $u_j \in \Pi_q(X_j, Y_j)$ and $T \in \Pi_p^n(Y_1, \dots, Y_n; Z)$ and let $1 \leq r < +\infty$ be such that $\frac{1}{r} = \frac{1}{p} + \frac{1}{q}$. Then $S = T(u_1, \dots, u_n)$ is multiple r -summing and $\pi_r(S) \leq \pi_p(T) \prod_{j=1}^n \pi_q(u_j)$.*

3. THE MAIN RESULTS

Given two Banach spaces X and Y , we will note by $I(X, Y)$ the space of integral linear operators from X to Y . It is a Banach space with the integral norm $\|\cdot\|_{int}$ (see [13, page 232] for the definitions).

A multilinear operator $T \in \mathcal{L}^n(X_1, \dots, X_n; Y)$ is said to be integral if there exists a regular Y^{**} -valued Borel measure G of bounded variation on the product $B_{X_1^*} \times \cdots \times B_{X_n^*}$ such that

$$T(x_1, \dots, x_n) = \int_{B_{X_1^*} \times \cdots \times B_{X_n^*}} x_1^*(x_1) \cdots x_n^*(x_n) dG(x_1^*, \dots, x_n^*)$$

for all $(x_1, \dots, x_n) \in X_1 \times \cdots \times X_n$. The space of integral multilinear operators $\mathcal{L}_I^n(X_1, \dots, X_n; Y)$ is a Banach space with the norm $\|T\|_{int} = \inf\{v(G), \text{ where } G \text{ represents } T \text{ as above}\}$.

These operators were defined in [30] (where they are called G-integral), although the definition is just a technical modification of a previous definition in [2]. In [30] it is proved that a multilinear operator $T : X_1 \times \cdots \times X_n \longrightarrow Y$ is integral if and only if its linearization \hat{T} is continuous for the ϵ topology and $\hat{T} : X_1 \hat{\otimes}_\epsilon \cdots \hat{\otimes}_\epsilon X_n \longrightarrow Y$ is an integral operator. Moreover, in that case $\|T\|_{int} = \|\hat{T}\|_{int}$.

We can prove now the following.

Proposition 3.1. *Let K_1, \dots, K_n be compact Hausdorff spaces, let $T : C(K_1) \times \cdots \times C(K_n) \longrightarrow Y$ be a multilinear operator and let γ be its representing polymeasure. Then, the following are equivalent:*

- i) T is multiple 1-summing.
- ii) $v(\gamma) < \infty$.
- iii) T is integral.

- iv) $T_1 \in \Pi_1(C(K_1), \Pi_1(C(K_2), \dots, \Pi_1(C(K_{n-1}), \Pi_1(C(K_n), Y)) \dots)))$.
 Moreover, in that case, all the norms coincide, i.e.

$$\pi_1(T) = v(\gamma) = \|T\|_{int} = \pi_1(T_1).$$

Proof. (i) \Rightarrow (ii) follows immediately from Theorem 2.3 and the fact that, if (Ω, Σ) is a measurable space and $(A_i)_{i=1}^m$ is a partition of Ω , then the sequence $(\chi_{A_i})_{i=1}^m \subset B(\Sigma)$ verifies $\|(\chi_{A_i})_{i=1}^m\|_1^\omega \leq 1$. The equivalence between (ii) and (iii) follows from [7, Corollary 4.2] and (iii) \Rightarrow (iv) is a consequence of [30, Proposition 2.9]. Finally, (iv) \Rightarrow (i) follows from Proposition 2.5. \square

As an immediate consequence we obtain the very surprising

Corollary 3.2. *Let X_j, Y_j and Z be Banach spaces ($1 \leq j \leq n$). Let $u_j \in \Pi_2(X_j, Y_j)$ and $T \in \Pi_2^n(Y_1, \dots, Y_n; Z)$. Then $S = T(u_1, \dots, u_n)$ is integral and*

$$\|S\|_{int} \leq \pi_2(T) \prod_{j=1}^n \pi_2(u_j).$$

Proof. It follows from the linear factorization theorem for 2-summing operators [12, Corollary 2.16] the existence of compact spaces K_j and 2-summing operators $b_j : C(K_j) \rightarrow Y_j$ such that $u_j = b_j \circ i_j$, where $i_j : X_j \hookrightarrow C(K_j)$ are isometric inclusions ($1 \leq j \leq n$). Let us consider the operator $R = T(b_1, \dots, b_n) \in \mathcal{L}^n(C(K_1), \dots, C(K_n); Z)$. Applying Theorem 2.6 and Proposition 3.1 we get that R is integral. Our result follows suit. \square

Remark 3.3. After the first version of this paper was written we have been able to prove that the operator S in Corollary 3.2 is actually nuclear (see [24]).

We can apply this corollary to prove a proposition that improves one of the results in [29] (see the remark below). We will say that a Banach space Y is a GT space, or that Y satisfies Grothendieck's theorem, if every linear operator from Y to ℓ_2 is 1-summing. According to Grothendieck's Theorem, \mathcal{L}_1 spaces are GT spaces, but there are several instances of GT spaces which are not \mathcal{L}_1 -spaces, for example L_1/H^1 or the quotient of an L_1 space by a subspace isomorphic to a Hilbert space (see [26]). All the known examples of GT spaces have cotype 2, and it remains an open question whether this must always happen.

Proposition 3.4. *For $1 \leq j \leq n$, let X_j be a \mathcal{L}_∞ space, Y_j a GT space with cotype 2 and $u_j : X_j \rightarrow Y_j$ a linear operator. Then, the operator*

$$u_1 \otimes \dots \otimes u_n : X_1 \hat{\otimes}_\epsilon \dots \hat{\otimes}_\epsilon X_n \rightarrow Y_1 \hat{\otimes}_\pi \dots \hat{\otimes}_\pi Y_n$$

is well defined and continuous.

Proof. By [7], it is sufficient to prove that, for every $T \in \mathcal{L}^n(Y_1, \dots, Y_n)$, the composition $T(u_1, \dots, u_n) \in \mathcal{L}_I^n(X_1, \dots, X_n)$. It is shown in [5] that T is multiple 2-summing and, by [12, Theorem 11.14], u_j is 2-summing for every j . Therefore, an appeal to Corollary 3.2 finishes the proof. \square

Remark 3.5. In [29], H.P. Rosenthal and S.J. Szarek mention that it would be desirable to determine pairs of (classes of) Banach spaces for which the conclusion of Proposition 3.4 holds. They obtained the result (in the case $n = 2$) for \mathcal{L}_∞ and \mathcal{L}_1 spaces. In that case, a direct proof can be given using induction. It is well known (see [15, Proposition 7] for a proof) that the projective tensor product of \mathcal{L}_1 spaces is an \mathcal{L}_1 space, and that the injective tensor product of \mathcal{L}_∞ spaces is an \mathcal{L}_∞ space. Therefore, all we have to do is to prove the case $n = 2$. Let X_1, X_2 be \mathcal{L}_∞ spaces, let Y_1, Y_2 be \mathcal{L}_1 spaces and let $u_j : X_j \rightarrow Y_j$ be a linear operator ($j = 1, 2$). As in Proposition 3.4, we have to prove that $S = T(u_1, u_2) : X_1 \times X_2 \rightarrow \mathbb{K}$ is integral for every $T \in \mathcal{L}^2(Y_1, Y_2)$. This is equivalent to prove that the associated linear operator $S_1 : X_1 \rightarrow X_2^*$ is integral. Now, we have the decomposition $S_1 = u_2^* \circ T_1 \circ u_1$. By Grothendieck's theorem [12, Theorem 3.7], u_1 and u_2^* are 2-summing. Then, [12, Theorem 2.22] tells us that S_1 is 1-summing and therefore integral [28, Theorem III.3].

It must be noticed that this argument gives also the case $n = 2$ of Proposition 3.4. However, the general case cannot be obtained by this simple induction reasoning since GT and cotype 2 spaces are not stable under projective tensor products. In fact, by [26, Theorem 10.6], there exists a GT space X with cotype 2 such that $X \hat{\otimes}_\pi X = X \hat{\otimes}_\epsilon X$. By [16, Remark 1] and [12, Theorem 14.1], this implies that $X \hat{\otimes}_\pi X$ does not have finite cotype and therefore (see [26, Corollary 6.13] and [12, Theorem 14.5]) $X \hat{\otimes}_\pi X$ cannot be a GT space.

Proposition 3.1 also allows us to give a new multilinear characterization of \mathcal{L}_∞ spaces.

Proposition 3.6. *Given X_1, \dots, X_n Banach spaces, the following are equivalent.*

- i) X_1, \dots, X_n are \mathcal{L}_∞ spaces.
- ii) *For every Banach space Y and for every multiple 1-summing n -linear operator $T : X_1 \times \dots \times X_n \rightarrow Y$, we have that T is integral.*

Proof. To see that *ii*) implies *i*) we consider an arbitrary Banach space Y and an arbitrary absolutely summing linear operator $u : X_1 \rightarrow Y$. By [28, Theorem III.3], if we prove that u is integral, we will obtain that X_1 is an \mathcal{L}_∞ space (we reason identically for $2 \leq j \leq n$). For $2 \leq j \leq n$ we consider $x_j \in B_{X_j}$ and $x_j^* \in B_{X_j^*}$ such that $x_j^*(x_j) = 1$. It is trivial that $T = u \otimes x_2^* \otimes \dots \otimes x_n^* : X_1 \times \dots \times X_n \rightarrow Y$ is multiple 1-summing. Using the hypothesis, we have that $\hat{T} : X_1 \hat{\otimes}_\epsilon \dots \hat{\otimes}_\epsilon X_n \rightarrow Y$ is integral, and so is $u = \hat{T}v$, where $v : X_1 \rightarrow X_1 \hat{\otimes}_\epsilon \dots \hat{\otimes}_\epsilon X_n$ is given by $v(x_1) = x_1 \otimes x_2 \otimes \dots \otimes x_n$.

To see that *i*) implies *ii*), we reason for the case $n = 2$ (the general case can be obtained similarly by induction). Choose a bilinear operator $T : X_1 \times X_2 \rightarrow Y$, and let $T_1 : X_1 \rightarrow \mathcal{L}(X_2, Y)$ be its associated linear operator. Using standard localization arguments we can deduce from Proposition 3.1 that, if $T \in \Pi_1^2(X_1, X_2; Y)$, then $T_1 \in \Pi_1(X_1, \Pi_1(X_2, Y))$.

Now, [28, Theorem III.3] tells us that $T_1 \in I(X_1, I(X_2, Y))$ and, by [30], we can conclude that T is integral. \square

Remark 3.7. Since 1-dominated multilinear operators (see [20] for definition and basic facts) are easily seen to be multiple 1-summing, Theorem 3.6 is weaker in one direction and stronger in the other direction than the main result in [9].

Next we are going to prove our main result relating multiple $(p, 1)$ -summing multilinear operators with the p -variation of their representing poly-measure.

Theorem 3.8. *Let (Ω_j, Σ_j) ($1 \leq j \leq n$) be measurable spaces, let $1 \leq p < \infty$ and let Y be a Banach space. Consider a multilinear operator $T : B(\Sigma_1) \times \cdots \times B(\Sigma_n) \rightarrow Y$ with representing polymeasure $\gamma : \Sigma_1 \times \cdots \times \Sigma_n \rightarrow Y$. Then T is multiple $(p, 1)$ -summing if and only if $v_p(\gamma) < \infty$. Moreover, in that case*

$$\begin{aligned} v_p(\gamma) &\leq \pi_{(p,1)}(T) \leq 2^{n(1-\frac{1}{p})} v_p(\gamma) && \text{(real case)} \\ v_p(\gamma) &\leq \pi_{(p,1)}(T) \leq 2^{n(2-\frac{1}{p})} v_p(\gamma) && \text{(complex case)} \end{aligned}$$

Proof. Let us first suppose that T is multiple $(p, 1)$ -summing and let us consider Σ_j partitions $(A_{k_j}^j)_{k_j=1}^{r_j}$ of Ω_j ($1 \leq j \leq n$). For every $\mu_j \in B(\Sigma_j)^*$ with $\|\mu_j\| \leq 1$ we have $\sum_{k_j=1}^{r_j} |\mu_j(A_{k_j}^j)| \leq 1$. Therefore

$$\begin{aligned} &\left(\sum_{k_1=1}^{r_1} \cdots \sum_{k_n=1}^{r_n} \|\gamma(A_{k_1}^1, \dots, A_{k_n}^n)\|^p \right)^{\frac{1}{p}} = \\ &= \left(\sum_{k_1=1}^{r_1} \cdots \sum_{k_n=1}^{r_n} \|T(\chi_{A_{k_1}^1}, \dots, \chi_{A_{k_n}^n})\|^p \right)^{\frac{1}{p}} \leq \pi_{(p,1)}(T). \end{aligned}$$

We prove now the converse in the real case, the complex case follows easily considering real and imaginary parts. Using density, it is enough to check for sequences in $S(\Sigma_i)$. So, let $(f_{i_j}^j)_{i_j=1}^{m_j} \subset S(\Sigma_j)$ ($1 \leq j \leq n$). There exist Σ_j -partitions $(A_{k_j}^j)_{k_j=1}^{r_j}$ of Ω_j ($1 \leq j \leq n$) and real numbers a_{i_j, k_j}^j such that

$$f_{i_j}^j = \sum_{k_j=1}^{r_j} a_{i_j, k_j}^j \chi_{A_{k_j}^j}.$$

Claim 1: $\|(f_{i_j}^j)_{i_j=1}^{m_j}\|_1^\omega \leq 1$ if and only if $\|a_j\| \leq 1$, where $a_j : \ell_1^{r_j} \rightarrow \ell_1^{m_j}$ is the operator defined by

$$a_j(e_{k_j}) = \sum_{i_j=1}^{m_j} a_{i_j, k_j}^j e_{i_j}.$$

Proof of the claim: Let us first suppose that $\|(f_{i_j}^j)_{i_j=1}^{m_j}\|_1^\omega \leq 1$, and consider $(c_{k_j})_{k_j=1}^{r_j} \in B_{\ell_1^{r_j}}$. For each $1 \leq k_j \leq r_j$, choose $\omega_{k_j} \in A_{k_j}^j$ and let $\mu_j = \sum_{k_j=1}^{r_j} c_{k_j} \delta_{\omega_{k_j}}$, where $\delta_{\omega_{k_j}}$ is the evaluation in ω_{k_j} . Then $\mu_j \in B_{B(\Sigma_j)^*}$ and

$$\|a_j((c_{k_j})_{k_j=1}^{r_j})\| = \sum_{i_j=1}^{m_j} \left| \sum_{k_j=1}^{r_j} c_{k_j} a_{i_j, k_j}^j \right| = \sum_{i_j=1}^{m_j} |\mu_j(f_{i_j}^j)| \leq 1,$$

which finishes this part of the proof.

For the converse, suppose $\|a_j\| \leq 1$ and choose $\mu_j \in B_{B(\Sigma_j)^*}$. Clearly $\sum_{k_j=1}^{r_j} |\mu_j(A_{k_j}^j)| \leq 1$ and we get

$$\sum_{i_j=1}^{m_j} |\mu_j(f_{i_j}^j)| = \sum_{i_j=1}^{m_j} \left| \sum_{k_j=1}^{r_j} a_{i_j, k_j}^j \mu_j(A_{k_j}^j) \right| = \|a_j((\mu_j(A_{k_j}^j))_{k_j=1}^{r_j})\| \leq 1,$$

which finishes the proof of the claim.

We consider now the (non-linear) mapping

$$F : \mathcal{L}(\ell_1^{r_1}, \ell_1^{m_1}) \times \cdots \times \mathcal{L}(\ell_1^{r_n}, \ell_1^{m_n}) \longrightarrow \mathbb{R}$$

defined by

$$F(c_1, \dots, c_n) = \left(\sum_{i_1, \dots, i_n=1}^{m_1, \dots, m_n} \left\| \sum_{k_1, \dots, k_n=1}^{r_1, \dots, r_n} c_{i_1, k_1}^1 \cdots c_{i_n, k_n}^n \gamma(A_{k_1}^1, \dots, A_{k_n}^n) \right\|^p \right)^{\frac{1}{p}}$$

where $c_{i_j, k_j}^j = \langle c_j(e_{k_j}), e_{i_j} \rangle$.

It is easy to see that F is continuous and separately convex. Therefore, its maximum in the compact set $B_{\mathcal{L}(\ell_1^{r_1}, \ell_1^{m_1})} \times \cdots \times B_{\mathcal{L}(\ell_1^{r_n}, \ell_1^{m_n})}$ is attained on the product of extremal points (b_1, \dots, b_n) .

Claim 2: If $b_j \in \text{ext}(B_{\mathcal{L}(\ell_1^{r_j}, \ell_1^{m_j})})$ then, for every $k_j \in \{1, \dots, r_j\}$, there exist $i_j(k_j) \in \{1, \dots, m_j\}$ and $\epsilon_{k_j}^j \in \{1, -1\}$ such that $b_{i_j, k_j}^j = \epsilon_{k_j}^j \delta_{i_j(k_j)}^{i_j}$. Obviously, $i_j(k_j)$ and $\epsilon_{k_j}^j$ are unique.

Proof of the claim:

If there is a k_j^0 such that $(b_{i_j, k_j^0}^j)_{i_j=1}^{m_j}$ is not of the form $\epsilon_{k_j^0}^j e_{i_j(k_j^0)}$, then $(b_{i_j, k_j^0}^j)_{i_j=1}^{m_j}$ is not an extremal point of $B_{\ell_1^{m_j}}$. Consequently, there exist two different sequences $(y_{i_j}^j)_{i_j=1}^{m_j}, (z_{i_j}^j)_{i_j=1}^{m_j} \in B_{\ell_1^{m_j}}$ with $b_{i_j, k_j^0}^j = \frac{1}{2} y_{i_j}^j + \frac{1}{2} z_{i_j}^j$ for all $i_j = 1, \dots, m_j$.

By doing

$$y_{i_j, k_j}^j = \begin{cases} b_{i_j, k_j}^j & k_j \neq k_j^0 \\ y_{i_j}^j & k_j = k_j^0 \end{cases} \quad z_{i_j, k_j}^j = \begin{cases} b_{i_j, k_j}^j & k_j \neq k_j^0 \\ z_{i_j}^j & k_j = k_j^0 \end{cases}$$

we have that $b_{i_j, k_j}^j = \frac{1}{2}y_{i_j, k_j}^j + \frac{1}{2}z_{i_j, k_j}^j$ for every i_j, k_j , that $(y_{i_j, k_j}^j)_{i_j, k_j} \neq (z_{i_j, k_j}^j)_{i_j, k_j}$ and that $(y_{i_j, k_j}^j)_{i_j, k_j}, (z_{i_j, k_j}^j)_{i_j, k_j} \in B_{\mathcal{L}(\ell_1^j, \ell_1^{m_j})}$. In conclusion, b_j is not extremal, which finishes the proof of the claim.

So, we have

$$\begin{aligned} & \left(\sum_{i_1, \dots, i_n=1}^{m_1, \dots, m_n} \|T(f_{m_1}^1, \dots, f_{m_n}^n)\|^p \right)^{\frac{1}{p}} = F(a_1, \dots, a_n) \leq F(b_1, \dots, b_n) = \\ & = \left(\sum_{i_1, \dots, i_n=1}^{m_1, \dots, m_n} \left\| \sum_{\{k_1: i_1(k_1)=i_1\}} \dots \sum_{\{k_n: i_n(k_n)=i_n\}} \epsilon_{k_1}^1 \dots \epsilon_{k_n}^n \gamma(A_{k_1}^1, \dots, A_{k_n}^n) \right\|^p \right)^{\frac{1}{p}} \leq \\ & \leq \left(\sum_{i_1, \dots, i_n=1}^{m_1, \dots, m_n} \left(\sum_{\phi \in \Phi} \left\| \sum_{k_1 \in \Gamma_{i_1, \phi(1)}^1} \dots \sum_{k_n \in \Gamma_{i_n, \phi(n)}^n} \gamma(A_{k_1}^1, \dots, A_{k_n}^n) \right\|^p \right) \right)^{\frac{1}{p}} \end{aligned}$$

with

$$\begin{aligned} \Gamma_{i_j, +}^j &= \{k_j : i_j(k_j) = i_j \text{ and } \epsilon_{k_j}^j = 1\} \\ \Gamma_{i_j, -}^j &= \{k_j : i_j(k_j) = i_j \text{ and } \epsilon_{k_j}^j = -1\} \end{aligned}$$

and Φ the set of mappings from $\{1, \dots, n\}$ to $\{+, -\}$.

We note by $B_{i_j, +}^j = \cup_{k_j \in \Gamma_{i_j, +}^j} A_{k_j}^j$ and by $B_{i_j, -}^j = \cup_{k_j \in \Gamma_{i_j, -}^j} A_{k_j}^j$. We have that, for each j , the sets $B_{i_j, +}^j$ and $B_{i_j, -}^j$ are all disjoint. So,

$$\begin{aligned} & \left(\sum_{i_1, \dots, i_n=1}^{m_1, \dots, m_n} \|T(f_{i_1}^1, \dots, f_{i_n}^n)\|^p \right)^{\frac{1}{p}} \\ & \leq \left(\sum_{i_1, \dots, i_n=1}^{m_1, \dots, m_n} \left(\sum_{\phi \in \Phi} \left\| \sum_{k_1 \in \Gamma_{i_1, \phi(1)}^1} \dots \sum_{k_n \in \Gamma_{i_n, \phi(n)}^n} \gamma(A_{k_1}^1, \dots, A_{k_n}^n) \right\|^p \right) \right)^{\frac{1}{p}} \\ & = \left(\sum_{i_1, \dots, i_n=1}^{m_1, \dots, m_n} \left(\sum_{\phi \in \Phi} \left\| \gamma(B_{i_1, \phi(1)}^1, \dots, B_{i_n, \phi(n)}^n) \right\|^p \right) \right)^{\frac{1}{p}} \\ & \leq 2^{n(1-\frac{1}{p})} \left(\sum_{i_1, \dots, i_n=1}^{m_1, \dots, m_n} \sum_{\phi \in \Phi} \left\| \gamma(B_{i_1, \phi(1)}^1, \dots, B_{i_n, \phi(n)}^n) \right\|^p \right)^{\frac{1}{p}} \\ & \leq 2^{n(1-\frac{1}{p})} v_p(\gamma) \end{aligned}$$

□

Using Theorems 2.3 and 3.8 and the comments above about polymeasures, it is very easy to obtain the $C(K)$ version of Theorem 3.8

Theorem 3.9. *Let K_j be compact Hausdorff spaces, Y a Banach space and $T : C(K_1) \times \cdots \times C(K_n) \rightarrow Y$ a multilinear operator with representing polymeasure $\gamma : \Sigma_1 \times \cdots \times \Sigma_n \rightarrow Y^{**}$. Then, T is multiple $(p, 1)$ -summing if and only if $v_p(\gamma) < \infty$.*

Moreover, in this case,

$$v_p(\gamma) \leq \pi_{(p,1)}(T) \leq 2^{n(1-\frac{1}{p})} v_p(\gamma) \quad (\text{real case})$$

$$v_p(\gamma) \leq \pi_{(p,1)}(T) \leq 2^{n(2-\frac{1}{p})} v_p(\gamma) \quad (\text{complex case})$$

Remark 3.10. The case $n = 1$ of Theorem 3.9 gives a new characterization of $(p, 1)$ -summing linear operators from $C(K)$ spaces in terms of their representing measure.

As a corollary, we obtain a new proof of a classical result ([21, page 14])

Corollary 3.11. *Let K be a compact Hausdorff space, $p \geq 1$ and Y a Banach space. A linear operator $T : C(K) \rightarrow Y$ is $(p, 1)$ -summing if and only if*

$$(3) \quad \sup \left\{ \left(\sum_{i=1}^m \|T(f_i)\|^p \right)^{\frac{1}{p}} : (f_i)_{i=1}^m \in B_{C(K)} \text{ with disjoint supports} \right\} < \infty.$$

Proof. First of all, it should be noticed that, if $(f_i)_{i=1}^m$ have disjoint supports, then $\|(f_i)_{i=1}^m\|_1^\omega = \max_i \|f_i\|$. So, by theorem 3.9, it is enough to see that $v_p(\gamma)$ is less or equal than (3), where $\gamma : \Sigma \rightarrow Y^{**}$ is the representing measure of T . The proof of this fact for $p = 1$ can be seen in [13, page 163]. The general case can be obtained with obvious modifications. \square

Remark 3.12. The constant $2^{n(1-\frac{1}{p})}$ in the real case for Theorems 3.8 and 3.9 is optimal. To see this, we can consider $T : \ell_\infty^2 \times \cdots \times \ell_\infty^2 \rightarrow \mathbb{R}$ given by $T((x_1, y_1), \dots, (x_n, y_n)) = \prod_{j=1}^n (y_n - x_n)$. In the complex case, however, we do not know (even in the case $n=1$) what the optimal constant is.

It is now a natural question whether we can obtain a result similar to Proposition 3.1 for multiple $(p, 1)$ -summing multilinear operators. The answer is no and the clue is [5, Theorem 3.2] (see Theorem 3.17 below).

Example 3.13. Let X, Y, Z be infinite dimensional \mathcal{L}_∞ spaces. Then we have that

$$\Pi_{(2,1)}(X, \Pi_{(2,1)}(Y, Z^*)) \subsetneq \Pi_{(2,1)}^2(X, Y; Z^*).$$

Proof. Using a version of Grothendieck's theorem ([12, Theorem 3.7]), we know that $\Pi_{(2,1)}(Y, Z^*)$ is isomorphic to $(Y \otimes_\pi Z)^*$. Moreover, it follows from Dvoretzki's theorem that, for any $\epsilon > 0$, $Y^* \hat{\otimes}_\epsilon Z^*$ contains the ℓ_∞^n 's $(1 + \epsilon)$ -uniformly (see [16, Remark 1]). Since $Y^* \hat{\otimes}_\epsilon Z^*$ is isometrically embedded

into $(Y \hat{\otimes}_\pi Z)^*$, we get that $(Y \hat{\otimes}_\pi Z)^*$ contains the ℓ_∞^n 's $(1 + \epsilon)$ -uniformly (complemented).

Let $i_n : \ell_\infty^n \hookrightarrow (Y \otimes_\pi Z)^*$, $p_n : (Y \otimes_\pi Z)^* \rightarrow \ell_\infty^n$ be such that $p_n i_n = Id_{\ell_\infty^n}$ and $\|i_n\| = 1$, $\|p_n\| \leq 2$. Let X be an $\mathcal{L}_{\infty, \lambda}$ space, then, for every $n \in \mathbb{N}$, we can consider projections $R_n : X \rightarrow \ell_\infty^n$ with $\|R_n\| \leq \lambda$ and $\pi_{(2,1)}(R_n) \geq \sqrt{n}$.

For every $n \in \mathbb{N}$, we consider the operator $T_n = i_n R_n : X \rightarrow (Y \otimes_\pi Z)^*$ and its associated bilinear operator $\bar{T}_n : X \times Y \rightarrow Z^*$. Since Z^* is an \mathcal{L}_1 space, it has cotype 2. So, [5, Theorem 3.2] tells us that there exists $C > 0$ such that, for every $n \in \mathbb{N}$,

$$\pi_{(2,1)}(\bar{T}_n) \leq C \|T_n\| \leq \lambda C.$$

As $\pi_{(2,1)}(R_n) \geq \sqrt{n}$, we have that $\pi_{(2,1)}(T_n) = \pi_{(2,1)}(i_n R_n) \geq \frac{\sqrt{n}}{2}$. This proves the non equivalence of the corresponding norms, and, hence, the existence of an operator $T \in \Pi_{(2,1)}^2(X, Y; Z^*)$ such that its associated operator $T_1 : X \rightarrow \Pi_{(2,1)}(Y, Z^*)$ is not $(2, 1)$ -summing.

To give an explicit counterexample, let $X = c_0$, $Y = Z = \ell_\infty$. Then $(Y \hat{\otimes}_\pi Z)^*$ contains an isomorphic copy of c_0 (see [1]) and we can consider $T : X \times Y \rightarrow Z^*$ as the bilinear operator associated to $T_1 : c_0 \hookrightarrow (Y \hat{\otimes}_\pi Z)^*$. \square

Remark 3.14. In fact, if we use the multilinear version of Grothendieck's theorem given in [5, Theorem 3.1] instead of [5, Theorem 3.2], we can prove, with the same argument, the existence of a multiple 2-summing bilinear operator $T : X \times Y \rightarrow Z$ such that $T_1 \notin \Pi_2(X, \Pi_2(Y, Z))$, solving a question stated in [27].

Now, we are going to extend to the multilinear setting another linear property that extends the field of application of the above results. First we need a proposition, whose proof follows immediately from the definitions.

Proposition 3.15. *Let $T : X_1 \times \dots \times X_n \rightarrow Y$ be a multilinear operator and let $1 \leq p_1, \dots, p_n \leq q < \infty$. The following are equivalent*

- i) T is multiple $(q; p_1, \dots, p_n)$ -summing
- ii) There exists a constant $K > 0$ such that for every $m_2, \dots, m_n \in \mathbb{N}$ and every choice of sequences $(x_{i_j}^j)_{i_j=1}^{m_j} \subset X_j$, with $\|(x_{i_j}^j)_{i_j=1}^{m_j}\|_{p_j}^\omega \leq 1$ ($2 \leq j \leq n$), we have that the associated linear operator

$$S : X_1 \rightarrow \ell_q^{m_2 \dots m_n}(Y)$$

given by

$$S(x_1) = (T(x_1, x_{i_2}^2, \dots, x_{i_n}^n))_{i_2, \dots, i_n=1}^{m_2, \dots, m_n}$$

is (q, p_1) -summing and it verifies

$$(4) \quad \pi_{(q, p_1)}(S) \leq K.$$

In that case, $\pi_{(q; p_1, \dots, p_n)}(T) = \min\{K : K \text{ verifies (4)}\}$

Proposition 3.16. *Let $1 \leq p_1, \dots, p_n < q < \infty$ and let K_1, \dots, K_n be compact Hausdorff spaces. A multilinear operator $T : C(K_1) \times \dots \times C(K_n) \longrightarrow Y$ is multiple $(q, 1)$ -summing if and only if it is multiple $(q; p_1, \dots, p_n)$ -summing.*

Proof. We reason in the bilinear case, the reasonings being similar in the general case. Suppose $T : C(K_1) \times C(K_2) \longrightarrow Y$ is multiple $(q, 1)$ -summing. Then, for any sequence $(x_{i_1}^1)_{i_1=1}^{m_1} \subset C(K_1)$ such that $\|(x_{i_1}^1)_{i_1=1}^{m_1}\|_1^\omega \leq 1$, the operator $S : C(K_2) \longrightarrow \ell_q^m(Y)$ defined as in Proposition 3.15 is $(q, 1)$ -summing and verifies that

$$\pi_{(q,1)}(S) \leq \pi_{(q,1)}(T).$$

Let $i : \ell_q^m(Y) \longrightarrow \ell_q(Y)$ be the natural inclusion. Applying [12, Theorem 10.9] to $i \circ S$, and using the injectivity of the operator ideal of the (q, p) summing operators, we get that S is (q, p_2) -summing and that

$$\pi_{(q,p_2)}(S) \leq K\pi_{(q,1)}(T)$$

where the constant K does not depend on the choice of $(x_{i_1}^1)_{i_1=1}^{m_1}$.

Therefore, T is multiple $(q; 1, p_2)$ -summing. We choose now any sequence $(x_{i_2}^2)_{i_2=1}^{m_2} \subset C(K_2)$ such that $\|(x_{i_2}^2)_{i_2=1}^{m_2}\|_{p_2}^\omega \leq 1$ and reason similarly. \square

To end the paper, we are going to state some results concerning the p -variation of polymeasures. The starting point is [5, Theorem 3.2], that says

Theorem 3.17. *Let X_j be a Banach space for $1 \leq j \leq n$ and let Y be a cotype q space. Then, every multilinear operator $T : X_1 \times \dots \times X_n \longrightarrow Y$ is multiple $(q, 1)$ -summing and*

$$\pi_{(q,1)}(T) \leq C_q(Y)^n \|T\|$$

where $C_q(Y)$ is the cotype q constant of Y .

Using this result, the proof of the following surprising corollary is trivial.

Corollary 3.18. *Let Y be a cotype q space and $\gamma : \Sigma_1 \times \dots \times \Sigma_n \longrightarrow Y$ a polymeasure of bounded semivariation. Then $v_q(\gamma) \leq C_q(Y)^n \|\gamma\|$. In particular, every scalar polymeasure of bounded semivariation has bounded 2-variation.*

Note that, in general, scalar polymeasures do not have bounded variation (see [7]).

We can improve the scalar case of the last two results. To this end, we consider the following classical theorem (see [4], [10], [17], [18]).

Theorem 3.19 (Littlewood-Bohnenblust-Hille). *If T is a continuous n -linear form on c_0 , then*

$$\left(\sum_{i_1, \dots, i_n=1}^{\infty} |T(e_{i_1}^1, \dots, e_{i_n}^n)|^{\frac{2n}{n+1}} \right)^{\frac{n+1}{2n}} \leq 2^{\frac{n-1}{2}} \|T\|.$$

This theorem allows us to prove the following

Corollary 3.20. *Let X_1, \dots, X_n be Banach spaces. Every n -linear form $T : X_1 \times \dots \times X_n \longrightarrow \mathbb{K}$ is multiple $(\frac{2n}{n+1}, 1)$ -summing and*

$$\pi_{(\frac{2n}{n+1}, 1)}(T) \leq 2^{\frac{n-1}{2}} \|T\|.$$

Proof. Let us consider, for $1 \leq j \leq n$, sequences $(x_{i_j}^j)_{i_j=1}^{m_j} \subset X_j$ with $\|(x_{i_j}^j)_{i_j=1}^{m_j}\|_1^\omega \leq 1$. The operator $u_j : \ell_\infty^{m_j} \longrightarrow X_j$ given by $u_j(e_{i_j}) = x_{i_j}^j$ verifies that

$$\|u_j\| = \|(x_{i_j}^j)_{i_j=1}^{m_j}\|_1^\omega \leq 1.$$

We can now apply Theorem 3.19 to the multilinear operator

$$S = T(u_1, \dots, u_n) : \ell_\infty^{m_1} \times \dots \times \ell_\infty^{m_n} \longrightarrow \mathbb{K}$$

to obtain that

$$\begin{aligned} \left(\sum_{i_1, \dots, i_n=1}^{m_1, \dots, m_n} |T(x_{i_1}^1, \dots, x_{i_n}^n)|^{\frac{2n}{n+1}} \right)^{\frac{n+1}{2n}} &= \left(\sum_{i_1, \dots, i_n=1}^{m_1, \dots, m_n} |S(e_{i_1}^1, \dots, e_{i_n}^n)|^{\frac{2n}{n+1}} \right)^{\frac{n+1}{2n}} \\ &\leq 2^{\frac{n-1}{2}} \|S\| \leq 2^{\frac{n-1}{2}} \|T\|. \end{aligned}$$

□

Finally, we have

Corollary 3.21. *Every scalar polymeasure $\gamma : \Sigma_1 \times \dots \times \Sigma_n \longrightarrow \mathbb{K}$ with bounded semivariation verifies that*

$$v_{\frac{2n}{n+1}}(\gamma) \leq 2^{\frac{n-1}{2}} \|\gamma\|.$$

REFERENCES

- [1] ALAMINOS, J., CHOI, Y.S., KIM, S.G. and PAYÁ, R., Norm attaining bilinear forms on spaces of continuous functions, *Glasgow Math. J.* **40** (1998), 359–365.
- [2] ALENCAR, R., Multilinear mappings of nuclear and integral type, *Proc. Amer. Math. Soc.* **94** (1985), 33–38.
- [3] ARON, R. and BERNER, P.D., A Hahn-Banach extension theorem for analytic mappings, *Bull. Soc. Math. France* **106** (1978), 3–24.
- [4] BOHNENBLUST, H.F. and HILLE, E., On the absolute convergence of Dirichlet series, *Ann. of Math.* **32** (1931), 600–622.
- [5] BOMBAL, F., PÉREZ-GARCÍA, D. and VILLANUEVA, I., Multilinear extensions of Grothendieck's theorem, *Preprint*.
- [6] BOMBAL, F. and VILLANUEVA, I., Multilinear operators on spaces of continuous functions, *Funct. Approx. Comment. Math.* **26** (1998), 117–126.
- [7] BOMBAL, F. and VILLANUEVA, I., Integral operators on the product of $C(K)$ spaces, *J. Math. Anal. Appl.* **264** (2001), 107–121.
- [8] CABELLO, F., GARCÍA, R. and VILLANUEVA, I., Extension of multilinear operators on Banach spaces, *Extrac. Math.* **15** (2000), 291–334.
- [9] CILIA, R., D'ANNA, M. and GUTIÉRREZ, J., Polynomial characterization of \mathcal{L}_∞ -spaces, *J. Math. Anal. Appl.* **275** (2002), 900–912.

- [10] DAVIE, A.M., Quotient algebras of uniform algebras, *J. London Math. Soc.* **7** (1973), 31–40.
- [11] DEFANT, A. and FLORET, K., *Tensor norms and operator ideals*, North-Holland, Amsterdam, 1993.
- [12] DIESTEL, J., JARCHOW, H. and TONGE, A., *Absolutely summing operators*, Cambridge Univ. Press, Cambridge, 1995.
- [13] DIESTEL, J. and UHL, J.J., *Vector measures*, Mathematical Surveys and Monographs **15**, Amer. Math. Soc., Providence, Rhode Island, 1977.
- [14] DIMANT, V., Strongly p -summing multilinear operators, *J. Math. Anal. Appl.* **278** (2003), 182–193.
- [15] GONZÁLEZ, M. and GUTIÉRREZ, J., The Dunford-Pettis property on tensor products, *Math. Proc. Camb. Phil. Soc.* **131** (2001), 185–192.
- [16] JOHN, K., Tensor powers of operators and nuclearity, *Math. Nachr.* **129** (1986), 115–121.
- [17] KAIJSER, S., Some results in the metric theory of tensor products, *Studia Math.* **63** (1978), 157–170.
- [18] LITTLEWOOD, J., On bounded bilinear forms in an infinite number of variables, *Q. J. Math.* **2** (1930), 167–171.
- [19] MATOS, M.C., Fully absolutely summing and Hilbert-Schmidt multilinear mappings, To appear in *Collect. Math.*
- [20] MATOS, M.C., On multilinear mappings of nuclear type, *Rev. Mat. Univ. Complut. Madrid* **6** (1993), 61–81.
- [21] MAUREY, B., *Type et cotype dans les espaces munis de structures locales inconditionnelles*, Séminaire Maurey-Schwartz 1973/74, Exp. XXIV-XXV, École Polytechnique, Paris.
- [22] MELÉNDEZ, Y. and TONGE, A., Polynomials and the Pietsch domination theorem, *Math. Proc. Roy. Irish Acad.* **99A** (1999), 195–212.
- [23] PÉREZ-GARCÍA, D., The inclusion theorem for multiple summing operators, *Preprint*.
- [24] PÉREZ-GARCÍA, D. and VILLANUEVA, I., A composition theorem for multiple summing operators, *Preprint*.
- [25] PÉREZ-GARCÍA, D. and VILLANUEVA, I., Multiple summing operators on Banach spaces, To appear in *J. Math. Anal. Appl.*
- [26] PISIER, G., *Factorization of linear operators and geometry of Banach spaces*, CBMS Regional Conference Series **60**, Amer. Math. Soc., Providence, Rhode Island, 1986.
- [27] RAMANUJAN, M.S. and SCHOCK, E., Operator ideals and spaces of bilinear operators, *Linear and Multilinear Algebra* **18** (1985), 307–318.
- [28] RETHERFORD, J.R. and STEGALL, C.P., Fully nuclear and completely nuclear operators with applications to \mathcal{L}_1 - and \mathcal{L}_∞ -spaces, *Trans. Amer. Math. Soc.* **163** (1972), 457–492.
- [29] ROSENTHAL, H.P. and SZAREK, S.J., On tensor products of operators from L^p to L^q , in *Functional Analysis* (E. Odell and H. Rosenthal, eds.), Lecture Notes in Math. **1470**, 108–132, Springer-Verlag, Berlin-Heidelberg, 1991.
- [30] VILLANUEVA, I., Integral mappings between Banach spaces, *J. Math. Anal. Appl.* **279** (2003), 56–70.

E-mail address: David.Perez@mat.ucm.es, ignaciov@mat.ucm.es

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, FACULTAD DE MATEMÁTICAS, UNIVERSIDAD COMPLUTENSE DE MADRID, MADRID 28040, SPAIN