



# When hadrons become unstable: A novel type of non-analyticity in chiral extrapolations

F.-K. Guo<sup>a</sup>, C. Hanhart<sup>b</sup>, F.J. Llanes-Estrada<sup>c,\*</sup>, U.-G. Meißner<sup>a,b</sup>

<sup>a</sup> Helmholtz-Institut für Strahlen- und Kernphysik and Bethe Center for Theoretical Physics, Universität Bonn, D-53115 Bonn, Germany

<sup>b</sup> Institut für Kernphysik, Institute for Advanced Simulation and Jülich Center for Hadron Physics, Forschungszentrum Jülich, D-52425 Jülich, Germany

<sup>c</sup> Dept. Física Teórica I, Universidad Complutense, 28040 Madrid, Spain

## ARTICLE INFO

### Article history:

Received 20 May 2011

Received in revised form 21 July 2011

Accepted 9 August 2011

Available online 12 August 2011

Editor: J.-P. Blaizot

### Keywords:

Analyticity of S-matrix

Lattice QCD

Chiral extrapolations

## ABSTRACT

Hadron masses show a specific dependence on the quark masses. Therefore, the variation of these masses can cause a resonance in a hadronic scattering amplitude to become a bound state. Consequently, the amplitude exhibits a non-analytic behavior at this transition. Crossed amplitudes, where the resonance can be exchanged in the  $t$ -channel, can be shown to exhibit the same phenomenon by  $s \rightarrow t$  analytic continuation. This entails possible kinks in lattice quark-mass extrapolations needed to compute hadronic observables.

© 2011 Elsevier B.V. All rights reserved.

## 1. Introduction

It is of current interest to obtain lattice Quantum Chromodynamics (QCD) predictions for hadronic observables, both to test QCD in the strong-coupling regime, and to compute QCD backgrounds to new physics searches. It is customary in these lattice gauge theory computations, due to the large numerical costs, to perform simulations with unphysically large masses of the light quarks. Then a smooth extrapolation formula to physical values, inspired by chiral perturbation theory is usually employed to obtain the physical results (for a recent review see Ref. [1]).<sup>1</sup> However, there are no theorems of S-matrix theory guaranteeing the analyticity of such an extrapolation for larger quark masses, denoted as  $m_q$  in what follows, beyond the regime where chiral perturbation theory is applicable.<sup>2</sup> Note that one can equally well talk of the pion mass instead of the quark mass, since they are related by the Gell-Mann–Oakes–Renner relation [6]  $m_\pi^2 f_\pi^2 = 2m_q(\bar{q}q) + \mathcal{O}(m_q^2)$ ,

\* Corresponding author.

E-mail address: fllanes@fis.ucm.es (F.J. Llanes-Estrada).

<sup>1</sup> We are well aware that first simulations with physical (or even lower) quark masses [2–4] become available, but these are still key exceptions. If future simulations are to access a large set of observables, working at unphysical pion masses will still be useful and economic if one has reliable extrapolation formulae. In addition, simulations with various quark masses can provide more information than those solely with the physical quark masses.


<sup>2</sup> It is well known that certain non-analyticities in the quark masses can be shown to hold for arbitrary momenta (see Ref. [5] and references therein), but these are not the effects we are dealing with.

where the corrections  $\sim m_q^2$  are known to be very small. On the contrary, in this Letter we expose non-analyticities (kinks in the  $m_q$ -extrapolation) that may arise when a resonance becomes bound upon varying  $m_q$ . For example, in the pion form factor or the  $B \rightarrow \pi\pi$  decay amplitude, the relevant resonance is the  $\rho(770)$ . For the weak  $K \rightarrow \pi$  transition form factor, the  $K^*$  matters. Let us stress that the value of  $s$  is of no concern; the non-analyticity in the variable  $m_q$ , if present, affects the entire amplitude or form factor. We will exemplify this with both time-like and space-like pion form factors. So far the analysis has been carried out for the  $m_q$  dependence of the resonance mass itself,  $m_\rho(m_q)$ , or its derivative, for example Refs. [7–9]. Here, we demonstrate the generality of the phenomenon affecting the computation of most hadron observables, which completes the preliminary results reported in [10].

Our results are relevant because there is much active lattice research in form factor determinations, see e.g. [11–13] and spectroscopy [14–17]. Extrapolation formulae are available for both form factors [18] and spectroscopy, e.g. [19]. Typically these extrapolations are smooth except for the usual chiral logarithms of the pion mass,  $\log(m_\pi^2/\mu^2)$ , that present a non-analyticity at  $m_\pi = 0$ .

## 2. Illustration: A simple model

To expose the feature in the simplest possible physical manner, we now focus on two-pion amplitudes, and the role of the  $\rho$ -resonance. In Section 3 it will however be shown that the results



**Fig. 1.** Contribution of  $\rho$ -meson one-loop vacuum polarization to  $\pi$  time-like form factor. Solid, double and wiggly lines denote in turn pions,  $\rho$ -mesons and photons.

are general. In a simple field theory where the two pions are coupled to the resonance and the latter is represented as an additional field, the threshold effect appears through the vacuum polarization of the resonance, whose imaginary part controls the decay width. Therefore, the size of the possible non-analyticities in any amplitude has to be proportional to the width of the resonance and appear only in, at least, one-loop calculations, such as depicted in Fig. 1 for the time-like pion form factor.

We therefore proceed to study the  $\pi$ - $\rho$  case at one loop. The bare  $\rho$ -meson propagator is  $1/(s - m_{0\rho}^2)$  – the one-loop vacuum polarization will renormalize the bare mass  $m_{0\rho}$  to  $m_\rho$ . The  $s$ -channel on-shell unitarization for the scattering amplitude reads

$$T^{11}(s) = \frac{V^{11}(s)}{1 - G(s)V^{11}(s)} = \frac{-\frac{4}{3}g_{\rho\pi\pi}^2|\mathbf{p}|^2}{z_0^\rho(s - m_{0\rho}^2) + \frac{4}{3}g_{\rho\pi\pi}^2|\mathbf{p}|^2G(s)} \quad (1)$$

in terms of the Born amplitude for  $\rho \rightarrow \pi\pi$  (with  $J = I = 1$ ),

$$V^{11}(s) = -\frac{4}{3} \frac{g_{\rho\pi\pi}^2}{z_0^\rho} \frac{|\mathbf{p}|^2}{s - m_{0\rho}^2}, \quad (2)$$

where the  $\rho$  wave function renormalization constant  $z_0^\rho$  comes from the relation between the bare coupling constant and the renormalized one. The factor  $|\mathbf{p}|^2 = s/4 - m_\pi^2$  stems from the on-shell p-wave derivative coupling. With this on-shell factorization, the denominator in Eq. (1) contains the unregularized scalar one-loop function

$$G(s) = \frac{1}{16\pi^2} \left( R + \log\left(\frac{m_\pi^2}{\mu^2}\right) + 1 - \bar{J}(s) \right). \quad (3)$$

We use the convention

$$\bar{J}(s) = 2 + \sigma \log\left(\frac{\sigma - 1}{\sigma + 1}\right) \quad (4)$$

with  $\sigma = 2|\mathbf{p}|/\sqrt{s} = \sqrt{1 - 4m_\pi^2/s}$  for the relativistic phase space. The divergence in dimensional regularization appears in

$$R = \frac{2}{d-4} - \log(4\pi) - \Gamma'(1) - 1$$


with the number of space-time dimensions  $d \rightarrow 4$ .

Returning to our main issue, consider the imaginary part of the vacuum polarization in the denominator of Eq. (1). It is given by

$$\text{Im} \Pi = \frac{\pi\sigma}{16\pi^2} \frac{4}{3} g_{\rho\pi\pi}^2 |\mathbf{p}|^2 \theta(s - 4m_\pi^2) \quad (5)$$

and leads to the well-known non-analyticity in  $s$  (branch point at  $s = 4m_\pi^2$ ). A trivial observation is that, reciprocally, there is a non-analyticity in  $m_\pi$  for fixed  $s$  at  $m_\pi = \sqrt{s}/4$ . More subtle is to notice that when  $2m_\pi = m_\rho(m_\pi)$  (in Appendix B, we show generally that such a situation will occur for the  $\rho$ ), a similar non-analyticity affects the amplitude for all  $s$  due to the dependence of the amplitude on the renormalized  $m_\rho$  (physical pole position) that suddenly changes from a bound state on the real axis to an unbound resonance.

We present in Fig. 2 the behavior of  $m_\rho(m_\pi)$ . The results agree qualitatively with those from the more sophisticated treatment



**Fig. 2.** Dependence of the  $\rho$  mass  $m_\rho$  (a) and its derivative  $dm_\rho/dm_\pi$  (b) on the pion mass obtained from Eq. (12). The dashed line in (a) denotes the motion of the  $\pi\pi$  threshold. Data are from a lattice calculation by the QCDSF Collaboration [20], and the physical  $\rho$  mass is represented by a thick circle.


in, e.g., Refs. [8,9]. Further details of the computation within the simple model are given in Appendix A. Eq. (12) contains the combination

$$\frac{1}{z_0^\rho} \text{Re} \bar{J}(m_\rho^2) \left( \frac{m_\rho^2}{4} - m_\pi^2 \right)$$

that is non-analytic for the value of the pion mass where the  $\rho$  becomes bound, cf. Eqs. (4), (18). This is seen especially as a kink in the derivative  $dm_\rho/dm_\pi$ , shown in Fig. 2(b). Therefore, the position of the pole in the pion scattering amplitude appears not to be an analytic function of the pion mass. The same phenomenon will appear in other amplitudes, such as the time-like electromagnetic form-factor. This can then be analytically continued (in  $s$ ) to the space-like side and the same phenomenon will appear for, say, the radius squared (which appears naturally in the low-energy expansion of the space-like form-factor). To illustrate this effect we now use the simplest realization of vector meson dominance (VMD), where the direct photon-pion coupling is neglected and the form factor is entirely given by the photon- $\rho$ -meson coupling to the intermediate resonance (with the strength  $g_{\rho\gamma}$ ) [21]. The tree level formula

$$F^{\text{tree}}(s) = \frac{m_\rho^2}{m_\rho^2 - s} \quad (6)$$

already suggests a kink if one substitutes the dependence of the pole mass  $m_\rho(m_\pi)$  that we have argued to be non-analytic. This



**Fig. 3.** The squared pion charge radius also presents a kink in its first derivative as a function of  $m_\pi$ , at the point where the  $\rho$ -resonance becomes bound. See Appendix A for computational details of the  $\rho\pi\pi$  model.

lack of analyticity appears then in the squared charge radius in the Breit frame  $\langle r^2 \rangle = 6/m_\rho^2$ . Although the mentioned VMD description is too simplistic to exhibit all pertinent features of the pion vector form factor, it is very useful for illustrative purposes. In Appendix A we demonstrate that the full one loop amplitude exhibits the same features and in the next section we demonstrate that our findings are indeed model independent.

The one-loop result for the charge radius squared as a function of the pion mass is displayed in Fig. 3. We have assumed that  $g_{\rho\pi\pi}$  is independent of the pion mass.<sup>3</sup> Thus, the threshold non-analyticity when the  $\rho$ -resonance becomes bound is inherited by the space-like form factor. This result may not seem intuitive, since the argument  $t$  of the space-like form factor is apparently very far from any thresholds associated with  $s$ , so perhaps it is helpful to think of the form factor as a function of two variables  $F(s, m_q)$ . The non-analyticity enters because of the implicit pion-mass dependence through  $m_\rho(m_q)$  and is not affected by the analytic continuation in the other variable  $s \rightarrow t$ .

### 3. Generalization of the results


We now turn to a model-independent discussion of the effect, introduced so far within a particular model, for a specific  $\pi\pi$  partial wave characterized by conserved angular momentum and isospin  $l = I = 1$ . In a model-independent way this effect can be studied by employing the Omnès representation for the form-factor as given e.g. in Refs. [23–26]. This renowned relation expresses the form factor as an integral over the scattering phase shift. In once-subtracted form it reads in the absence of bound states

$$F(t, m_\pi^2) = \Omega(t, m_\pi^2) = \exp\left(\frac{t}{\pi} \int_{4m_\pi^2}^{\infty} ds \frac{\delta_{11}(s, m_\pi^2)}{s(s-t-i\epsilon)}\right). \quad (7)$$

Then the charge radius in the Omnès representation is expressed in terms of the  $\pi\pi$  scattering phase shift as [25]

$$\langle r^2 \rangle = \frac{6}{\pi} \int_{4m_\pi^2}^{\infty} ds \frac{\delta_{11}(s, m_\pi^2)}{s^2}. \quad (8)$$

<sup>3</sup> The pion-mass dependence of  $g_{\rho\pi\pi}$  is very moderate from both the unitarized chiral perturbation theory [8] and very recent lattice simulations [22].



**Fig. 4.** Pion-mass dependence of  $\delta_{11}(s, m_\pi^2)$  from the one-loop model, as the pion mass approaches the value where the  $\rho$  becomes stable. Shown are the phases for  $m_\pi = 420$  MeV (dotted line), 427 MeV (dashed line) and 431 MeV (solid line).

In the presence of a bound state there is an additional singularity on the first sheet and thus the dispersion integral needs to be modified. It now reads [27],

$$F_b(t, m_\pi^2) = \left(1 + \frac{t g_{\gamma\rho} g_{\rho\pi\pi}}{m_\rho^2(m_\rho^2 - t)} \frac{1}{\Omega(m_\rho^2, m_\pi^2)}\right) \Omega(t, m_\pi^2). \quad (9)$$

Eq. (8) needs to be adapted accordingly

$$\langle r^2 \rangle_b = \frac{6}{\Omega(m_\rho^2, m_\pi^2)} \frac{g_{\gamma\rho} g_{\rho\pi\pi}}{m_\rho^4} + \frac{6}{\pi} \int_{4m_\pi^2}^{\infty} ds \frac{\delta_{11}(s, m_\pi^2)}{s^2}. \quad (10)$$

Here, we introduced the subscript  $b$  to distinguish the quantities defined in the presence of a bound state from those given in Eqs. (7), (8). We stress that the form factor is continuous at the value of the pion mass for which the  $\rho$  become a stable state, for all  $t \neq m_\rho^2$ . To see this we first observe that the integrals over the phases in Eqs. (8), (10) converge towards each other, as the  $\rho$  mass approaches the two-pion threshold. This follows directly from the behavior of the phases shown in Fig. 4 – as soon as the  $\rho$  appears as a stable state at  $m_\pi = 430$  MeV, according to Levinson's theorem the phase shift starts from  $\pi$ . In addition, when approaching the point  $m_\rho(m_\pi) = 2m_\pi$  from larger pion masses one finds

$$\begin{aligned} \log \Omega(t, m_\pi^2) &\simeq \frac{t}{\pi} \delta_{11} \int_{4m_\pi^2}^{\infty} ds \frac{1}{s(s-t-i\epsilon)} \\ &= \frac{\delta_{11}}{\pi} \log \frac{4m_\pi^2}{|4m_\pi^2 - t|}, \end{aligned} \quad (11)$$

where we used that in this limit  $\delta_{11}$  is a slowly varying function of  $s$  in the energy range of interest. Evidently,  $1/\Omega(t, m_\pi^2)$  vanishes, when  $t = 4m_\pi^2$ . However, as becomes apparent in Eq. (10), continuity of the function is not accompanied by continuity of its derivatives: the non-analyticity in, e.g.,  $m_\rho$  directly influences the quark-mass dependence of the squared radius [28].

Since causality permits the Omnès dispersive representation of form factors, and in particular the charge radius, it is not surprising *a posteriori* that the  $s$ -channel cusp already known in the literature affects these quantities. The  $\rho\pi\pi$  model in Section 2 leads to this effect by construction, since it is based on a one-loop Feynman perturbation theory diagram, that satisfies the Mandelstam analyticity in the  $s$  variable, so it is a particular instance of the general discussion here.

#### 4. Summary and conclusions

In this work we discussed a non-analyticity in the chiral extrapolation of physical quantities emerging when a change in the quark mass transforms a resonance – pole on the second sheet – to a physical state with a pole on the first sheet. We established that the analytic continuation in the kinematic variable  $s \rightarrow t$  does carry over the non-analyticity in  $m_\pi$  from the time-like to the space-like domain. This kind of behavior is model-independent.

In this Letter we analyzed the case of the pion form factor, but the same phenomenon should appear in other form factors. For example, in the  $K \rightarrow \pi$  weak vector transition form factor [29,30], when  $m_\pi \simeq 350$  MeV the  $K^*$  resonance should also become bound, and develop a non-analyticity. The same phenomenon appears also in  $c$ - and  $b$ -quark physics where resonance masses are much closer to or even below the thresholds with the physical pion mass.

The situation is even more interesting for scalar form factors, where one has a kink directly in the function (be it the mass or the squared radius) and not in its derivative with respect to the pion mass. This is simply because the factor  $|\mathbf{p}|^2$  from the  $J = 1$  derivative coupling is absent (as already discussed in detail in Ref. [23]). Then it will be easier for lattice data to isolate such a structure (that is not yet visible in existing simulations, see e.g. [13]). If a relative drop  $(\delta F^{(n)})/F^{(n)}$  in the  $n$ -th derivative of a function is to be identified visually, the error acceptable in the lattice computation of  $F$  itself is, as a rule of thumb,  $(\delta F)/F \leq (\delta F^{(n)})/(2^n F^{(n)})$  since each derivative with a good mid-point numerical method requires two evaluations. Hence, we would propose that the scalar pion form factor be computed with smaller statistical error bars and smaller  $t$ -intervals, as a favorable system to try to find the non-analyticity, given that there is no phase-space suppression and that the coupling  $g_{\sigma\pi\pi}$  is large. Calculations using unitarized chiral perturbation theory predict that the  $\sigma$ -meson becomes bound at about  $m_\pi \simeq 350$  MeV [8]. Full QCD simulations for scalar quantities at sufficiently low pion masses will, however, not appear in the near future, for those are a lot more computer-time intensive compared to the ones discussed here so far, due to the presence of disconnected diagrams.

We now examine to what extent non-analyticities have been stressed in earlier studies. Very old work focused on the particle virtuality for fixed mass, be it in perturbation theory or with the Lehman representation [31], or for scaling deeply-inelastic scattering functions [32]. In both cases the phenomenon of a resonance becoming a bound state during the particle mass variation is absent, and those authors found analyticities in the transferred momentum plane with the physical values of the particle masses. Closer in spirit to our work, features in quark-mass extrapolations due to presence of a threshold have already been discussed in Ref. [33], which focuses on the avoided level crossing in a finite volume. A small cusp in the pion-mass dependence of the mass of the  $\Delta$  resonance [34] is produced. In Ref. [35] it was stressed that the kind of non-analyticity discussed in Ref. [34] also shows up in electro-magnetic properties such as the magnetic moment for unstable particles. The effect we discussed introduces an *additional* non-analyticity in the radii *even for stable particles* as long as they couple to resonances that might become bound upon varying  $m_q$ . Finally, another kind of non-analyticity – possible discontinuities – in the chiral extrapolation of hadron masses was proposed in [36].

The non-analyticity that we uncover is a feature of continuum field theory, and not an artifact of lattice quantization. It is possible that accurate lattice data should be able to isolate these non-analyticities, provided the volume is large enough that the resonance is not bound by the minimum momentum possible on the lattice (though the alternative non-analyticities of Ref. [34] mask

the effect). They should be taken into account when attempting to extrapolate lattice data to physical pion masses when high precision is expected.

#### Acknowledgements

We thank Stephan Dürr and Akaki Rusetsky for useful discussions. This work was supported in part by grants provided by the HGF to the virtual institute “Spin and strong QCD” (VH-VI-231), the DFG (SFB/TR 16) and the EU I3HP “Study of Strongly Interacting Matter” under the Seventh Framework Program of the EU, FPA 2008-00592, 2007-29115-E, FIS2008-01323 (Spain) and 227431, HadronPhysics2 (EU). F.J.L.E. thanks the members of the Nuclear Theory Center at Forschungszentrum Jülich for their hospitality during this work. U.G.M. also thanks the BMBF for support (Grant No. 06BN9006).

#### Appendix A. One-loop renormalization of the $\rho\pi\pi$ model

Our choice of renormalization is meant to expose the pole mass and decay coupling constant in the amplitude, so this one is expressed in terms of directly measurable quantities. To achieve this, we add and subtract to the denominator of Eq. (1) the vacuum polarization evaluated at the (still unknown)  $\rho$ -pole mass, that is,

$$\frac{4}{3} g_{\rho\pi\pi}^2 G(m_\rho^2) \left( \frac{m_\rho^2}{4} - m_\pi^2 \right).$$

Imposing now the renormalization condition that the position of the pole in the denominator be at  $m_\rho$  yields the equation

$$m_\rho^2 = m_{0\rho}^2 - \frac{4}{3} \frac{g_{\rho\pi\pi}^2}{z_0^2} G(m_\rho^2) \left( \frac{m_\rho^2}{4} - m_\pi^2 \right). \quad (12)$$

Above the two-pion threshold,  $m_\rho$  is complex. But in the numerical calculations, for simplicity, we take  $m_\rho$  to be real. This amounts to neglecting  $\text{Im} G$  – which is a very good approximation near the kink where phase space closes. Note that  $G$  contains an infinity that needs to be absorbed into the bare mass. Since the divergence is multiplied by  $(m_\rho^2/4 - m_\pi^2)/z_0^2$  the subtraction procedure calls for introducing a pion-mass dependent mass term. Thus, since we want to keep the pion-mass dependence explicit, the Lagrangian density for the model needs to contain a counterterm proportional to  $m_\pi^2 \rho^\dagger \rho$ .

We therefore define the renormalized mass and its mass derivative with respect to  $m_\pi^2$  through

$$\bar{m}_{0\rho}^2 + \bar{m}'_{0\rho} m_\pi^2 = m_{0\rho}^2 - \frac{4}{3} \frac{g_{\rho\pi\pi}^2}{z_0^2} \left( \frac{m_\rho^2}{4} - m_\pi^2 \right) \frac{R+1}{16\pi^2} \quad (13)$$

and demand that the two constants  $\bar{m}_{0\rho}^2$  and  $\bar{m}'_{0\rho}$  be pion-mass independent. These two parameters can be fixed using the physical  $\rho$ -meson mass at the physical point for  $m_\pi$  and lattice data. We choose as renormalization scale, which enters Eq. (12) through  $G(m_\rho^2)$  (cf. Eq. (3)),  $\mu = m_\rho$ , the pole mass itself. Then  $z_0^2$  becomes known (see below), and Eq. (12) can be solved. The best fit to the lattice data from the QCDSF Collaboration [20] with the constraint from the physical  $\rho$  mass gives  $\bar{m}_{0\rho} = 0.707$  GeV, and  $\bar{m}'_{0\rho} = 1.13$ .

At this point we have guaranteed that the pion-pion scattering amplitudes has a pole at physical  $m_\rho$  for the physical pion mass, and we can compute the variation of the pole position with the pion mass if this dependence is known for  $z_0^2$ , so we also need to solve for it.

The second renormalization condition we impose is that  $g_{\rho\pi\pi}$  be the physical coupling at the  $\rho$  pole, obtainable from the residue of the pion scattering amplitude

$$\text{Res } T^{11}(s) = \lim_{s \rightarrow m_\rho^2} (s - m_\rho^2) T^{11}(s). \quad (14)$$

Imposing that the residue be

$$\text{Res } T^{11}(s) = -\frac{4}{3} g_{\rho\pi\pi}^2 \left( \frac{m_\rho^2}{4} - m_\pi^2 \right) \quad (15)$$

and taking into account that, in terms of the pole mass, we have

$$T^{11}(s) = \frac{-\frac{4}{3} g_{\rho\pi\pi}^2 |\mathbf{p}|^2}{z_0^\rho (s - m_\rho^2) + \Delta(s) + i \text{Im } \Pi(s)} \quad (16)$$

with

$$\Delta(s) \equiv \text{Re } \Pi(s) - \text{Re } \Pi(m_\rho^2), \quad (17)$$

where

$$\Pi(s) = \frac{1}{12\pi^2} g_{\rho\pi\pi}^2 \bar{J}(s) \left( \frac{s}{4} - m_\pi^2 \right).$$

We find

$$z_0^\rho = 1 - \frac{d\Delta(s=m_\rho^2)}{ds}. \quad (18)$$

Note the  $m_\rho$  in the last expression is not the physical value but  $m_\rho(m_\pi)$ , to guarantee that  $m_\rho(m_\pi)$  is always defined as the pole mass in the propagator. As discussed below Eq. (12), although  $m_\rho$  is complex, for simplicity we take  $m_\rho$  to be real.

Resumming the Dyson series originating from the vacuum polarization of the  $\rho$ -meson one obtains for the form factor

$$F(s) = \frac{-g_{\rho\gamma} g_{\rho\pi\pi} \sqrt{z_0^A/z_0^\rho}}{z_0^\rho (s - m_\rho^2) + \Delta(s) + i \text{Im } \Pi(s)}. \quad (19)$$

The denominator is of course the same as in Eq. (16), and since the numerator is real this guarantees the same phase for form factor and scattering amplitude. In order to get the proper normalization in this most simple formulation of VMD one needs to impose

$$g_{\rho\gamma} g_{\rho\pi\pi} \sqrt{z_0^A/z_0^\rho} = z_0^\rho m_\rho^2 - \Delta(0) \quad (20)$$

on the photon- $\rho$  coupling. The resulting form factor

$$F(s) = \frac{-z_0^\rho m_\rho^2 + \Delta(0)}{z_0^\rho (s - m_\rho^2) + \Delta(s) + i \text{Im } \Pi(s)} \quad (21)$$

satisfies now  $F(0) = 1$  and has the correct unitarity cut. Through all the one-loop quantities  $m_\rho$ ,  $z_0^\rho$  and  $\Delta$ , as shown in Fig. 2 and Fig. 5, the form-factor acquires a non-analyticity in  $m_\pi$ . This non-analyticity appears at one loop and is therefore proportional to  $g_{\rho\pi\pi}^2$  and hence the physical resonance width. Using Eq. (18) the squared charge radius becomes now

$$\langle r^2 \rangle = 6 \frac{z_0^\rho + \Delta'(0)}{z_0^\rho m_\rho^2 - \Delta(0)}. \quad (22)$$

The derivative of  $\Delta(s)$  at  $s = 0$  is a constant  $\Delta'(0) = -g_{\rho\pi\pi}^2 / (72\pi^2)$ .

## Appendix B. Position of the point $m_\rho(m_\pi) = 2m_\pi$

In this appendix, we will show that the  $\rho$  mass grows more slowly than the two-pion threshold when increasing the pion mass, and hence there must be a certain point after which the  $\rho$  pole will be below the two-pion threshold.




Fig. 5. Pion-mass dependence of auxiliary  $\Delta(0)$  (solid) and  $\Delta'(m_\rho^2) = 1 - z_0^\rho$  (dashed).

Expanding the  $\rho$  mass in terms of  $m_\pi$ , to the order  $\mathcal{O}(m_\pi^2)$  (it is sufficient for our purpose to work to this order; for the expansion to higher orders, see [19]), one has

$$m_\rho(m_\pi) = m_{\rho 0} + c_1 m_\pi^2 = m_{\rho 0} + 2c_1 B_0 \hat{m}, \quad (23)$$

where  $m_{\rho 0}$  is the  $\rho$  mass in the chiral limit,  $c_1$  is a low-energy constant related to the quark-mass term in the chiral expansion,  $B_0 = -\langle 0 | \bar{q}q | 0 \rangle / f_\pi^2$  and  $\hat{m} = (m_u + m_d)/2$ . Generally, since the  $\rho$  has a non-vanishing (and not small) mass even in the chiral limit, one has  $m_{\rho 0} > 2m_\pi$  for small values of the pion mass. The points where  $m_\rho$  coincides with  $2m_\pi$  are then simply given by the solutions of  $m_{\rho 0} + c_1 m_\pi^2 = 2m_\pi$ , i.e.

$$m_\pi = \frac{1}{c_1} (1 \pm \sqrt{1 - c_1 m_{\rho 0}}). \quad (24)$$

Hence a crossing happens if and only if

$$c_1 \leq \frac{1}{m_{\rho 0}}. \quad (25)$$

To determine  $c_1$  we resort to quark-mass controlled  $SU(3)$  breaking, and expand the mass of the  $K^*$  in analogy with Eq. (23)

$$m_{K^*}(m_\pi) = m_{\rho 0} + c_1 B_0 (m_s + \hat{m}) = m_{\rho 0} + c_1 m_K^2. \quad (26)$$

Because  $c_1$  is independent of the quark mass by definition, it can be used for unphysical pion masses after determining it using physical meson masses by

$$c_1 = \frac{m_{K^*} - m_\rho}{m_K^2 - m_\pi^2} = 0.51 \text{ GeV}^{-1}. \quad (27)$$

Therefore, as long as  $m_{\rho 0} \leq 1960$  MeV, the inequality  $c_1 \leq 1/m_{\rho 0}$  can be fulfilled. It is believed that  $m_{\rho 0}$  is not far from the physical mass  $m_\rho = 770$  MeV, so that the  $\rho$  mass will coincide with  $2m_\pi$  at some value(s) of  $m_\pi$ . One can even estimate that value. Taking, e.g.,  $m_{\rho 0} \approx 700$  MeV, the crossing point will be at around  $m_\pi \approx 400$  MeV, in agreement with recent detailed estimates [37].

Finally, we note that the second solution of Eq. (24) is far beyond the applicability region of the chiral expansion.

## References

- [1] V. Bernard, Prog. Part. Nucl. Phys. 60 (2008) 82.
- [2] S. Aoki, et al., PACS-CS Collaboration, Phys. Rev. D 81 (2010) 074503; S. Aoki, et al., PACS-CS Collaboration, Phys. Rev. D 79 (2009) 034503.

- [3] S. Dürr, et al., Phys. Lett. B 701 (2011) 265;  
S. Dürr, et al., arXiv:1011.2711 [hep-lat].
- [4] W. Bietenholz, et al., arXiv:1102.5300 [hep-lat].
- [5] J. Gasser, A. Zepeda, Nucl. Phys. B 174 (1980) 445.
- [6] M. Gell-Mann, R.J. Oakes, B. Renner, Phys. Rev. 175 (1968) 2195.
- [7] D.B. Leinweber, A.W. Thomas, K. Tsushima, S.V. Wright, Phys. Rev. D 64 (2001) 094502.
- [8] C. Hanhart, J.R. Pelaez, G. Rios, Phys. Rev. Lett. 100 (2008) 152001.
- [9] J. Nebreda, J.R. Pelaez, Contributed to the 12th International Conference on Meson–Nucleon Physics and the Structure of the Nucleon, arXiv:1010.3982 [hep-ph];  
J. Nebreda, J.R. Pelaez, G. Rios, Phys. Rev. D 83 (2011) 094011.
- [10] F. Llanes-Estrada, et al., PoS EFT09 (2009) 054.
- [11] P.A. Boyle, et al., JHEP 0807 (2008) 112.
- [12] R. Frezzotti, V. Lubicz, S. Simula, ETM Collaboration, Phys. Rev. D 79 (2009) 074506.
- [13] S. Aoki, et al., JLQCD Collaboration, TWQCD Collaboration, Phys. Rev. D 80 (2009) 034508.
- [14] S. Dürr, et al., Science 322 (2008) 1224.
- [15] H.W. Lin, et al., Hadron Spectrum Collaboration, Phys. Rev. D 79 (2009) 034502.
- [16] R. Baron, et al., JHEP 1006 (2010) 111.
- [17] G.P. Engel, et al., BGR Collaboration, Phys. Rev. D 82 (2010) 034505.
- [18] J. Bijnens, G. Colangelo, P. Talavera, JHEP 9805 (1998) 014.
- [19] P.C. Bruns, U.-G. Meißner, Eur. Phys. J. C 40 (2005) 97.
- [20] M. Göckeler, et al., QCDSF Collaboration, PoS LATTICE2008 (2008) 136.
- [21] J.J. Sakurai, Currents and Mesons, University of Chicago Press, Chicago, 1969.
- [22] X. Feng, K. Jansen, D.B. Renner, Phys. Rev. D 83 (2011) 094505.
- [23] J. Gasser, U.-G. Meißner, Nucl. Phys. B 357 (1991) 90.
- [24] J.F. De Troconiz, F.J. Yndurain, Phys. Rev. D 65 (2002) 093001.
- [25] J.A. Oller, L. Roca, Phys. Lett. B 651 (2007) 139.
- [26] B. Ananthanarayan, I. Caprini, I.S. Imsong, Phys. Rev. D 83 (2011) 096002.
- [27] G. Barton, Introduction to Dispersion Techniques in Field Theory, W.A. Benjamin Inc., Amsterdam, 1965.
- [28] F.-K. Guo, et al., Phys. Lett. B 678 (2009) 90.
- [29] D.J. Antonio, et al., PoS LAT2006 (2006) 101, arXiv:hep-lat/0610080.
- [30] D. Becirevic, et al., Nucl. Phys. B 705 (2005) 339.
- [31] I.M. Dremin, Zh. Eksp. Teor. Fiz. 41 (1961) 821.
- [32] B.V. Geshkenbein, B.L. Ioffe, Pis'ma Zh. Eksp. Teor. Fiz. 18 (1973) 646, JETP Lett. 70 (1999) 161.
- [33] V. Bernard, U.-G. Meißner, A. Rusetsky, Nucl. Phys. B 788 (2008) 1.
- [34] V. Bernard, D. Hoja, U.-G. Meißner, A. Rusetsky, JHEP 0906 (2009) 061.
- [35] T. Ledwig, et al., arXiv:1104.3877 [hep-ph];  
T. Ledwig, V. Pascalutsa, M. Vanderhaeghen, Phys. Rev. D 82 (2010) 091301.
- [36] A. Semke, M.F.M. Lutz, Nucl. Phys. A 789 (2007) 251.
- [37] J.R. Pelaez, G. Rios, Phys. Rev. D 82 (2010) 114002.