

Matrix biorthogonal polynomials on the real line: Geronimus transformations

Gerardo Ariznabarreta^{*,§}, Juan C. García-Ardila^{†,¶}, Manuel Mañas^{*,||} and
Francisco Marcellán^{†,‡,**}

**Departamento de Física Teórica II (Métodos Matemáticos de la Física)
Universidad Complutense de Madrid, Ciudad Universitaria
Plaza de Ciencias 1, 28040 Madrid, Spain*

*†Departamento de Matemáticas, Universidad Carlos III de Madrid
Avd/Universidad 30, 28911 Leganés, Spain*

*‡Instituto de Ciencias Matemáticas (ICMAT)
C/Nicolás Cabrera 13-15, 28049 Canto Blanco, Spain*

§gariznab@ucm.es

¶jugarcia@math.uc3m.es

||manuel.manas@ucm.es

***pacomarc@ing.uc3m.es*

Received 6 February 2017

Accepted 26 July 2018

Published 8 May 2019

Communicated by Ari Laptev

In this paper, Geronimus transformations for matrix orthogonal polynomials in the real line are studied. The orthogonality is understood in a broad sense, and is given in terms of a nondegenerate continuous sesquilinear form, which in turn is determined by a quasi-definite matrix of bivariate generalized functions with a well-defined support. The discussion of the orthogonality for such a sesquilinear form includes, among others, matrix Hankel cases with linear functionals, general matrix Sobolev orthogonality and discrete orthogonal polynomials with an infinite support. The results are mainly concerned with the derivation of Christoffel-type formulas, which allow to express the perturbed matrix biorthogonal polynomials and its norms in terms of the original ones. The basic tool is the Gauss–Borel factorization of the Gram matrix, and particular attention is paid to the non-associative character, in general, of the product of semi-infinite matrices. The Geronimus transformation in which a right multiplication by the inverse of a matrix polynomial and an addition of adequate masses are performed, is considered. The resolvent matrix and connection formulas are given. Two different methods are developed. A spectral one, based on the spectral properties of the perturbing polynomial, and constructed in terms of the second kind functions. This approach requires the perturbing matrix polynomial to have a nonsingular leading term. Then, using spectral techniques and spectral jets, Christoffel–Geronimus formulas for the transformed polynomials and

**Corresponding author.

This is an Open Access article published by World Scientific Publishing Company. It is distributed under the terms of the Creative Commons Attribution 4.0 (CC-BY) License. Further distribution of this work is permitted, provided the original work is properly cited.

norms are presented. For this type of transformations, the paper also proposes an alternative method, which does not require of spectral techniques, that is valid also for singular leading coefficients. When the leading term is nonsingular, a comparison of between both methods is presented. The nonspectral method is applied to unimodular Christoffel perturbations, and a simple example for a degree one massless Geronimus perturbation is given.

Keywords: Matrix biorthogonal polynomials; spectral theory of matrix polynomials; quasi-definite matrix of generalized kernels; nondegenerate continuous sesquilinear forms; Gauss–Borel factorization; matrix Geronimus transformations; matrix linear spectral transformations; Christoffel-type formulas; quasideterminants; spectral jets; unimodular matrix polynomials.

Mathematics Subject Classification: 42C05, 15A23

1. Introduction

Perturbations of a linear functional u in the linear space of polynomials with real coefficients have been extensively studied in the theory of orthogonal polynomials on the real line (scalar OPRL). In particular, when you deal with the positive definite case and linear functionals associated with probability measures supported in an infinite subset of the real line are considered, such perturbations provide interesting information in the framework of Gaussian quadrature rules taking into account the perturbation yields new nodes and Christoffel numbers, see [25, 26]. Three perturbations have attracted the interest of the researchers. Christoffel perturbations, that appear when you consider a new functional $\hat{u} = p(x)u$, where $p(x)$ is a polynomial, were studied in 1858 by the German mathematician Christoffel in [13], in the framework of Gaussian quadrature rules. He found explicit formulas relating the corresponding sequences of orthogonal polynomials with respect to two measures, the Lebesgue measure $d\mu$ supported in the interval $(-1, 1)$ and $d\hat{\mu}(x) = p(x)d\mu(x)$, with $p(x) = (x - q_1) \cdots (x - q_N)$ a signed polynomial in the support of $d\mu$, as well as the distribution of their zeros as nodes in such quadrature rules. Nowadays, these are called Christoffel formulas, and can be considered a classical result in the theory of orthogonal polynomials which can be found in a number of textbooks, see for example [12, 26, 71]. Explicit relations between the corresponding sequences of orthogonal polynomials have been extensively studied, see [25], as well as the connection between the corresponding monic Jacobi matrices in the framework of the so-called Darboux transformations based on the LU factorization of such matrices [9]. In the theory of orthogonal polynomials, connection formulas between two families of orthogonal polynomials allow to express any polynomial of a given degree n as a linear combination of all polynomials of degree less than or equal to n in the second family. A noteworthy fact regarding the Christoffel finding is that in some cases the number of terms does not grow with the degree n but remarkably, and on the contrary, remain constant, equal to the degree of the perturbing polynomial. See [25, 26] for more on the Christoffel-type formulas. Christoffel transformations for orthogonal polynomials on the unit circle based on LU factorizations of CMV matrices have been studied in [10].

Geronimus transformation appears when you are dealing with perturbed functionals v defined by $p(x)v = u$, where $p(x)$ is a polynomial. Such a kind of transformations were used by the Russian mathematician Geronimus, see [35], in order to have a nice proof of a result by Hahn [43] concerning the characterization of classical orthogonal polynomials (Hermite, Laguerre, Jacobi, and Bessel) as those orthogonal polynomials whose first derivatives are also orthogonal polynomials, for an English account of Geronimus' paper [35] see [40]. Again, as happened for the Christoffel transformation, within the Geronimus transformation one can find Christoffel-type formulas, now in terms of the second kind functions, relating the corresponding sequences of orthogonal polynomials, see for example the work of Maroni [55] for a perturbation of the type $p(x) = x - a$.

Krein in [48] was the first to discuss matrix orthogonal polynomials, for a review on the subject see [15]. The great activity in this scientific field has produced a vast bibliography, treating among other things subjects like inner products defined on the linear space of polynomials with matrix coefficients or aspects as the existence of the corresponding sequences of matrix orthogonal polynomials in the real line, see [18, 19, 56, 63, 70]) and their applications in Gaussian quadrature for matrix-valued functions [69], scattering theory [5, 34] and system theory [24]. The seminal paper [20] gave the key for further studies in this subject and, subsequently, some relevant advances have been achieved in the study of families of matrix orthogonal polynomials associated to second-order linear differential operators as eigenfunctions and their structural properties [18, 21, 41, 42]. In [11], sequences of orthogonal polynomials satisfying a first-order linear matrix differential equation were found, which is a remarkable difference with the scalar scenario, where such a situation does not appear. The spectral problem for second-order linear difference operators with polynomial coefficients has been considered in [4]. Therein four families of matrix orthogonal polynomials (as matrix relatives of Charlier, Meixner, Krawtchouk scalar polynomials and another one that seems not have any scalar relative) are obtained as illustrative examples of the method described therein.

We continue this introduction with two introductory subsections. One is focused on the spectral theory of matrix polynomials, we follow [39]. The other is a basic background on matrix orthogonal polynomials, see [15]. In the Sec. 2, we extend the Geronimus transformations to the matrix realm, and find connection formulas for the biorthogonal polynomials and the Christoffel–Darboux kernels. These developments allow for the finding of the Christoffel–Geronimus formula for matrix perturbations of Geronimus type. As we said we present two different schemes. In the first one, which can be applied when the perturbing polynomial has a nonsingular leading coefficient, we express the perturbed objects in terms of spectral jets of the primitive second kind functions and Christoffel–Darboux kernels. We present a second approach, applicable even when the leading coefficient is singular. For each method we consider two different situations, the less interesting case of biorthogonal polynomials of degree less than the degree of the perturbing polynomial, and the much more interesting situation whence the degrees of the families of biorthogonal

polynomials are greater than or equal to the degree of the perturbing polynomial. To end the section, we compare spectral versus nonspectral methods and present a number of applications. In particular, we deal with unimodular polynomial matrix perturbations and degree one matrix Geronimus transformations. Notice that in [6] we have extended these results to the matrix linear spectral case, i.e. to Uvarov–Geronimus–Christoffel formulas for certain matrix rational perturbations. Finally, an appendix with the definitions of Schur complements and quasideterminants is also included in order to have a perspective of these basic tools in the theory of matrix orthogonal polynomials.

1.1. On spectral theory of matrix polynomials

Here we give some background material regarding the spectral theory of matrix polynomials [39, 52].

Definition 1. Let $A_0, A_1, \dots, A_N \in \mathbb{C}^{p \times p}$ be square matrices of size $p \times p$ with complex entries and $A_N \neq 0_p$. Then

$$W(x) = A_N x^N + A_{N-1} x^{N-1} + \dots + A_1 x + A_0 \tag{1}$$

is said to be a matrix polynomial of degree N , $\deg(W(x)) = N$. The matrix polynomial is said to be monic when $A_N = I_p$, where $I_p \in \mathbb{C}^{p \times p}$ denotes the identity matrix. The linear space — a bimodule for the ring of matrices $\mathbb{C}^{p \times p}$ — of matrix polynomials with coefficients in $\mathbb{C}^{p \times p}$ will be denoted by $\mathbb{C}^{p \times p}[x]$.

Definition 2 (Eigenvalues). The spectrum, or the set of eigenvalues, $\sigma(W(x))$ of a matrix polynomial W is the zero set of $\det W(x)$, i.e.

$$\sigma(W(x)) := \{x \in \mathbb{C} : \det W(x) = 0\}.$$

Proposition 1. A monic matrix polynomial $W(x)$, $\deg(W(x)) = N$, has Np (counting multiplicities) eigenvalues or zeros, i.e. we can write

$$\det W(x) = \prod_{a=1}^q (x - x_a)^{\alpha_a},$$

with $Np = \alpha_1 + \dots + \alpha_q$.

Proposition 2. Any nonsingular matrix polynomial $W(x) \in \mathbb{C}^{p \times p}[x]$, $\det W(x) \neq 0$, can be represented as

$$W(x) = E_{x_0}(x) \operatorname{diag}((x - x_0)^{\kappa_1}, \dots, (x - x_0)^{\kappa_m}) F_{x_0}(x)$$

at $x = x_0 \in \mathbb{C}$, where $E_{x_0}(x)$ and $F_{x_0}(x)$ are nonsingular matrices and $\kappa_1 \leq \dots \leq \kappa_m$ are nonnegative integers. Moreover, $\{\kappa_1, \dots, \kappa_m\}$ are uniquely determined by $W(x)$ and they are known as partial multiplicities of $W(x)$ at x_0 .

Definition 3. For an eigenvalue x_0 of a monic matrix polynomial $W(x) \in \mathbb{C}^{p \times p}[x]$, then:

- (i) A nonzero vector $r_0 \in \mathbb{C}^p$ is said to be a right eigenvector, with eigenvalue $x_0 \in \sigma(W(x))$, whenever $W(x_0)r_0 = 0$, i.e. $r_0 \in \operatorname{Ker} W(x_0) \neq \{0\}$.

- (ii) A nonzero covector $l_0 \in (\mathbb{C}^p)^*$ is said to be a left eigenvector, with eigenvalue $x_0 \in \sigma(W(x))$, whenever $l_0 W(x_0) = 0, (l_0)^\top \in (\text{Ker}(W(x_0)))^\perp = \text{Ker}((W(x_0))^\top) \neq \{0\}$.
- (iii) A sequence of vectors $\{r_0, r_1, \dots, r_{m-1}\}$ is said to be a right Jordan chain of length m corresponding to the eigenvalue $x_0 \in \sigma(W(x))$, if r_0 is a right eigenvector of $W(x_0)$ and

$$\sum_{s=0}^j \frac{1}{s!} \frac{d^s W}{dx^s} \Big|_{x=x_0} r_{j-s} = 0, \quad j \in \{0, \dots, m-1\}.$$

- (iv) A sequence of covectors $\{l_0, l_1, \dots, l_{m-1}\}$ is said to be a left Jordan chain of length m , corresponding to $x_0 \in \sigma(W^\top)$, if $\{(l_0)^\top, (l_1)^\top, \dots, (l_{m-1})^\top\}$ is a right Jordan chain of length m for the matrix polynomial $(W(x))^\top$.
- (v) A right root polynomial at x_0 is a nonzero vector polynomial $r(x) \in \mathbb{C}^p[x]$ such that $W(x)r(x)$ has a zero of certain order at $x = x_0$, the order of this zero is called the order of the root polynomial. Analogously, a left root polynomial is a nonzero covector polynomial $l(x) \in (\mathbb{C}^p)^*[x]$ such that $l(x_0)W(x_0) = 0$.
- (vi) The maximal lengths, either of right or left Jordan chains corresponding to the eigenvalue x_0 , are called the multiplicity of the eigenvector r_0 or l_0 and are denoted by $m(r_0)$ or $m(l_0)$, respectively.

Proposition 3. *Given an eigenvalue $x_0 \in \sigma(W(x))$ of a monic matrix polynomial $W(x)$, multiplicities of right and left eigenvectors coincide and they are equal to the corresponding partial multiplicities κ_i .*

The above definition generalizes the concept of Jordan chain for degree one matrix polynomials.

Proposition 4. *The Taylor expansion of a right root polynomial $r(x)$, respectively of a left root polynomial $l(x)$, at a given eigenvalue $x_0 \in \sigma(W(x))$ of a monic matrix polynomial $W(x)$,*

$$r(x) = \sum_{j=0}^{\kappa-1} r_j(x-x_0)^j, \quad \text{respectively, } l(x) = \sum_{j=0}^{\kappa-1} l_j(x-x_0)^j,$$

provides us with right Jordan chain

$$\{r_0, r_1, \dots, r_{\kappa-1}\}, \quad \text{respectively, left Jordan chain } \{l_0, l_1, \dots, l_{\kappa-1}\}.$$

Proposition 5. *Given an eigenvalue $x_0 \in \sigma(W(x))$ of a monic matrix polynomial $W(x)$, with multiplicity $s = \dim \text{Ker } W(x_0)$, we can construct s right root polynomials, respectively left root polynomials, for $i \in \{1, \dots, s\}$,*

$$r_i(x) = \sum_{j=0}^{\kappa_i-1} r_{i,j}(x-x_0)^j, \quad \text{respectively, } l_i(x) = \sum_{j=0}^{\kappa_i-1} l_{i,j}(x-x_0)^j,$$

where $r_i(x)$ are right root polynomials (respectively, $l_i(x)$ are left root polynomials) with the largest order κ_i among all right root polynomials, whose right eigenvector

does not belong to $\mathbb{C}\{r_{0,1}, \dots, r_{0,i-1}\}$ (respectively, left root polynomials whose left eigenvector does not belong to $\mathbb{C}\{l_{0,1}, \dots, l_{0,i-1}\}$).

Definition 4 (Canonical Jordan chains). A canonical set of right Jordan chains (respectively left Jordan chains) of the monic matrix polynomial $W(x)$ corresponding to the eigenvalue $x_0 \in \sigma(W(x))$ is, in terms of the right root polynomials (respectively left root polynomials) described in Proposition 5, the following sets of vectors

$$\{r_{1,0}, \dots, r_{1,\kappa_1-1}, \dots, r_{s,0}, \dots, r_{s,\kappa_s-1}\}, \quad \text{respectively, covectors}$$

$$\{l_{1,0}, \dots, l_{1,\kappa_1-1}, \dots, l_{s,0}, \dots, l_{s,\kappa_s-1}\}.$$

Proposition 6. For a monic matrix polynomial $W(x)$ the lengths $\{\kappa_1, \dots, \kappa_s\}$ of the Jordan chains in a canonical set of Jordan chains of $W(x)$ corresponding to the eigenvalue x_0 , see Definition 4, are the nonzero partial multiplicities of $W(x)$ at $x = x_0$ described in Proposition 2.

Definition 5 (Canonical Jordan chains and root polynomials). For each eigenvalue $x_a \in \sigma(W(x))$ of a monic matrix polynomial $W(x)$, with multiplicity α_a and $s_a = \dim \text{Ker } W(x_a)$, $a \in \{1, \dots, q\}$, we choose a canonical set of right Jordan chains, respectively left Jordan chains,

$$\left\{ r_{j,0}^{(a)}, \dots, r_{j,\kappa_j^{(a)}-1}^{(a)} \right\}_{j=1}^{s_a}, \quad \text{respectively} \quad \left\{ l_{j,0}^{(a)}, \dots, l_{j,\kappa_j^{(a)}-1}^{(a)} \right\}_{j=1}^{s_a},$$

and, consequently, with partial multiplicities satisfying $\sum_{j=1}^{s_a} \kappa_j^{(a)} = \alpha_a$. Thus, we can consider the following right root polynomials:

$$r_j^{(a)}(x) = \sum_{l=0}^{\kappa_j^{(a)}-1} r_{j,l}^{(a)}(x-x_a)^l, \quad \text{respectively, left root polynomials}$$

$$l_j^{(a)}(x) = \sum_{l=0}^{\kappa_j^{(a)}-1} l_{j,l}^{(a)}(x-x_a)^l. \tag{2}$$

Definition 6 (Canonical Jordan pairs). We also define the corresponding canonical Jordan pair (X_a, J_a) with X_a the matrix

$$X_a := \left[r_{1,0}^{(a)}, \dots, r_{1,\kappa_1^{(a)}-1}^{(a)}, \dots, r_{s_a,0}^{(a)}, \dots, r_{s_a,\kappa_{s_a}^{(a)}-1}^{(a)} \right] \in \mathbb{C}^{p \times \alpha_a},$$

and J_a the matrix

$$J_a := \text{diag}(J_{a,1}, \dots, J_{a,s_a}) \in \mathbb{C}^{\alpha_a \times \alpha_a},$$

where $J_{a,j} \in \mathbb{C}^{\kappa_j^{(a)} \times \kappa_j^{(a)}}$ are the Jordan blocks of the eigenvalue $x_a \in \sigma(W(x))$. Then, we say that (X, J) with

$$X := [X_1, \dots, X_q] \in \mathbb{C}^{p \times Np}, \quad J := \text{diag}(J_1, \dots, J_q) \in \mathbb{C}^{Np \times Np},$$

is a canonical Jordan pair for $W(x)$.

We have the important result, see [39],

Proposition 7. *The Jordan pairs of a monic matrix polynomial $W(x)$ satisfy*

$$A_0 X_a + A_1 X_a J_a + \cdots + A_{N-1} X_a (J_a)^{N-1} + X_a (J_a)^N = 0_{p \times \alpha_a},$$

$$A_0 X + A_1 X J + \cdots + A_{N-1} X J^{N-1} + X J^N = 0_{p \times Np}.$$

A key property, see [39, Theorem 1.20] is the following.

Proposition 8. *For any Jordan pair (X, J) of a monic matrix polynomial $W(x) = I_p x^N + A_{N-1} x^{N-1} + \cdots + A_0$ the matrix*

$$\begin{bmatrix} X \\ XJ \\ \vdots \\ XJ^{N-1} \end{bmatrix} \in \mathbb{C}^{Np \times Np}$$

is nonsingular.

Definition 7 (Jordan triple). Given

$$Y = \begin{bmatrix} Y_1 \\ \vdots \\ Y_q \end{bmatrix} \in \mathbb{C}^{Np \times p},$$

with $Y_a \in \mathbb{C}^{\alpha_a \times p}$, we say that (X, J, Y) is a Jordan triple whenever

$$\begin{bmatrix} X \\ XJ \\ \vdots \\ XJ^{N-1} \end{bmatrix} Y = \begin{bmatrix} 0_p \\ \vdots \\ 0_p \\ I_p \end{bmatrix}.$$

Moreover, [39, Theorem 1.23], gives the following characterization.

Proposition 9. *Two matrices $X \in \mathbb{C}^{p \times Np}$ and $J \in \mathbb{C}^{Np \times Np}$ constitute a Jordan pair of a monic matrix polynomial $W(x) = I_p x^N + A_{N-1} x^{N-1} + \cdots + A_0$ if and only if the two following properties hold:*

(i) *The matrix*

$$\begin{bmatrix} X \\ XJ \\ \vdots \\ XJ^{N-1} \end{bmatrix}$$

is nonsingular.

(ii)

$$A_0X + A_1XJ + \dots + A_{N-1}XJ^{N-1} + XJ^N = 0_{p \times Np}.$$

Proposition 10. *Given a monic matrix polynomial $W(x)$ the adapted root polynomials given in Definition 5 satisfy*

$$\begin{aligned} (W(x)r_j^{(a)}(x))_{x_a}^{(m)} = 0, \quad (l_j^{(a)}(x)W(x))_{x_a}^{(m)} = 0, \\ m \in \{0, \dots, \kappa_j^{(a)} - 1\}, \quad j \in \{1, \dots, s_a\}. \end{aligned} \tag{3}$$

Here, given a function $f(x)$ we use the following notation for its derivatives evaluated at an eigenvalue $x_a \in \sigma(W(x))$

$$(f)_{x_a}^{(m)} := \lim_{x \rightarrow x_a} \frac{d^m f}{dx^m}.$$

In this paper, we assume that the partial multiplicities are ordered in an increasing way, i.e. $\kappa_1^{(a)} \leq \kappa_2^{(a)} \leq \dots \leq \kappa_{s_a}^{(a)}$.

Proposition 11. *If $r_i^{(a)}$ and $l_j^{(a)}$ are right and left root polynomials corresponding to the eigenvalue $x_a \in \sigma(W(x))$, then a polynomial*

$$w_{i,j}^{(a)}(x) = \sum_{m=0}^{d_{i,j}^{(a)}} w_{i,j;m}^{(a)} x^m \in \mathbb{C}[x], \quad d_{i,j}^{(a)} := \kappa_{\min(i,j)}^{(a)} + N - 2,$$

exists such that

$$l_i^{(a)}(x)W(x)r_j^{(a)}(x) = (x - x_a)^{\kappa_{\max(i,j)}^{(a)}} w_{i,j}^{(a)}(x). \tag{4}$$

Definition 8 (Spectral jets). Given a matrix function $f(x)$ smooth in region $\Omega \subset \mathbb{C}$ with $x_a \in \overline{\Omega}$, a point in the closure of Ω we consider its matrix spectral jets

$$\mathcal{J}_f^{(i)}(x_a) := \lim_{x \rightarrow x_a} \left[f(x), \dots, \frac{f^{(\kappa_i^{(a)}-1)}(x)}{(\kappa_i^{(a)}-1)!} \right] \in \mathbb{C}^{p \times p \kappa_i^{(a)}},$$

$$\mathcal{J}_f(x_a) := [\mathcal{J}_f^{(1)}(x_a), \dots, \mathcal{J}_f^{(s_a)}(x_a)] \in \mathbb{C}^{p \times p \alpha_a},$$

$$\mathcal{J}_f := [\mathcal{J}_f(x_1), \dots, \mathcal{J}_f(x_q)] \in \mathbb{C}^{p \times Np^2},$$

and given a Jordan pair the root spectral jet vectors

$$\mathcal{J}_f^{(i)}(x_a) := \lim_{x \rightarrow x_a} \left[f(x)r_i^{(a)}(x), \dots, \frac{(f(x)r_i^{(a)}(x))_{x_a}^{(\kappa_i^{(a)}-1)}}{(\kappa_i^{(a)}-1)!} \right] \in \mathbb{C}^{p \times \kappa_i^{(a)}},$$

$$\mathcal{J}_f(x_a) := [\mathcal{J}_f^{(1)}(x_a), \dots, \mathcal{J}_f^{(s_a)}(x_a)] \in \mathbb{C}^{p \times \alpha_a},$$

$$\mathcal{J}_f := [\mathcal{J}_f(x_1), \dots, \mathcal{J}_f(x_q)] \in \mathbb{C}^{p \times Np}.$$

Definition 9. We consider the following jet matrices

$$\begin{aligned} \mathcal{Q}_{n;i}^{(a)} &:= \mathcal{J}_{I_p x^n}^{(i)}(x_a) = \left[\begin{array}{c} (x_a)^n r_i^{(a)}(x_a), (x^n r_i^{(a)}(x))_{x_a}^{(1)}, \dots, \\ \dots \frac{(x^n r_i^{(a)}(x))_{x_a}^{(\kappa_i^{(a)} - 1)}}{(\kappa_i^{(a)} - 1)!} \end{array} \right] \in \mathbb{C}^{p \times \kappa_i^{(a)}}, \\ \mathcal{Q}_n^{(a)} &:= \mathcal{J}_{I_p x^n}(x_a) = [\mathcal{Q}_{n;1}^{(a)}, \dots, \mathcal{Q}_{n;s_a}^{(a)}] \in \mathbb{C}^{p \times \alpha_a}, \\ \mathcal{Q}_n &:= \mathcal{J}_{I_p x^n} = [\mathcal{Q}_n^{(1)}, \dots, \mathcal{Q}_n^{(q)}] \in \mathbb{C}^{p \times Np}, \\ \mathcal{Q} &:= \mathcal{J}_{\chi_{[N]}} = \begin{bmatrix} \mathcal{Q}_0 \\ \vdots \\ \mathcal{Q}_{N-1} \end{bmatrix} \in \mathbb{C}^{Np \times Np}, \end{aligned}$$

where $(\chi_{[N]}(x))^\top := [I_p, \dots, I_p x^{N-1}] \in \mathbb{C}^{p \times Np}[x]$.

Lemma 1 (Root spectral jets and Jordan pairs). *Given a canonical Jordan pair (X, J) for the monic matrix polynomial $W(x)$ we have that*

$$\mathcal{Q}_n = XJ^n, \quad n \in \{0, 1, \dots\}.$$

Thus, any polynomial $P_n(x) = \sum_{j=0}^n P_j x^j$ has as its spectral jet vector corresponding to $W(x)$ the following matrix.

$$\mathcal{J}_P = P_0 X + P_1 XJ + \dots + P_n XJ^{n-1}.$$

Definition 10. If $W(x) = \sum_{k=0}^N A_k x^k \in \mathbb{C}^{p \times p}[x]$ is a matrix polynomial of degree N , we introduce the matrix

$$\mathcal{B} := \begin{bmatrix} A_1 & A_2 & A_3 & \dots & A_{N-1} & A_N \\ A_2 & A_3 & \vdots & \ddots & A_N & 0_p \\ A_3 & \dots & A_{N-1} & \ddots & 0_p & 0_p \\ \vdots & \ddots & \ddots & \ddots & & \vdots \\ A_{N-1} & A_N & 0_p & & & \\ A_N & 0_p & 0_p & \dots & 0_p & \end{bmatrix} \in \mathbb{C}^{Np \times Np}.$$

Lemma 2. *Given a Jordan triple (X, J, Y) for the monic matrix polynomial $W(x)$ we have*

$$\mathcal{Q} = \begin{bmatrix} X \\ XJ \\ \vdots \\ XJ^{N-1} \end{bmatrix}, \quad (\mathcal{B}\mathcal{Q})^{-1} = [Y, JY, \dots, J^{N-1}Y] =: \mathcal{R}.$$

Proof. From Lemma 1, we deduce that

$$Q = \begin{bmatrix} X \\ XJ \\ \vdots \\ XJ^{N-1} \end{bmatrix}$$

which is nonsingular, see Propositions 8 and 9. The biorthogonality condition (2.6) of [39] for \mathcal{R} and Q is

$$\mathcal{R}BQ = I_{Np},$$

and if (X, J, Y) is a canonical Jordan triple, then

$$\mathcal{R} = [Y, JY, \dots, J^{N-1}Y]. \tag{5}$$

□

Proposition 12. *The matrix $\mathcal{R}_n := [Y, JY, \dots, J^{n-1}Y] \in \mathbb{C}^{Np \times np}$ has full rank.*

Regarding the matrix \mathcal{B} , we have the following.

Definition 11. Let us consider the bivariate matrix polynomial

$$\mathcal{V}(x, y) := ((\chi(y))_{[N]})^\top \mathcal{B}(\chi(x))_{[N]} \in \mathbb{C}^{p \times p}[x, y],$$

where A_j are the matrix coefficients of $W(x)$, see (1).

We consider the complete homogeneous symmetric polynomials in two variables

$$h_n(x, y) = \sum_{j=0}^n x^j y^{n-j}.$$

For example, the first four polynomials are

$$\begin{aligned} h_0(x, y) &= 1, & h_1(x, y) &= x + y, & h_2(x, y) &= x^2 + xy + y^2, \\ h_3(x, y) &= x^3 + x^2y + xy^2 + y^3. \end{aligned}$$

Proposition 13. *In terms of complete homogeneous symmetric polynomials in two variables we can write*

$$\mathcal{V}(x, y) = \sum_{j=1}^N A_j h_{j-1}(x, y).$$

1.2. On orthogonal matrix polynomials

The polynomial ring $\mathbb{C}^{p \times p}[x]$ is a free bimodule over the ring of matrices $\mathbb{C}^{p \times p}$ with a basis given by $\{I_p, I_p x, I_p x^2, \dots\}$. Important free bisubmodules are the sets $\mathbb{C}_m^{p \times p}[x]$ of matrix polynomials of degree less than or equal to m . A basis, which has

cardinality $m + 1$, for $\mathbb{C}^{p \times p}[x]$ is $\{I_p, I_p x, \dots, I_p x^m\}$; as \mathbb{C} has the invariant basis number (IBN) property so does $\mathbb{C}^{p \times p}$, see [64]. Therefore, being $\mathbb{C}^{p \times p}$ an IBN ring, the rank of the free module $\mathbb{C}^{p \times p}[x]$ is unique and equal to $m + 1$, i.e. any other basis has the same cardinality. Its algebraic dual $(\mathbb{C}^{p \times p}[x])^*$ is the set of homomorphisms $\phi : \mathbb{C}^{p \times p}[x] \rightarrow \mathbb{C}^{p \times p}$ which are, for the right module, of the form

$$\langle \phi, P(x) \rangle = \phi_0 p_0 + \dots + \phi_m p_m, \quad P(x) = p_0 + \dots + p_m x^m,$$

where $\phi_k \in \mathbb{C}^{p \times p}$. Thus, we can identify the dual of the right module with the corresponding left submodule. This dual is a free module with a unique rank, equal to $m + 1$, and a dual basis $\{(I_p x^k)^*\}_{k=0}^m$ given by

$$\langle (I_p x^k)^*, I_p x^l \rangle = \delta_{k,l} I_p.$$

We have similar statements for the left module $\mathbb{C}^{p \times p}[x]$, being its dual a right module

$$\langle P(x), \phi \rangle = P_0 \phi_0 + \dots + P_m \phi_m, \quad \langle I_p x^l, (I_p x^k)^* \rangle = \delta_{k,l} I_p.$$

Definition 12 (Sesquilinear form). A sesquilinear form $\langle \cdot, \cdot \rangle$ on the bimodule $\mathbb{C}^{p \times p}[x]$ is a continuous map

$$\langle \cdot, \cdot \rangle : \mathbb{C}^{p \times p}[x] \times \mathbb{C}^{p \times p}[x] \rightarrow \mathbb{C}^{p \times p},$$

$$(P(x), Q(x)) \mapsto \langle P(x), Q(x) \rangle,$$

such that for any triple $P(x), Q(x), R(x) \in \mathbb{C}^{p \times p}[x]$ the following properties are fulfilled:

- (i) $\langle AP(x) + BQ(x), R(y) \rangle = A \langle P(x), R(y) \rangle + B \langle Q(x), R(y) \rangle, \forall A, B \in \mathbb{C}^{p \times p},$
- (ii) $\langle P(x), AQ(y) + BR(y) \rangle = \langle P(x), Q(y) \rangle A^\top + \langle P(x), R(y) \rangle B^\top, \forall A, B \in \mathbb{C}^{p \times p}.$

The reader probably has noticed that, despite dealing with complex polynomials in a real variable, we have followed [26] and chosen the transpose instead of the Hermitian conjugated. For any couple of matrix polynomials $P(x) = \sum_{k=0}^{\deg P} p_k x^k$ and $Q(x) = \sum_{l=0}^{\deg Q} q_l x^l$ the sesquilinear form is defined by

$$\langle P(x), Q(y) \rangle = \sum_{\substack{k=1, \dots, \deg P \\ l=1, \dots, \deg Q}} p_k G_{k,l} (q_l)^\top,$$

where the coefficients are the values of the sesquilinear form on the basis of the module

$$G_{k,l} = \langle I_p x^k, I_p y^l \rangle.$$

The corresponding semi-infinite matrix

$$G = \begin{bmatrix} G_{0,0} & G_{0,1} & \dots \\ G_{1,0} & G_{1,1} & \dots \\ \vdots & \vdots & \ddots \end{bmatrix}$$

is named as the Gram matrix of the sesquilinear form.

1.2.1. *Hankel sesquilinear forms*

Now, we present a family of examples of sesquilinear forms in $\mathbb{C}^{p \times p}[x]$ that we call Hankel sesquilinear forms. A first example is given by matrices with complex (or real) Borel measures in \mathbb{R} as entries

$$\mu = \begin{bmatrix} \mu_{1,1} & \cdots & \mu_{1,p} \\ \vdots & & \vdots \\ \mu_{p,1} & \cdots & \mu_{p,p} \end{bmatrix},$$

i.e. a $p \times p$ matrix of Borel measures supported in \mathbb{R} . Given any pair of matrix polynomials $P(x), Q(x) \in \mathbb{C}^{p \times p}[x]$ we introduce the following sesquilinear form:

$$\langle P(x), Q(x) \rangle_\mu = \int_{\mathbb{R}} P(x) d\mu(x) (Q(x))^\top.$$

A more general sesquilinear form can be constructed in terms of generalized functions (or continuous linear functionals). In [53, 54], a linear functional setting for orthogonal polynomials is given. We consider the space of polynomials $\mathbb{C}[x]$, with an appropriate topology, as the space of fundamental functions, in the sense of [32, 33], and take the space of generalized functions as the corresponding continuous linear functionals. It is remarkable that the topological dual space coincides with the algebraic dual space. On the other hand, this space of generalized functions is the space of formal series with complex coefficients $(\mathbb{C}[x])' = \mathbb{C}[[x]]$.

In this paper, we use generalized functions with a well-defined support and, consequently, the previously described setting requires a suitable modification. Following [32, 33, 67], let us recall that the space of distributions is a space of generalized functions when the space of fundamental functions is constituted by the complex-valued smooth functions of compact support $\mathcal{D} := C_0^\infty(\mathbb{R})$, the so-called space of test functions. In this context, the set of zeros of a distribution $u \in \mathcal{D}'$ is the region $\Omega \subset \mathbb{R}$ if for any fundamental function $f(x)$ with support in Ω we have $\langle u, f \rangle = 0$. Its complement, a closed set, is what is called support, $\text{supp } u$, of the distribution u . Distributions of compact support, $u \in \mathcal{E}'$, are generalized functions for which the space of fundamental functions is the topological space of complex-valued smooth functions $\mathcal{E} = C^\infty(\mathbb{R})$. As $\mathbb{C}[x] \subsetneq \mathcal{E}$ we also know that $\mathcal{E}' \subsetneq (\mathbb{C}[x])' \cap \mathcal{D}'$. The set of distributions of compact support is a first example of an appropriate framework for the consideration of polynomials and supports simultaneously. More general settings appear within the space of tempered distributions \mathcal{S}' , $\mathcal{S}' \subsetneq \mathcal{D}'$. The space of fundamental functions is given by the Schwartz space \mathcal{S} of complex-valued fast decreasing functions, see [32, 33, 67]. We consider the space of fundamental functions constituted by smooth functions of slow growth $\mathcal{O}_M \subset \mathcal{E}$, whose elements are smooth functions with derivatives bounded by polynomials. As $\mathbb{C}[x], \mathcal{S} \subsetneq \mathcal{O}_M$, for the corresponding set of generalized functions we find that $\mathcal{O}'_M \subset (\mathbb{C}[x])' \cap \mathcal{S}'$. Therefore, these distributions give a second appropriate framework. Finally, for

a third suitable framework, including the two previous ones, we need to introduce bounded distributions. Let us consider as space of fundamental functions, the linear space \mathcal{B} of bounded smooth functions, i.e. with all its derivatives in $L^\infty(\mathbb{R})$, being the corresponding space of generalized functions \mathcal{B}' the bounded distributions. From $\mathcal{D} \subsetneq \mathcal{B}$ we conclude that bounded distributions are distributions $\mathcal{B}' \subsetneq \mathcal{D}'$. Then, we consider the space of fast decreasing distributions \mathcal{O}'_c given by those distributions $u \in \mathcal{D}'$ such that for each positive integer k , we have that $(\sqrt{1+x^2})^k u \in \mathcal{B}'$ is a bounded distribution. Any polynomial $P(x) \in \mathbb{C}[x]$, with $\deg P = k$, can be written as $P(x) = (\sqrt{1+x^2})^k F(x)$ and $F(x) = \frac{P(x)}{(\sqrt{1+x^2})^k} \in \mathcal{B}$. Therefore, given a fast decreasing distribution $u \in \mathcal{O}'_c$ we may consider

$$\langle u, P(x) \rangle = \langle (\sqrt{1+x^2})^k u, F(x) \rangle$$

which makes sense as $(\sqrt{1+x^2})^k u \in \mathcal{B}'$, $F(x) \in \mathcal{B}$. Thus, $\mathcal{O}'_c \subset (\mathbb{C}[x])' \cap \mathcal{D}'$. Moreover, it can be proven that $\mathcal{O}'_M \subsetneq \mathcal{O}'_c$, see [53]. Summarizing this discussion, we have found three generalized function spaces suitable for the discussion of polynomials and supports simultaneously: $\mathcal{E}' \subset \mathcal{O}'_M \subset \mathcal{O}'_c \subset ((\mathbb{C}[x])' \cap \mathcal{D}')$.

The linear functionals could have discrete and, as the corresponding Gram matrix is required to be quasi-definite, infinite support. Then, we are faced with discrete orthogonal polynomials, see for example [57]. Two classical examples are those of Charlier and Meixner. For $\mu > 0$ we have the Charlier (or Poisson–Charlier) linear functional

$$u = \sum_{k=0}^{\infty} \frac{\mu^k}{k!} \delta(x - k),$$

and $\beta > 0$ and $0 < c < 1$, the Meixner linear functional is

$$u = \sum_{k=0}^{\infty} \frac{\beta(\beta + 1) \cdots (\beta + k - 1)}{k!} c^k \delta(x - k).$$

See [4] for matrix extensions of these discrete linear functionals and corresponding matrix orthogonal polynomials.

Definition 13 (Hankel sesquilinear forms). Given a matrix of generalized functions as entries

$$u = \begin{bmatrix} u_{1,1} & \cdots & u_{1,p} \\ \vdots & & \vdots \\ u_{p,1} & \cdots & u_{p,p} \end{bmatrix},$$

i.e. $u_{i,j} \in (\mathbb{C}[x])'$, then the associated sesquilinear form $\langle P(x), Q(x) \rangle_u$ is given by

$$\langle \langle P(x), Q(x) \rangle_u \rangle_{i,j} := \sum_{k,l=1}^p \langle u_{k,l}, P_{i,k}(x) Q_{j,l}(x) \rangle.$$

When $u_{k,l} \in \mathcal{O}'_c$, we write $u \in (\mathcal{O}'_c)^{p \times p}$ and say that we have a matrix of fast decreasing distributions. In this case the support is defined as $\text{supp}(u) := \bigcup_{k,l=1}^N \text{supp}(u_{k,l})$.

Observe that in this Hankel case, we could also have continuous and discrete orthogonality.

Proposition 14. *In terms of the moments*

$$m_n := \begin{bmatrix} \langle u_{1,1}, x^n \rangle & \dots & \langle u_{1,p}, x^n \rangle \\ \vdots & & \vdots \\ \langle u_{p,1}, x^n \rangle & \dots & \langle u_{p,p}, x^n \rangle \end{bmatrix}$$

the Gram matrix of the sesquilinear form given in Definition 13 is the following moment matrix:

$$G := \begin{bmatrix} m_0 & m_1 & m_2 & \dots \\ m_1 & m_2 & m_3 & \dots \\ m_2 & m_3 & m_4 & \dots \\ \vdots & \vdots & \vdots & \end{bmatrix},$$

of Hankel-type.

1.2.2. Matrices of generalized kernels and sesquilinear forms

The previous examples all have in common the same Hankel block symmetry for the corresponding matrices. However, there are sesquilinear forms which do not have this particular Hankel-type symmetry. Let us stop for a moment at this point, and elaborate on bilinear and sesquilinear forms for polynomials. We first recall some facts regarding the scalar case with $p = 1$, and bilinear forms instead of sesquilinear forms. Given $u_{x,y} \in (\mathbb{C}[x, y])' = (\mathbb{C}[x, y])^* \cong \mathbb{C}[x, y]$, we can consider the continuous bilinear form $B(P(x), Q(y)) = \langle u_{x,y}, P(x) \otimes Q(y) \rangle$. This gives a continuous linear map $\mathcal{L}_u : \mathbb{C}[y] \rightarrow (\mathbb{C}[x])'$ such that $B(P(x), Q(y)) = \langle \mathcal{L}_u(Q(y)), P(x) \rangle$. The Gram matrix of this bilinear form has coefficients $G_{k,l} = B(x^k, y^l) = \langle u_{x,y}, x^k \otimes y^l \rangle = \langle \mathcal{L}_u(y^l), x^k \rangle$. Here we follow Schwartz discussion on kernels and distributions [66], see also [47]. A kernel $u(x, y)$ is a complex-valued locally integrable function, that defines an integral operator $f(x) \mapsto g(x) = \int u(x, y)f(y)dy$. Following [67] we denote $(\mathcal{D})_x$ and $(\mathcal{D}')_x$ the test functions and the corresponding distributions in the variable x , and similarly for the variable y . We extend this construction considering a bivariate distribution in the variables x, y , $u_{x,y} \in (\mathcal{D}')_{x,y}$, that Schwartz called *noyau-distribution*, and as we use a wider range of generalized functions we will call generalized kernel. This $u_{x,y}$ generates a continuous bilinear form $B_u(\phi(x), \psi(y)) = \langle u_{x,y}, \phi(x) \otimes \psi(y) \rangle$. It also generates a continuous linear map $\mathcal{L}_u : (\mathcal{D})_y \rightarrow (\mathcal{D}')_x$ with $\langle (\mathcal{L}_u(\psi(y)))_x, \phi(x) \rangle = \langle u_{x,y}, \phi(x) \otimes \psi(y) \rangle$. The Schwartz kernel theorem states that every generalized kernel $u_{x,y}$ defines a continuous linear transformation \mathcal{L}_u from $(\mathcal{D})_y$ to $(\mathcal{D}')_x$, and to each of such continuous linear transformations we can associate one and only one generalized kernel. According to the prolongation scheme

developed in [66], the generalized kernel $u_{x,y}$ is such that $\mathcal{L}_u : (\mathcal{E})_y \rightarrow (\mathcal{E}')_x$ if and only if the support of $u_{x,y}$ in \mathbb{R}^2 is compact.^a

We can extend these ideas to the matrix scenario of this paper, where instead of bilinear forms we have sesquilinear forms.

Definition 14. Given a matrix of generalized kernels

$$u_{x,y} := \begin{bmatrix} (u_{x,y})_{1,1} & \cdots & (u_{x,y})_{1,p} \\ \vdots & & \vdots \\ (u_{x,y})_{p,1} & \cdots & (u_{x,y})_{p,p} \end{bmatrix}$$

with $(u_{x,y})_{k,l} \in (\mathbb{C}[x,y])'$ or, if a notion of support is required, $(u_{x,y})_{k,l} \in (\mathcal{E}')_{x,y}, (\mathcal{O}'_M)_{x,y}, (\mathcal{O}'_c)_{x,y}$ provides a continuous sesquilinear form with entries given by

$$\begin{aligned} \langle P(x), Q(y) \rangle_u &= \sum_{k,l=1}^p \langle (u_{x,y})_{k,l}, P_{i,k}(x) \otimes Q_{j,l}(y) \rangle \\ &= \sum_{k,l=1}^p \langle \mathcal{L}_{u_{k,l}}(Q_{j,l}(y)), P_{i,k}(x) \rangle, \end{aligned}$$

where $\mathcal{L}_{u_{k,l}} : \mathbb{C}[y] \rightarrow (\mathbb{C}[x])'$ — or depending on the setting $\mathcal{L}_{u_{k,l}} : (\mathcal{E})_y \rightarrow (\mathcal{E}')_x$, $\mathcal{L}_{u_{k,l}} : (\mathcal{O}'_M)_y \rightarrow (\mathcal{O}'_c)_x$, for example — is a continuous linear operator. We can condensate it in a matrix form, for $u_{x,y} \in (\mathbb{C}^{p \times p}[x,y])' = (\mathbb{C}^{p \times p}[x,y])^* \cong \mathbb{C}^{p \times p}[x,y]$, a sesquilinear form is given

$$\langle P(x), Q(y) \rangle_u = \langle u_{x,y}, P(x) \otimes Q(y) \rangle = \langle \mathcal{L}_u(Q(y)), P(x) \rangle,$$

with $\mathcal{L}_u : \mathbb{C}^{p \times p}[y] \rightarrow (\mathbb{C}^{p \times p}[x])'$ a continuous linear map. Or, in other scenarios $\mathcal{L}_u : ((\mathcal{E})_y)^{p \times p} \rightarrow ((\mathcal{E}')_x)^{p \times p}$ or $\mathcal{L}_u : ((\mathcal{O}'_M)_y)^{p \times p} \rightarrow ((\mathcal{O}'_c)_x)^{p \times p}$.

If, instead of a matrix of bivariate distributions, we have a matrix of bivariate measures then we could write for the sesquilinear form $\langle P(x), Q(y) \rangle = \iint P(x) d\mu(x,y) (Q(y))^\top$, where $\mu(x,y)$ is a matrix of bivariate measures.

For the scalar case $p = 1$, Adler and van Moerbeke discussed in [1] different possibilities of non-Hankel Gram matrices. Their Gram matrix has as coefficients $G_{k,l} = \langle u_l, x^k \rangle$, for an infinite sequence of generalized functions u_l , that recovers the Hankel scenario for $u_l = x^l u$. They studied in more detail the following cases:

- (i) Banded case: $u_{l+km} = x^{km} u_l$.

^aUnderstood as a prolongation problem, see [66, §5], we have similar results if we require $\mathcal{L}_u : \mathcal{O}_M \rightarrow \mathcal{O}'_c$ or $\mathcal{L}_u : \mathcal{O}_c \rightarrow \mathcal{O}'_c$ or any other possibility that makes sense for polynomials and support.

- (ii) Concatenated solitons: $u_l(x) = \delta(x - p_{l+1}) - (\lambda_{l+1})^2 \delta(x - q_{k+1})$.
- (iii) Nested Calogero–Moser systems: $u_l(x) = \delta'(x - p_{l+1}) + \lambda_{l+1} \delta(x - p_{l+1})$.
- (iv) Discrete KdV soliton type: $u_l(x) = (-1)^k \delta^{(l)}(x - p) - \delta^{(l)}(x + p)$.

We see that the three last weights are generalized functions. To compare with the Schwartz’s approach we observe that $\langle u_{x,y}, x^k \otimes y^l \rangle = \langle u_l, x^k \rangle$ and, consequently, we deduce $u_l = \mathcal{L}_u(y^l)$ (and for continuous kernels $u_l(x) = \int u(x, y) y^l dy$). The first case has a banded structure and its Gram matrix fulfills $\Lambda^m G = G(\Lambda^\top)^m$. In [3], different examples are discussed for the matrix orthogonal polynomials, like bigraded Hankel matrices $\Lambda^n G = G(\Lambda^\top)^m$, where n, m are positive integers, can be realized as $G_{k,l} = \langle u_l, I_p x^k \rangle$, in terms of matrices of linear functionals u_l which satisfy the following periodicity condition $u_{l+m} = u_l x^n$. Therefore, given the linear functionals u_0, \dots, u_{m-1} we can recover all the others.

1.2.3. Sesquilinear forms supported by the diagonal and Sobolev sesquilinear forms

First we consider the scalar case

Definition 15. A generalized kernel $u_{x,y}$ is supported by the diagonal $y = x$ if

$$\langle u_{x,y}, \phi(x, y) \rangle = \sum_{n,m} \left\langle u_x^{(n,m)}, \frac{\partial^{n+m} \phi(x, y)}{\partial x^n \partial y^m} \Big|_{y=x} \right\rangle$$

for a locally finite sum and generalized functions $u_x^{(n,m)} \in (\mathcal{D}')_x$.

Proposition 15 (Sobolev bilinear forms). *The bilinear form corresponding to a generalized kernel supported by the diagonal is $B(\phi(x), \psi(x)) = \sum_{n,m} \langle u_x^{(n,m)}, \phi^{(n)}(x) \psi^{(m)}(x) \rangle$, which is of Sobolev type.*

For order zero $u_x^{(n,m)}$ generalized functions, i.e. for a set of Borel measures $\mu^{(n,m)}$, we have

$$B(\phi(x), \psi(x)) = \sum_{n,m} \int \phi^{(n)}(x) \psi^{(m)}(x) d\mu^{(n,m)}(x),$$

which is of Sobolev type. Thus, in the scalar case, generalized kernels supported by the diagonal are just Sobolev bilinear forms. The extension of these ideas to the matrix case is immediate, we only need to require to all generalized kernels to be supported by the diagonal.

Proposition 16 (Sobolev sesquilinear forms). *A matrix of generalized kernels supported by the diagonal provides Sobolev sesquilinear forms*

$$\langle \langle P(x), Q(x) \rangle_u \rangle_{i,j} = \sum_{k,l=1}^p \sum_{n,m} \left\langle u_{k,l}^{(n,m)}, P_{i,k}^{(n)}(x) Q_{j,l}^{(m)}(x) \right\rangle$$

for a locally finite sum, in the derivatives of order n, m , and of generalized functions $u_x^{(n,m)} \in (\mathbb{C}[x])'$. All Sobolev sesquilinear forms are obtained in this form.

For a recent review on scalar Sobolev orthogonal polynomials see [51]. Observe that with this general framework we could consider matrix discrete Sobolev orthogonal polynomials, that will appear whenever the linear functionals $u^{(m,n)}$ have infinite discrete support, as far as u is quasi-definite.

1.2.4. *Biorthogonality, quasi-definiteness and Gauss–Borel factorization*

Definition 16 (Biorthogonal matrix polynomials). Given a sesquilinear form $\langle \cdot, \cdot \rangle$, two sequences of matrix polynomials $\{P_n^{[1]}(x)\}_{n=0}^\infty$ and $\{P_n^{[2]}(x)\}_{n=0}^\infty$ are said to be biorthogonal with respect to $\langle \cdot, \cdot \rangle$ if

- (i) $\deg(P_n^{[1]}(x)) = \deg(P_n^{[2]}(x)) = n$ for all $n \in \{0, 1, \dots\}$,
- (ii) $\langle P_n^{[1]}(x), P_m^{[2]}(y) \rangle = \delta_{n,m} H_n$ for all $n, m \in \{0, 1, \dots\}$,

where H_n are nonsingular matrices and $\delta_{n,m}$ is the Kronecker delta.

Definition 17 (Quasi-definiteness). A Gram matrix of a sesquilinear form $\langle \cdot, \cdot \rangle_u$ is said to be quasi-definite whenever $\det G_{[k]} \neq 0$, $k \in \{0, 1, \dots\}$. Here $G_{[k]}$ denotes the truncation

$$G_{[k]} := \begin{bmatrix} G_{0,0} & \dots & G_{0,k-1} \\ \vdots & & \vdots \\ G_{k-1,0} & \dots & G_{k-1,k-1} \end{bmatrix}.$$

We say that the bivariate generalized function $u_{x,y}$ is quasi-definite and the corresponding sesquilinear form is nondegenerate whenever its Gram matrix is quasi-definite.

Proposition 17 (Gauss–Borel factorization, see [7]). *If the Gram matrix of a sesquilinear form $\langle \cdot, \cdot \rangle_u$ is quasi-definite, then there exists a unique Gauss–Borel factorization given by*

$$G = (S_1)^{-1} H (S_2)^{-\top}, \tag{6}$$

where S_1, S_2 are lower unitriangular block matrices and H is a diagonal block matrix

$$S_i = \begin{bmatrix} I_p & 0_p & 0_p & \dots \\ (S_i)_{1,0} & I_p & 0_p & \dots \\ (S_i)_{2,0} & (S_i)_{2,1} & I_p & \ddots \\ \vdots & \vdots & \ddots & \ddots \end{bmatrix}, \quad i = 1, 2, \quad H = \text{diag}(H_0, H_1, H_2, \dots),$$

with $(S_i)_{n,m}$ and $H_n \in \mathbb{C}^{p \times p}$, $\forall n, m \in \{0, 1, \dots\}$.

For $l \geq k$ we will also use the following bordered truncated Gram matrix:

$$G_{[k,l]}^{[1]} := \begin{bmatrix} G_{0,0} & \cdots & G_{0,k-1} \\ \vdots & & \vdots \\ G_{k-2,0} & \cdots & G_{k-2,k-1} \\ \hline G_{l,0} & \cdots & G_{l,k-1} \end{bmatrix},$$

where we have replaced the last row of blocks of the truncated Gram matrix $G_{[k]}$ by the row of blocks $[G_{l,0}, \dots, G_{l,k-1}]$. We also need a similar matrix but replacing the last block column of $G_{[k]}$ by a column of blocks as indicated

$$G_{[k,l]}^{[2]} := \left[\begin{array}{ccc|c} G_{0,0} & \cdots & G_{0,k-2} & G_{0,l} \\ \vdots & & \vdots & \vdots \\ G_{k-1,0} & \cdots & G_{k-1,k-2} & G_{k-1,l} \end{array} \right].$$

Using last quasideterminants, see [27, 58] and Appendix A, we find the following.

Proposition 18. *If the last quasideterminants of the truncated moment matrices are nonsingular, i.e.*

$$\det \Theta_*(G_{[k]}) \neq 0, \quad k = 1, 2, \dots,$$

then, the Gauss–Borel factorization can be performed and the following expressions are fulfilled:

$$H_k = \Theta_* \begin{bmatrix} G_{0,0} & G_{0,1} & \cdots & G_{0,k-1} \\ G_{1,0} & G_{1,1} & \cdots & G_{1,k-1} \\ \vdots & \vdots & & \vdots \\ G_{k-1,0} & G_{k-1,1} & \cdots & G_{k-1,k-1} \end{bmatrix},$$

$$(S_1)_{k,l} = \Theta_* \begin{bmatrix} G_{0,0} & G_{0,1} & \cdots & G_{0,k-1} & 0_p \\ G_{1,0} & G_{1,1} & \cdots & G_{1,k-1} & 0_p \\ \vdots & \vdots & & \vdots & \vdots \\ G_{l-1,0} & G_{l,1} & \cdots & G_{l-1,k-1} & 0_p \\ G_{l,0} & G_{l,1} & \cdots & G_{l,k-1} & I_p \\ G_{l+1,0} & G_{l+1,1} & \cdots & G_{l+1,k-1} & 0_p \\ \vdots & \vdots & & \vdots & \vdots \\ G_{k,0} & G_{k,1} & \cdots & G_{k,k-1} & 0_p \end{bmatrix},$$

$$((S_2)^\top)_{k,l} = \Theta_* \begin{bmatrix} G_{0,0} & G_{0,1} & \dots & G_{0,l-1} & G_{0,l} & G_{0,l+1} & \dots & G_{0,k} \\ G_{1,0} & G_{1,1} & \dots & G_{1,l-1} & G_{1,l} & G_{1,l+1} & \dots & G_{1,k} \\ \vdots & \vdots & & \vdots & \vdots & \vdots & & \vdots \\ G_{k-1,0} & G_{k-1,1} & \dots & G_{k-1,l-1} & G_{k-1,l} & G_{k-1,l+1} & \dots & G_{k-1,k} \\ 0_p & 0_p & \dots & 0_p & I_p & 0_p & \dots & 0_p \end{bmatrix},$$

and for the inverse elements [58] the formulas

$$(S_1^{-1})_{k,l} = \Theta_*(G_{[k,l+1]}^{[1]})\Theta_*(G_{[l+1]})^{-1},$$

$$(S_2^{-1})_{k,l} = (\Theta_*(G_{[l+1]})^{-1}\Theta_*(G_{[k,l+1]}^{[2]}))^\top,$$

hold true.

We see that the matrices H_k are quasideterminants, and following [7, 8] we refer to them as quasi-tau matrices.

1.2.5. *Biorthogonal polynomials, second kind functions and Christoffel–Darboux kernels*

Definition 18. We define $\chi(x) := [I_p, I_p x, I_p x^2, \dots]^\top$, and for $x \neq 0$, $\chi^*(x) := [I_p x^{-1}, I_p x^{-2}, I_p x^{-3}, \dots]^\top$.

Remark 1. Observe that the Gram matrix can be expressed as

$$G = \langle \chi(x), \chi(y) \rangle_u = \langle u_{x,y}, \chi(x) \otimes \chi(y) \rangle \tag{7}$$

and its block entries are

$$G_{k,l} = \langle I_p x^k, I_p y^l \rangle_u.$$

If the sesquilinear form derives from a matrix of bivariate measures $\mu(x, y) = [\mu_{i,j}(x, y)]$ we have for the Gram matrix blocks

$$G_{k,l} = \iint x^k d\mu(x, y) y^l,$$

which reduces for absolutely continuous measures with respect to the Lebesgue measure $dx dy$ to a matrix of weights $w(x, y) = [w_{i,j}(x, y)]$, and When the matrix of generalized kernels is Hankel we recover the classical Hankel structure, and the Gram matrix is a moment matrix. For example, for a matrix of measures we will have $G_{k,l} = \int x^{k+l} d\mu(x)$.

Definition 19. Given a quasi-definite matrix of generalized kernels $u_{x,y}$ and the Gauss–Borel factorization (17) of its Gram matrix, the corresponding first and

second families of matrix polynomials are

$$P^{[1]}(x) = \begin{bmatrix} P_0^{[1]}(x) \\ P_1^{[1]}(x) \\ \vdots \end{bmatrix} := S_1\chi(x), \quad P^{[2]}(y) = \begin{bmatrix} P_0^{[2]}(y) \\ P_1^{[2]}(y) \\ \vdots \end{bmatrix} := S_2\chi(y), \quad (8)$$

respectively.

Proposition 19 (Biorthogonality). *Given a quasi-definite matrix of generalized kernels $u_{x,y}$, the first and second families of monic matrix polynomials $\{P_n^{[1]}(x)\}_{n=0}^\infty$ and $\{P_n^{[2]}(x)\}_{n=0}^\infty$ are biorthogonal*

$$\langle P_n^{[1]}(x), P_m^{[2]}(y) \rangle_u = \delta_{n,m} H_n, \quad n, m \in \{0, 1, \dots\}. \quad (9)$$

Remark 2. The biorthogonal relations yield the orthogonality relations

$$\langle P_n^{[1]}(x), y^m I_p \rangle_u = 0_p, \quad \langle x^m I_p, P_n^{[2]}(y) \rangle_u = 0_p, \quad m \in \{1, \dots, n-1\}, \quad (10)$$

$$\langle P_n^{[1]}(x), y^n I_p \rangle_u = H_n, \quad \langle x^n I_p, P_n^{[2]}(y) \rangle_u = H_n. \quad (11)$$

Remark 3 (Symmetric generalized kernels). If $u_{x,y} = (u_{y,x})^\top$, the Gram matrix is symmetric $G = G^\top$ and we are dealing with a Cholesky block factorization with $S_1 = S_2$ and $H = H^\top$. Now $P_n^{[1]}(x) = P_n^{[2]}(x) =: P_n(x)$, and $\{P_n(x)\}_{n=0}^\infty$ is a set of monic orthogonal matrix polynomials. In this case $C_n^{[1]}(x) = C_n^{[2]}(x) =: C_n(x)$.

The shift matrix is the following semi-infinite block matrix:

$$\Lambda := \begin{bmatrix} 0_p & I_p & 0_p & 0_p & \dots \\ 0_p & 0_p & I_p & 0_p & \ddots \\ 0_p & 0_p & 0_p & I_p & \ddots \\ 0_p & 0_p & 0_p & 0_p & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \end{bmatrix}$$

which satisfies the spectral property

$$\Lambda\chi(x) = x\chi(x).$$

Proposition 20. *The symmetry of the block Hankel moment matrix reads $\Lambda G = G\Lambda^\top$.*

Notice that this symmetry completely characterizes Hankel block matrices.

Definition 20. The matrices $J_1 := S_1\Lambda(S_1)^{-1}$ and $J_2 := S_2\Lambda(S_2)^{-1}$ are the Jacobi matrices associated with the Gram matrix G .

The reader must notice the abuse in the notation. But for the sake of simplicity we have used the same letter for Jacobi and Jordan matrices. The type of matrix will be clear from the context.

Proposition 21. *The biorthogonal polynomials are eigenvectors of the Jacobi matrices*

$$J_1 P^{[1]}(x) = xP^{[1]}(x), \quad J_2 P^{[2]}(x) = xP^{[2]}(x).$$

and the second kind functions á la Gram satisfy

$$(H(J_2)^\top H^{-1})C^{[1]}(x) = xC^{[1]}(x) - H_0 \begin{bmatrix} I_p \\ 0_p \\ \vdots \end{bmatrix},$$

$$(H^\top (J_1)^\top H^{-\top})C^{[2]}(x) = xC^{[2]}(x) - H_0^\top \begin{bmatrix} I_p \\ 0_p \\ \vdots \end{bmatrix}.$$

Proposition 22. *For Hankel-type Gram matrices (i.e. associated with a matrix of univariate generalized functionals) the two Jacobi matrices are related by $H^{-1}J_1 = J_2^\top H^{-1}$, being, therefore, a tridiagonal matrix. This yields the three-term relation for biorthogonal polynomials and second kind functions, respectively.*

Proposition 23. *We have the following last quasideterminantal expressions:*

$$P_n^{[1]}(x) = \Theta_* \begin{bmatrix} G_{0,0} & G_{0,1} & \cdots & G_{0,n-1} & I_p \\ G_{1,0} & G_{1,1} & \cdots & G_{1,n-1} & I_p x \\ \vdots & \vdots & & \vdots & \vdots \\ G_{n-1,0} & G_{n-1,1} & \cdots & G_{n-1,n-1} & I_p x^{n-1} \\ G_{n,0} & G_{n,1} & \cdots & G_{n,n-1} & I_p x^n \end{bmatrix},$$

$$(P_n^{[2]}(y))^\top = \Theta_* \begin{bmatrix} G_{0,0} & G_{0,1} & \cdots & G_{0,n-1} & G_{0,n} \\ G_{1,0} & G_{1,1} & \cdots & G_{1,n-1} & G_{1,n} \\ \vdots & \vdots & & \vdots & \vdots \\ G_{n-1,0} & G_{n-1,1} & \cdots & G_{n-1,n-1} & G_{n-1,n} \\ I_p & I_p y & \cdots & I_p y^{n-1} & I_p y^n \end{bmatrix}.$$

Definition 21 (Christoffel–Darboux kernel [15, 68]). Given two sequences of matrix biorthogonal polynomials $\{P_k^{[1]}(x)\}_{k=0}^\infty$ and $\{P_k^{[2]}(y)\}_{k=0}^\infty$, with respect to

the sesquilinear form $\langle \cdot, \cdot \rangle_u$, we define the n th Christoffel–Darboux kernel matrix polynomial

$$K_n(x, y) := \sum_{k=0}^n (P_k^{[2]}(y))^\top (H_k)^{-1} P_k^{[1]}(x), \tag{12}$$

and the mixed Christoffel–Darboux kernel

$$K_n^{(pc)}(x, y) := \sum_{k=0}^n (P_k^{[2]}(y))^\top (H_k)^{-1} C_k^{[1]}(x).$$

Proposition 24. (i) *For a quasi-definite matrix of generalized kernels $u_{x,y}$, the corresponding Christoffel–Darboux kernel gives the projection operator*

$$\begin{aligned} \left\langle K_n(x, z), \sum_{0 \leq j \ll \infty} C_j P_j^{[2]}(y) \right\rangle_u &= \left(\sum_{j=0}^n C_j P_j^{[2]}(z) \right)^\top, \\ \left\langle \sum_{0 \leq j \ll \infty} C_j P_j^{[1]}(x), (K_n(z, y))^\top \right\rangle_u &= \sum_{j=0}^n C_j P_j^{[1]}(z). \end{aligned} \tag{13}$$

(ii) *In particular, we have*

$$\langle K_n(x, z), I_p y^l \rangle_u = I_p z^l, \quad l \in \{0, 1, \dots, n\}. \tag{14}$$

Proposition 25 (Christoffel–Darboux formula). *When the sesquilinear form is Hankel (now u is a matrix of univariate generalized functions with its Gram matrix of block Hankel-type) the Christoffel–Darboux kernel satisfies*

$$(x - y)K_n(x, y) = (P_n^{[2]}(y))^\top (H_n)^{-1} P_{n+1}^{[1]}(x) - (P_{n+1}^{[2]}(y))^\top (H_n)^{-1} P_n^{[1]}(x),$$

and the mixed Christoffel–Darboux kernel fulfills

$$(x - y)K_n^{(pc)}(x, y) = (P_n^{[2]}(y))^\top H_n^{-1} C_{n+1}^{[1]}(x) - (P_{n+1}^{[2]}(y))^\top H_n^{-1} C_n^{[1]}(x) + I_p.$$

Proof. We only prove the second formula, for the first one proceeds similarly. It is obviously a consequence of the three-term relation. First, let us notice that

$$J_2^\top H^{-1} C^{[1]}(x) = xH^{-1}C^{[1]}(x) - \begin{bmatrix} I_p \\ 0_p \\ \vdots \end{bmatrix}, \quad (P^{[2]}(y))^\top J_2^\top H^{-1} = y(P^{[2]}(y))^\top H^{-1}.$$

Second, we have

$$J_2^\top H^{-1} = \left[\begin{array}{ccc|ccc} & & & & 0 & 0 & \dots \\ & & & & \vdots & \vdots & \\ & & & & 0 & 0 & \dots \\ & & & & H_n^{-1} & 0 & \dots \\ \hline 0 & \dots & 0 & H_n^{-1} & & & \\ 0 & \dots & 0 & 0 & & & \\ \vdots & & \vdots & \vdots & & * & \end{array} \right].$$

Using this, we calculate the $(P_{[n]}^{[2]}(y))^\top [J_2^\top H^{-1}]_{[n]} C_{[n]}^{[1]}(x)$, first by computing the action of middle matrix on its left and then on its right to get

$$\begin{aligned} xK_{n-1}^{(pc)}(x, y) - (P_{n-1}^{[2]}(y))^\top H_n^{-1} C_n^{[1]}(x) - P_0 \\ = yK_{n-1}^{(pc)}(x, y) - (P_n^{[2]}(y))^\top H_n^{-1} C_{n-1}^{[1]}(x), \end{aligned}$$

and since $P_0 = I_p$ the proposition is proven. □

Next, we deal with the fact that our definition of second kind functions implies non-admissible products and do involve series.

Definition 22. For the support of the matrix of generalized kernels $\text{supp}(u_{x,y}) \subset \mathbb{C}^2$ we consider the action of the component projections $\pi_1, \pi_2: \mathbb{C}^2 \rightarrow \mathbb{C}$ on its first and second variables, $(x, y) \xrightarrow{\pi_1} x$, $(x, y) \xrightarrow{\pi_2} y$, respectively, and introduce the projected supports $\text{supp}_x(u) := \pi_1(\text{supp}(u_{x,y}))$ and $\text{supp}_y(u) := \pi_2(\text{supp}(u_{x,y}))$, both subsets of \mathbb{C} . We will assume that $r_x := \sup\{|z| : z \in \text{supp}_x u\} < \infty$ and $r_y := \sup\{|z| : z \in \text{supp}_y u\} < \infty$. We also consider the disks about infinity, or annulus around the origin, $D_x := \{z \in \mathbb{C} : |z| > r_x\}$ and $D_y := \{z \in \mathbb{C} : |z| > r_y\}$.

Definition 23 (Second kind functions á la Cauchy). For a generalized kernel it is such that $u_{x,y} \in ((\mathcal{O}'_c)_{x,y})^{p \times p}$ we define two families of second kind functions á la Cauchy given by

$$\begin{aligned} C_n^{[1]}(z) &= \left\langle P_n^{[1]}(x), \frac{I_p}{z - y} \right\rangle_u, \quad z \notin \text{supp}_y(u), \\ (C_n^{[2]}(z))^\top &= \left\langle \frac{I_p}{z - x}, P_n^{[2]}(y) \right\rangle_u, \quad z \notin \text{supp}_x(u). \end{aligned}$$

2. Matrix Geronimus Transformations

Geronimus transformations for scalar orthogonal polynomials were first discussed in [35], where some determinantal formulas were found, see [55, 73]. Geronimus perturbations of degree two of scalar bilinear forms have been very recently treated

in [17] and in the general case in [16]. Here we discuss its matrix extension for general sesquilinear forms.

Definition 24. Given a matrix of generalized kernels $u_{x,y} = ((u_{x,y})_{i,j}) \in ((\mathcal{O}'_c)_{x,y})^{p \times p}$ with a given support $\text{supp } u_{x,y}$, and a matrix polynomial $W(y) \in \mathbb{C}^{p \times p}[y]$ of degree N , such that $\sigma(W(y)) \cap \text{supp}_y(u) = \emptyset$, a matrix of bivariate generalized functions $\check{u}_{x,y}$ is said to be a matrix Geronimus transformation of the matrix of generalized kernels $u_{x,y}$ if

$$\check{u}_{x,y}W(y) = u_{x,y}. \tag{15}$$

Proposition 26. *In terms of sesquilinear forms a Geronimus transformation fulfills*

$$\langle P(x), Q(y)(W(y))^\top \rangle_{\check{u}} = \langle P(x), Q(y) \rangle_u,$$

while, in terms of the corresponding Gram matrices, satisfies

$$\check{G}W(\Lambda^\top) = G.$$

We will assume that the perturbed moment matrix has a Gauss–Borel factorization $\check{G} = \check{S}_1^{-1}\check{H}(\check{S}_2)^{-\top}$, where \check{S}_1, \check{S}_2 are lower unitriangular block matrices and \check{H} is a diagonal block matrix

$$\check{S}_i = \begin{bmatrix} I_p & 0_p & 0_p & \cdots \\ (\check{S}_i)_{1,0} & I_p & 0_p & \cdots \\ (\check{S}_i)_{2,0} & (\check{S}_i)_{2,1} & I_p & \ddots \\ & & \ddots & \ddots \end{bmatrix}, \quad i = 1, 2, \quad \check{H} = \text{diag}(\check{H}_0, \check{H}_1, \check{H}_2, \dots).$$

Hence, the Geronimus transformation provides the family of matrix biorthogonal polynomials

$$\check{P}^{[1]}(x) = \check{S}_1\chi(x), \quad \check{P}^{[2]}(y) = \check{S}_2\chi(y),$$

with respect to the perturbed sesquilinear form $\langle \cdot, \cdot \rangle_{\check{u}}$.

Observe that the matrix generalized kernels $v_{x,y}$ such that $v_{x,y}W(y) = 0_p$ can be added to a Geronimus transformed matrix of generalized kernels $\check{u}_{x,y} \mapsto \check{u}_{x,y} + v_{x,y}$, to get a new Geronimus transformed matrix of generalized kernels. We call masses these type of terms.

2.1. The resolvent and connection formulas

Definition 25. The resolvent matrix is

$$\omega := \check{S}_1(S_1)^{-1}. \tag{16}$$

The key role of this resolvent matrix is determined by the following properties.

Proposition 27. (i) *The resolvent matrix can be also expressed as*

$$\omega = \check{H}(\check{S}_2)^{-\top} W(\Lambda^\top)(S_2)^\top H^{-1}, \tag{17}$$

where the products in the right-hand side are associative.

(ii) *The resolvent matrix is a lower unitriangular block banded matrix — with only the first N block subdiagonals possibly not zero, i.e.*

$$\omega = \begin{bmatrix} I_p & 0_p & \cdots & 0_p & 0_p & \cdots \\ \omega_{1,0} & I_p & \ddots & 0_p & 0_p & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \\ \omega_{N,0} & \omega_{N,1} & \cdots & I_p & 0_p & \ddots \\ 0_p & \omega_{N+1,1} & \cdots & \omega_{N+1,N} & I_p & \ddots \\ \vdots & \ddots & \ddots & & \ddots & \ddots \end{bmatrix}.$$

(iii) *The following connection formulas are satisfied*

$$\check{P}^{[1]}(x) = \omega P^{[1]}(x), \tag{18}$$

$$(\check{H}^{-1}\omega H)^\top \check{P}^{[2]}(y) = P^{[2]}(y)W^\top(y). \tag{19}$$

(iv) *For the last subdiagonal of the resolvent we have*

$$\omega_{N+k,k} = \check{H}_{N+k}A_N(H_k)^{-1}. \tag{20}$$

Proof. (i) From Proposition 26 and the Gauss–Borel factorization of G and \check{G} we get

$$(S_1)^{-1}H(S_2)^{-\top} = ((\check{S}_1)^{-1}\check{H}(\check{S}_2)^{-\top})W(\Lambda^\top),$$

so that

$$\check{S}_1(S_1)^{-1}H = \check{H}(\check{S}_2)^{-\top}W(\Lambda^\top)(S_2)^\top.$$

(ii) The resolvent matrix, being a product of lower unitriangular matrices, is a lower unitriangular matrix. However, from (17) we deduce that it is a matrix with all its subdiagonals with zero coefficients but for the first N . Thus, it must have the described band structure.

(iii) From the definition we have (18). Let us notice that (17) can be written as

$$\omega^\top \check{H}^{-\top} = H^{-\top} S_2 W^\top(\Lambda)(\check{S}_2)^{-1},$$

so that

$$\omega^\top \check{H}^{-\top} \check{P}^{[2]}(y) = H^{-\top} S_2 W^\top(\Lambda)\chi(y),$$

and (19) follows.

(iv) It is a consequence of (17). □

The connection formulas (18) and (19) can be written as

$$\check{P}_n^{[1]}(x) = P_n^{[1]}(x) + \sum_{k=n-N}^{n-1} \omega_{n,k} P_k^{[1]}(x), \tag{21}$$

$$W(y)(P_n^{[2]}(y))^\top (H_n)^{-1} = (\check{P}_n^{[2]}(y))^\top (\check{H}_n)^{-1} + \sum_{k=n+1}^{n+N} (\check{P}_k^{[2]}(y))^\top (\check{H}_k)^{-1} \omega_{k,n}. \tag{22}$$

Lemma 3. *We have that*

$$W(\Lambda^\top) \chi^*(x) = \chi^*(x) W(x) - \begin{bmatrix} \mathcal{B}(\chi(x))_{[N]} \\ 0_p \\ \vdots \end{bmatrix}, \tag{23}$$

with \mathcal{B} given in Definition 10.

Proposition 28. *The Geronimus transformation of the second kind functions satisfies*

$$\check{C}^{[1]}(x) W(x) - \begin{bmatrix} (\check{H}(\check{S}_2)^{-\top})_{[N]} \mathcal{B}(\chi(x))_{[N]} \\ 0_p \\ \vdots \end{bmatrix} = \omega C^{[1]}(x), \tag{24}$$

$$(\check{C}^{[2]}(x))^\top \check{H}^{-1} \omega = (C^{[2]}(x))^\top H^{-1}. \tag{25}$$

Proof. To get (24) we argue as follows

$$\begin{aligned} \check{C}^{[1]}(z) W(z) - \omega C^{[1]}(z) &= \left\langle \check{P}_1(x), \frac{I_p}{z-y} \right\rangle_{\check{u}} W(z) - \left\langle \check{P}_1(x), \frac{I_p}{z-y} \right\rangle_{\check{u}W} \\ &\text{use (18) and (15)} \\ &= \left\langle \check{P}_1(x), \frac{W(z) - W(y)}{z-y} \right\rangle_{\check{u}}. \end{aligned}$$

But, we have

$$\begin{aligned} \frac{W(z) - W(y)}{z-y} &= I_p \frac{z^N - y^N}{z-y} + A_{N-1} \frac{z^{N-1} - y^{N-1}}{z-y} + \dots + A_1 \\ &= I_p h_{N-1}(z, y) + A_{N-1} h_{N-2}(z, y) + \dots + A_1 \\ &= (\chi(y))^\top \begin{bmatrix} \mathcal{B}(\chi(z))_N \\ 0 \end{bmatrix} \end{aligned}$$

so that

$$\begin{aligned} \check{C}^{[1]}(z) W(z) - \omega C^{[1]}(z) &= \check{S}_1 \langle \chi(x), \chi(y) \rangle_{\check{u}} \begin{bmatrix} \mathcal{B}(\chi(z))_N \\ 0 \end{bmatrix} \\ &= \check{S}_1 \check{G} \begin{bmatrix} \mathcal{B}(\chi(x))_N \\ 0 \end{bmatrix} \end{aligned}$$

and using the Gauss–Borel factorization the result follows. For (25) we have

$$\begin{aligned}
 & (\check{C}^{[2]}(x))^\top \check{H}^{-1} \omega - (C^{[2]}(x))^\top H^{-1} \\
 &= \left\langle \frac{I_p}{z-x}, \check{P}^{[2]}(y) \right\rangle_{\check{u}} \check{H}^{-1} \omega - \left\langle \frac{I_p}{z-x}, P^{[2]}(y) \right\rangle_u H^{-1} \\
 &= \left\langle \frac{I_p}{z-x}, (\check{H}^{-1} \omega)^\top \check{P}^{[2]}(y) \right\rangle_{\check{u}} - \left\langle \frac{I_p}{z-x}, H^{-\top} P^{[2]}(y) \right\rangle_u \\
 &= \left\langle \frac{I_p}{z-x}, H^{-\top} P^{[2]}(y) (W(y))^\top \right\rangle_{\check{u}} - \left\langle \frac{I_p}{z-x}, H^{-\top} P^{[2]}(y) \right\rangle_u \\
 &= 0.
 \end{aligned}$$

□

Observe that the corresponding entries are

$$(C_n^{[2]}(y))^\top (H_k)^{-1} = (\check{C}_n^{[2]}(y))^\top (\check{H}_n)^{-1} + \sum_{k=n+1}^{n+N} (\check{C}_k^{[2]}(y))^\top (\check{H}_k)^{-1} \omega_{n,k}. \quad (26)$$

2.2. Geronimus transformations and Christoffel–Darboux kernels

Definition 26. The resolvent wing is the matrix

$$\Omega[n] = \begin{cases} \begin{bmatrix} \omega_{n,n-N} & \dots & \dots & \omega_{n,n-1} \\ 0_p & \ddots & & \vdots \\ \vdots & \ddots & \ddots & \vdots \\ 0_p & \dots & 0_p & \omega_{n+N-1,n-1} \end{bmatrix} & \in \mathbb{C}^{Np \times Np}, \quad n \geq N, \\ \begin{bmatrix} \omega_{n,0} & \dots & \dots & \omega_{n,n-1} \\ \vdots & & & \vdots \\ \omega_{N,0} & & & \omega_{N,n-1} \\ 0_p & \ddots & & \\ \vdots & \ddots & \ddots & \vdots \\ 0_p & \dots & 0_p & \omega_{n+N-1,n-1} \end{bmatrix} & \in \mathbb{C}^{Np \times np}, \quad n < N. \end{cases}$$

Theorem 1. For $m = \min(n, N)$, the perturbed and original Christoffel–Darboux kernels are related by the following connection formula:

$$\begin{aligned}
 \check{K}_{n-1}(x, y) &= W(y) K_{n-1}(x, y) - [(\check{P}_n^{[2]}(y))^\top \check{H}_n^{-1}, \dots, (\check{P}_{n+N-1}^{[2]}(y))^\top \check{H}_{n+N-1}^{-1}] \\
 &\quad \times \Omega[n] \begin{bmatrix} P_{n-m}^{[1]}(x) \\ \vdots \\ P_{n-1}^{[1]}(x) \end{bmatrix}. \quad (27)
 \end{aligned}$$

For $n \geq N$, the connection formula for the mixed Christoffel–Darboux kernels is

$$\check{K}_{n-1}^{(pc)}(x, y)W(x) = W(y)K_{n-1}^{(pc)}(x, y) - [(\check{P}_n^{[2]}(y))^\top \check{H}_n^{-1}, \dots, (\check{P}_{n+N-1}^{[2]}(y))^\top \check{H}_{n+N-1}^{-1}] \Omega[n] \begin{bmatrix} C_{n-N}^{[1]}(x) \\ \vdots \\ C_{n-1}^{[1]}(x) \end{bmatrix} + \mathcal{V}(x, y), \quad (28)$$

where $\mathcal{V}(x, y)$ was introduced in Definition 11.

Proof. For the first connection formula (27) we consider the pairing

$$\mathcal{K}_{n-1}(x, y) := [(\check{P}_0^{[2]}(y))^\top (\check{H}_0)^{-1}, \dots, (\check{P}_{n-1}^{[2]}(y))^\top (\check{H}_{n-1})^{-1}] \omega_{[n]} \begin{bmatrix} P_0^{[1]}(x) \\ \vdots \\ P_{n-1}^{[1]}(x) \end{bmatrix},$$

and compute it in two different ways. From (21) we get

$$\omega_{[n]} \begin{bmatrix} P_0^{[1]}(x) \\ \vdots \\ P_{n-1}^{[1]}(x) \end{bmatrix} = \begin{bmatrix} \check{P}_0^{[1]}(x) \\ \vdots \\ \check{P}_{n-1}^{[1]}(x) \end{bmatrix},$$

and, therefore, $\mathcal{K}_{n-1}(x, y) = \check{K}_{n-1}(x, y)$. Relation (22) leads to

$$\mathcal{K}_{n-1}(x, y) = W(y)K_{n-1}(x, y) - [(\check{P}_n^{[2]}(y))^\top (\check{H}_n)^{-1}, \dots, (\check{P}_{n+N-1}^{[2]}(y))^\top (\check{H}_{n+N-1})^{-1}] \Omega[n] \begin{bmatrix} P_{n-m}^{[1]}(x) \\ \vdots \\ P_{n-1}^{[1]}(x) \end{bmatrix},$$

and (27) is proven.

To derive (28) we consider the pairing

$$\mathcal{K}_{n-1}^{(pc)}(x, y) := [(\check{P}_0^{[2]}(y))^\top (\check{H}_0)^{-1}, \dots, (\check{P}_{n-1}^{[2]}(y))^\top (\check{H}_{n-1})^{-1}] \omega_{[n]} \begin{bmatrix} C_0^{[1]}(x) \\ \vdots \\ C_{n-1}^{[1]}(x) \end{bmatrix},$$

which, as before, can be computed in two different forms. On the one hand, using (24) we get

$$\begin{aligned} \mathcal{K}_{n-1}^{(pc)}(x, y) &= [(\check{P}_0^{[2]}(y))^\top (\check{H}_0)^{-1}, \dots, (\check{P}_{n-1}^{[2]}(y))^\top (\check{H}_{n-1})^{-1}] \\ &\quad \times \left(\begin{bmatrix} \check{C}_0^{[1]}(x)W(x) \\ \vdots \\ \check{C}_{n-1}^{[1]}(x)W(x) \end{bmatrix} - (\check{H}(\check{S}_2)^{-\top})_{[n,N]} \mathcal{B}(\chi(x))_{[N]} \right) \\ &= \check{K}_{n-1}^{(pc)}(x, y)W(x) - ((\chi(y))_{[n]})^\top ((\check{S}_2)^\top \check{H}^{-1})_{[n]} \\ &\quad \times (\check{H}(\check{S}_2)^{-\top})_{[n,N]} \mathcal{B}(\chi(x))_{[N]}, \end{aligned}$$

where $(\check{H}(\check{S}_2)^{-\top})_{[n,N]}$ is the truncation to the first n block rows and first N block columns of $\check{H}(\check{S}_2)^{-\top}$. This simplifies for $n \geq N$ to

$$\mathcal{K}_{n-1}^{(pc)}(x, y) = \check{K}_{n-1}^{(pc)}(x, y)W(x) - ((\chi(y))_{[N]})^\top \mathcal{B}(\chi(x))_{[N]}.$$

On the other hand, from (22) we conclude

$$\begin{aligned} \mathcal{K}_{n-1}^{(pc)}(x, y) &= W(y)K_{n-1}^{(pc)}(x, y) - [(\check{P}_n^{[2]}(y))^\top (\check{H}_n)^{-1}, \dots, \\ &\quad (\check{P}_{n+N-1}^{[2]}(y))^\top (\check{H}_{n+N-1})^{-1}] \Omega[n] \begin{bmatrix} C_{n-N}^{[1]}(x) \\ \vdots \\ C_{n-1}^{[1]}(x) \end{bmatrix}, \end{aligned}$$

and, consequently, we obtain

$$\begin{aligned} \check{K}_{n-1}^{(pc)}(x, y)W(x) &= W(y)K_{n-1}^{(pc)}(x, y) - [(\check{P}_n^{[2]}(y))^\top (\check{H}_n)^{-1}, \dots, \\ &\quad (\check{P}_{n+N-1}^{[2]}(y))^\top (\check{H}_{n+N-1})^{-1}] \Omega[n] \begin{bmatrix} C_{n-N}^{[1]}(x) \\ \vdots \\ C_{n-1}^{[1]}(x) \end{bmatrix} \\ &\quad + ((\chi(y))_{[N]})^\top \mathcal{B}(\chi(x))_{[N]}. \end{aligned}$$

□

2.3. Spectral jets and relations for the perturbed polynomials and its second kind functions

For the time being we will assume that the perturbing polynomial is monic, $W(x) = I_p x^N + \sum_{k=0}^{N-1} A_k x^k \in \mathbb{C}^{p \times p}[x]$.

Definition 27. Given a perturbing monic matrix polynomial $W(y)$ the most general mass term will have the form

$$v_{x,y} := \sum_{a=1}^q \sum_{j=1}^{s_a} \sum_{m=0}^{\kappa_j^{(a)}-1} \frac{(-1)^m}{m!} (\xi_{j,m}^{[a]})_x \otimes \delta^{(m)}(y - x_a) l_j^{(a)}(y), \tag{29}$$

expressed in terms of derivatives of Dirac linear functionals and adapted left root polynomials $l_j^{(a)}(x)$ of $W(x)$, and for vectors of generalized functions $(\xi_{j,m}^{[a]})_x \in ((\mathbb{C}[x])')^p$. Discrete Hankel masses appear when these terms are supported by the diagonal with

$$v_{x,x} := \sum_{a=1}^q \sum_{j=1}^{s_a} \sum_{m=0}^{\kappa_j^{(a)}-1} (-1)^m \delta^{(m)}(x - x_a) \frac{\xi_{j,m}^{[a]}}{m!} l_j^{(a)}(x), \tag{30}$$

with $\xi_{j,m}^{[a]} \in \mathbb{C}^p$.

Remark 4. Observe that the Hankel masses (30) are particular cases of (29) with

$$v_{x,y} := \sum_{a=1}^q \sum_{j=1}^{s_a} \sum_{m=0}^{\kappa_j^{(a)}-1} (-1)^m \frac{\xi_{j,m}^{[a]}}{m!} \sum_{k=0}^m \binom{m}{k} \delta^{(m-k)}(x - x_a) \otimes \delta^{(k)}(y - x_a) l_j^{(a)}(y),$$

so that, with the particular choice in (29),

$$(\xi_{j,k}^{[a]})_x = \sum_{n=0}^{\kappa_j^{(a)}-1-k} (-1)^n \frac{\xi_{j,k+n}^{[a]}}{n!} \delta^{(n)}(x - x_a),$$

we get the diagonal case.

Remark 5. For the sesquilinear forms we have

$$\begin{aligned} \langle P(x), Q(y) \rangle_{\bar{u}} &= \langle P(x), Q(y) (W(y))^{-\top} \rangle_u \\ &+ \sum_{a=1}^q \sum_{j=1}^{s_a} \sum_{m=0}^{\kappa_j^{(a)}-1} \langle P(x), (\xi_{j,m}^{[a]})_x \rangle \frac{1}{m!} (l_j^{(a)}(y) (Q(y))^\top)_{x_a}^{(m)}. \end{aligned}$$

Observe that the distribution $v_{x,y}$ is associated with the eigenvalues and left root vectors of the perturbing polynomial $W(x)$. Needless to say that, when $W(x)$ has a singular leading coefficient, this spectral part could even disappear, for example if $W(x)$ is unimodular; i.e. with constant determinant, not depending on x . Notice

that, in general, we have $Np \geq \sum_{a=1}^q \sum_{i=1}^{s_a} \kappa_j^{(a)}$ and we cannot ensure the equality, up to the nonsingular leading coefficient case.

Definition 28. Given a set of generalized functions $(\xi_{i,m}^{[a]})_x$, we introduce the matrices

$$\begin{aligned} \langle \check{P}_n^{[1]}(x), (\xi_i^{[a]})_x \rangle &:= [\langle \check{P}_n^{[1]}(x), (\xi_{i,0}^{[a]})_x \rangle, \langle \check{P}_n^{[1]}(x), (\xi_{i,1}^{[a]})_x \rangle, \dots, \\ &\quad \langle \check{P}_n^{[1]}(x), (\xi_{i,\kappa_i^{(a)}-1}^{[a]})_x \rangle] \in \mathbb{C}^{p \times \kappa_i^{(a)}}, \\ \langle \check{P}_n^{[1]}(x), (\xi^{[a]})_x \rangle &:= [\langle \check{P}_n^{[1]}(x), (\xi_1^{[a]})_x \rangle, \langle \check{P}_n^{[1]}(x), (\xi_2^{[a]})_x \rangle, \dots, \\ &\quad \langle \check{P}_n^{[1]}(x), (\xi_{s_a}^{[a]})_x \rangle] \in \mathbb{C}^{p \times \alpha_a}, \\ \langle \check{P}_n^{[1]}(x), (\xi)_x \rangle &:= [\langle \check{P}_n^{[1]}(x), (\xi^{[1]})_x \rangle, \langle \check{P}_n^{[1]}(x), (\xi^{[2]})_x \rangle, \dots, \\ &\quad \langle \check{P}_n^{[1]}(x), (\xi^{[q]})_x \rangle] \in \mathbb{C}^{p \times Np}. \end{aligned}$$

Definition 29. The exchange matrix is

$$\eta_i^{(a)} := \begin{bmatrix} 0 & 0 & \dots & 0 & 1 \\ 0 & 0 & \dots & 1 & 0 \\ \vdots & & \ddots & & \vdots \\ 0 & 1 & \dots & 0 & 0 \\ 1 & 0 & \dots & 0 & 0 \end{bmatrix} \in \mathbb{C}^{\kappa_i^{(a)} \times \kappa_i^{(a)}}.$$

Definition 30. The left Jordan chain matrix is given by

$$\mathcal{L}_i^{(a)} := \begin{bmatrix} l_{i,0}^{(a)} & l_{i,1}^{(a)} & l_{i,2}^{(a)} & \dots & l_{i,\kappa_i^{(a)}-1}^{(a)} \\ 0_{1 \times p} & l_{i,0}^{(a)} & l_{i,1}^{(a)} & \dots & l_{i,\kappa_i^{(a)}-2}^{(a)} \\ 0_{1 \times p} & 0_{1 \times p} & l_{i,0}^{(a)} & \dots & l_{i,\kappa_i^{(a)}-3}^{(a)} \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0_{1 \times p} & 0_{1 \times p} & & & l_{i,0}^{(a)} \end{bmatrix} \in \mathbb{C}^{\kappa_i^{(a)} \times p\kappa_i^{(a)}}.$$

For $z \neq x_a$, we also introduce the $p \times p$ matrices

$$\check{C}_{n,i}^{(a)}(z) := \langle \check{P}_n^{[1]}(x), (\xi_i^{[a]})_x \rangle \eta_i^{(a)} \mathcal{L}_i^{(a)} \begin{bmatrix} I_p \\ \hline (z - x_a)^{\kappa_i^{(a)}} \\ \vdots \\ I_p \\ \hline z - x_a \end{bmatrix}, \tag{31}$$

where $i = 1, \dots, s_a$.

Remark 6. Assume that the mass matrix is as in (30). Then, in terms of

$$\mathcal{X}_i^{(a)} := \begin{bmatrix} \xi_{i,\kappa_i^{(a)}-1}^{[a]} & \xi_{i,\kappa_i^{(a)}-2}^{[a]} & \xi_{i,\kappa_i^{(a)}-3}^{[a]} & \cdots & \xi_{i,0}^{[a]} \\ 0_{p \times 1} & \xi_{i,\kappa_i^{(a)}-1}^{[a]} & \xi_{i,\kappa_i^{(a)}-2}^{[a]} & \cdots & \xi_{i,1}^{[a]} \\ 0_{p \times 1} & 0_{p \times 1} & \xi_{i,\kappa_i^{(a)}-1}^{[a]} & \cdots & \xi_{i,2}^{[a]} \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0_{p \times 1} & 0_{p \times 1} & & & \xi_{i,\kappa_i^{(a)}-1}^{[a]} \end{bmatrix} \in \mathbb{C}^{p\kappa_i^{(a)} \times \kappa_i^{(a)}}, \quad (32)$$

we can write

$$\langle \check{P}_n^{[1]}(x), (\xi_i^{[a]})_x \rangle \eta_i^{(a)} = \mathcal{J}_{\check{P}_n^{[1]}}^{(i)}(x_a) \mathcal{X}_i^{(a)}. \quad (33)$$

Consequently,

$$\check{C}_{n;i}^{(a)}(z) := \mathcal{J}_{\check{P}_n^{[1]}}^{(i)}(x_a) \mathcal{X}_i^{(a)} \mathcal{L}_i^{(a)} \begin{bmatrix} I_p \\ (z-x_a)^{\kappa_i^{(a)}} \\ \vdots \\ I_p \\ z-x_a \end{bmatrix}.$$

Observe that $\mathcal{X}_i^{(a)} \mathcal{L}_i^{(a)} \in \mathbb{C}^{p\kappa_i^{(a)} \times p\kappa_i^{(a)}}$ is a block upper triangular matrix, with blocks in $\mathbb{C}^{p \times p}$.

Proposition 29. For $z \notin \text{supp}_y(\check{u}) = \text{supp}_y(u) \cup \sigma(W(y))$, the following expression

$$\check{C}_n^{[1]}(z) = \left\langle \check{P}_n^{[1]}(x), \frac{I_p}{z-y} \right\rangle_{uW^{-1}} + \sum_{a=1}^q \sum_{i=1}^{s_a} \check{C}_{n;i}^{(a)}(z)$$

holds.

Proof. We have

$$\begin{aligned} \check{C}_n^{[1]}(z) &= \left\langle \check{P}_n^{[1]}(x), \frac{I_p}{z-y} \right\rangle_{\check{u}} \\ &= \left\langle \check{P}_n^{[1]}(x), \frac{I_p}{z-y} \right\rangle_{uW^{-1}} + \sum_{a=1}^q \sum_{i=1}^{s_a} \sum_{m=0}^{\kappa_i^{(a)}-1} \langle \check{P}_n^{[1]}(x), (\xi_{j,m}^{[a]})_x \rangle \\ &\quad \times \left(\frac{1}{m!} \frac{l_i^{(a)}(x)}{z-x} \right)_{x_a}^{(m)}. \end{aligned}$$

Now, taking into account that

$$\left(\frac{1}{m!} \frac{l_i^{(a)}(x)}{z-x} \right)_{x=x_a}^{(m)} = \sum_{k=0}^m \left(\frac{l_i^{(a)}(x)}{(m-k)!} \right)_{x_a}^{(m-k)} \frac{1}{(z-x_a)^{k+1}},$$

we deduce the result. □

Lemma 4. Let $r_j^{(a)}(x)$ be right root polynomials of the monic matrix polynomial $W(x)$ given in (2), then

$$\begin{aligned} \mathcal{L}_i^{(a)} \begin{bmatrix} I_p \\ \hline (x - x_a)^{\kappa_i^{(a)}} \\ \vdots \\ \hline I_p \\ \hline x - x_a \end{bmatrix} W(x)r_j^{(a)}(x) \\ = \begin{bmatrix} 1 \\ \hline (x - x_a)^{\kappa_i^{(a)}} \\ \vdots \\ \hline 1 \\ \hline x - x_a \end{bmatrix} l_i^{(a)}(x)W(x)r_j^{(a)}(x) + (x - x_a)^{\kappa_j^{(a)}} T(x), \quad T(x) \in \mathbb{C}^{\kappa_j^{(a)}}[x]. \end{aligned}$$

Proof. Notice that we can write

$$\begin{aligned} \mathcal{L}_i^{(a)} \begin{bmatrix} I_p \\ \hline (x - x_a)^{\kappa_i^{(a)}} \\ \vdots \\ \hline I_p \\ \hline x - x_a \end{bmatrix} W(x)r_j^{(a)}(x) \\ = \begin{bmatrix} l_{i,0}^{(a)} & l_{i,1}^{(a)} & l_{i,2}^{(a)} & \cdots & l_{i,\kappa_i^{(a)}-1}^{(a)} \\ 0_{1 \times p} & l_{i,0}^{(a)} & l_{i,1}^{(a)} & \cdots & l_{i,\kappa_i^{(a)}-2}^{(a)} \\ 0_{1 \times p} & 0_{1 \times p} & l_{i,0}^{(a)} & & l_{i,\kappa_i^{(a)}-3}^{(a)} \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0_{1 \times p} & 0_{1 \times p} & & & l_{i,0}^{(a)} \end{bmatrix} \begin{bmatrix} I_p \\ \hline (x - x_a)^{\kappa_i^{(a)}} \\ \vdots \\ \hline I_p \\ \hline x - x_a \end{bmatrix} W(x)r_j^{(a)}(x) \\ = \begin{bmatrix} \frac{l_i^{(a)}(x)}{(x - x_a)^{\kappa_i^{(a)}}} \\ \frac{l_i^{(a)}(x)}{(x - x_a)^{\kappa_i^{(a)}-1} - l_{i,\kappa_i^{(a)}-1}^{(a)}} \\ \vdots \\ \frac{l_i^{(a)}(x)}{x - x_a} - l_{i,1}^{(a)} - \cdots - l_{i,\kappa_i^{(a)}-1}^{(a)}(x - x_a)^{\kappa_i^{(a)}-2} \end{bmatrix} W(x)r_j^{(a)}(x). \end{aligned}$$

□

Lemma 5. *The function $\check{C}_{n;i}^{(a)}(x)W(x)r_j^{(b)}(x) \in \mathbb{C}^p[x]$ satisfies*

$$\check{C}_{n;i}^{(a)}(x)W(x)r_j^{(b)}(x) = \begin{cases} \left\langle \check{P}_n^{[1]}(x), (\xi_i^{[a]})_x \eta_i^{(a)} \begin{bmatrix} (x-x_a)^{\kappa_{\max(i,j)}^{(a)} - \kappa_i^{(a)}} \\ \vdots \\ (x-x_a)^{\kappa_{\max(i,j)}^{(a)} - 1} \end{bmatrix} \right\rangle w_{i,j}^{(a)}(x) \\ + (x-x_a)^{\kappa_j^{(a)}} T^{(a,a)}(x) \quad \text{if } a = b, \\ (x-x_b)^{\kappa_j^{(b)}} T^{(a,b)}(x) \quad \text{if } a \neq b, \end{cases} \tag{34}$$

where the \mathbb{C}^p -valued function $T^{(a,b)}(x)$ is analytic at $x = x_b$ and, in particular, $T^{(a,a)}(x) \in \mathbb{C}^p[x]$.

Proof. First, for the function $\check{C}_{n;i}^{(a)}(x)W(x)r_j^{(b)}(x) \in \mathbb{C}^p[x]$, with $a \neq b$, we have

$$\begin{aligned} \check{C}_{n;i}^{(a)}(x)W(x)r_j^{(b)}(x) &= \left\langle \check{P}_n^{[1]}(x), (\xi_i^{[a]})_x \eta_i^{(a)} \mathcal{L}_i^{(a)} \begin{bmatrix} \frac{I_p}{(x-x_a)^{\kappa_i^{(a)}}} \\ \vdots \\ \frac{I_p}{x-x_a} \end{bmatrix} \right\rangle W(x)r_j^{(b)}(x) \\ &= (x-x_b)^{\kappa_j^{(b)}} T^{(a,b)}(x), \end{aligned}$$

where the \mathbb{C}^p -valued function $T^{(a,b)}(x)$ is analytic at $x = x_b$. Second, from (31) and Lemma 4 we deduce that

$$\begin{aligned} \check{C}_{n;i}^{(a)}(x)W(x)r_j^{(a)}(x) &= \left\langle \check{P}_n^{[1]}(x), (\xi_i^{[a]})_x \eta_i^{(a)} \mathcal{L}_i^{(a)} \begin{bmatrix} \frac{I_p}{(x-x_a)^{\kappa_i^{(a)}}} \\ \vdots \\ \frac{I_p}{x-x_a} \end{bmatrix} \right\rangle W(x)r_j^{(a)}(x) \\ &= \left\langle \check{P}_n^{[1]}(x), (\xi_i^{[a]})_x \eta_i^{(a)} \begin{bmatrix} \frac{I_p}{(x-x_a)^{\kappa_i^{(a)}}} \\ \vdots \\ \frac{I_p}{x-x_a} \end{bmatrix} \right\rangle l_i^{(a)}(x)W(x)r_j^{(a)}(x) \\ &\quad + (x-x_a)^{\kappa_j^{(a)}} \left\langle \check{P}_n^{[1]}(x), (\xi_i^{[a]})_x \eta_i^{(a)} \right\rangle T^{(a,a)}(x), \end{aligned}$$

for some $T^{(a,a)}(x) \in \mathbb{C}^p[x]$. Therefore, from Proposition 11 we get

$$\begin{aligned} \check{C}_{n;i}^{(a)}(x)W(x)r_j^{(a)}(x) &= \left\langle \check{P}_n^{[1]}(x), (\xi_i^{[a]})_x \eta_i^{(a)} \begin{bmatrix} (x-x_a)^{\kappa_{\max(i,j)}^{(a)} - \kappa_i^{(a)}} \\ \vdots \\ (x-x_a)^{\kappa_{\max(i,j)}^{(a)} - 1} \end{bmatrix} \right\rangle \\ &\times (w_{i,j;0}^{(a)} + w_{i,j;1}^{(a)}(x-x_a) + \dots + w_{i,j;\kappa_{\min(i,j)}^{(a)}+N-2}^{(a)}) \\ &\times (x-x_a)^{\kappa_{\min(i,j)}^{(a)}+N-2} \\ &+ (x-x_a)^{\kappa_j^{(a)}} \left\langle \check{P}_n^{[1]}(x), (\xi_i^{[a]})_x \eta_i^{(a)} T^{(a,a)}(x) \right\rangle, \end{aligned}$$

and the result follows. □

We evaluate now the spectral jets of the second kind functions $\check{C}^{[1]}(z)$ á la Cauchy, thus we must take limits of derivatives precisely in points of the spectrum of $W(x)$, which do not lay in the region of definition but on the border of it. Notice that these operations are not available for the second kind functions á la Gram.

Lemma 6. For $m = 0, \dots, \kappa_j^{(a)} - 1$, the following relations hold:

$$(\check{C}_n^{[1]}(z)W(z)r_j^{(a)}(z))_{x_a}^{(m)} = \sum_{i=1}^{s_a} (\check{C}_{n;i}^{(a)}(z)W(z)r_j^{(a)}(z))_{x_a}^{(m)}. \tag{35}$$

Proof. For $z \notin \text{supp}_y(u) \cup \sigma(W(y))$, a consequence of Proposition 29 is that

$$\begin{aligned} (\check{C}_n^{[1]}(z)W(z)r_j^{(a)}(z))_{x_a}^{(m)} &= \left\langle \check{P}_n^{[1]}(x), \frac{I_p}{z-y} \right\rangle_{uW^{-1}} W(z)r_j^{(a)}(z) \Big|_{x_a}^{(m)} \\ &+ \sum_{b=1}^q \sum_{i=1}^{s_b} (\check{C}_{n;i}^{(b)}(z)W(z)r_j^{(a)}(z))_{x_a}^{(m)}. \end{aligned}$$

But, as $\sigma(W(y)) \cap \text{supp}_y(u) = \emptyset$, the derivatives of the Cauchy kernel $1/(z-y)$ are analytic functions at $z = x_a$. Therefore,

$$\begin{aligned} \left\langle \check{P}_n^{[1]}(x), \frac{I_p}{z-y} \right\rangle_{uW^{-1}} W(z)r_j^{(a)}(z) \Big|_{x_a}^{(m)} &= \left\langle \check{P}_n^{[1]}(x), \left(\frac{W(z)r_j^{(a)}(z)}{z-y} \right)_{x_a}^{(m)} \right\rangle_{uW^{-1}} \\ &= \left\langle \check{P}_n^{[1]}(x), \sum_{k=0}^m \binom{m}{k} (W(z)r_j^{(a)}(z))_{x_a}^{(k)} \right. \\ &\quad \left. \times \frac{(-1)^{m-k} (m-k)!}{(x_a-y)^{m-k+1}} \right\rangle_{uW^{-1}} \\ &= 0_{p \times 1}, \end{aligned}$$

for $m = 0, \dots, \kappa_j^{(a)} - 1$. Equation (34) shows that $\check{C}_{n;i}^{(b)}(x)W(x)r_j^{(a)}(x)$ for $b \neq a$ has a zero at $z = x_a$ of order $\kappa_j^{(a)}$ and, consequently,

$$\left(\check{C}_{n;i}^{(b)}(x)W(x)r_j^{(a)}(x)\right)_{x_a}^{(m)} = 0, \quad b \neq a,$$

for $m = 0, \dots, \kappa_j^{(a)} - 1$. □

Definition 31. Given the functions $w_{i,j;k}^{(a)}$ introduced in Proposition 11, let us introduce the matrix $\mathcal{W}_{j,i}^{(a)} \in \mathbb{C}^{\kappa_j^{(a)} \times \kappa_i^{(a)}}$

$$\mathcal{W}_{j,i}^{(a)} := \begin{cases} \eta_j^{(a)} \begin{bmatrix} w_{i,j;0}^{(a)} & w_{i,j;1}^{(a)} & \cdots & w_{i,j;\kappa_j^{(a)}-1}^{(a)} \\ 0_p & w_{i,j;0}^{(a)} & \cdots & w_{i,j;\kappa_j^{(a)}-2}^{(a)} \\ \vdots & & \ddots & \vdots \\ 0_p & 0_p & & w_{i,j;0}^{(a)} \end{bmatrix}, & i \geq j, \\ \eta_j^{(a)} \begin{bmatrix} w_{i,j;\kappa_j^{(a)}-\kappa_i^{(a)}}^{(a)} & w_{i,j;\kappa_j^{(a)}-\kappa_i^{(a)}+1}^{(a)} & \cdots & w_{i,j,\kappa_j^{(a)}-1}^{(a)} \\ \vdots & & & \vdots \\ w_{i,j;0}^{(a)} & w_{i,j;1}^{(a)} & \cdots & w_{i,j,\kappa_i^{(a)}-1}^{(a)} \\ 0_p & w_{i,j;0}^{(a)} & & \vdots \\ \vdots & & \ddots & \\ 0_p & 0_p & & w_{i,j;0}^{(a)} \end{bmatrix}, & i \leq j, \end{cases}$$

and the matrix $\mathcal{W}_j^{(a)} \in \mathbb{C}^{\kappa_j^{(a)} \times \alpha_a}$ given by

$$\mathcal{W}_j^{(a)} := [\mathcal{W}_{j,1}^{(a)}, \dots, \mathcal{W}_{j,s_a}^{(a)}].$$

We also consider the matrices $\mathcal{W}^{(a)} \in \mathbb{C}^{\alpha_a \times \alpha_a}$ and $\mathcal{W} \in \mathbb{C}^{Np \times Np}$

$$\mathcal{W}^{(a)} := \begin{bmatrix} \mathcal{W}_1^{(a)} \\ \vdots \\ \mathcal{W}_{s_a}^{(a)} \end{bmatrix}, \quad \mathcal{W} := \text{diag}(\mathcal{W}^{(1)}, \dots, \mathcal{W}^{(g)}). \tag{36}$$

Proposition 30. *The following relations among the spectral jets, introduced in Definition 8, of the perturbed polynomials and second kind functions*

$$\mathcal{J}_{\check{C}_n^{[1]}W}^{(j)}(x_a) = \sum_{i=1}^{s_a} \mathcal{J}_{\check{C}_{n;i}W}^{(j)}(x_a), \quad \mathcal{J}_{\check{C}_n^{[1]}W}(x_a) = \sum_{i=1}^{s_a} \mathcal{J}_{\check{C}_{n;i}W}(x_a), \tag{37}$$

$$\mathcal{J}_{\check{C}_{n;i}W}^{(j)}(x_a) = \langle \check{P}_n^{[1]}(x), (\xi_i^{[a]})_x \rangle \mathcal{W}_{i,j}^{(a)}, \quad \mathcal{J}_{\check{C}_{n;i}W}(x_a) = \langle \check{P}_n^{[1]}(x), (\xi_i^{[a]})_x \rangle \mathcal{W}_i^{(a)}, \tag{38}$$

$$\mathcal{J}_{\check{C}_n^{[1]}W}(x_a) = \langle \check{P}_n^{[1]}(x), (\xi^{[a]})_x \rangle \mathcal{W}^{(a)}, \quad \mathcal{J}_{\check{C}_n^{[1]}W} = \langle \check{P}_n^{[1]}(x), (\xi)_x \rangle \mathcal{W}, \tag{39}$$

are satisfied.

Proof. Equation (37) is a direct consequence of (35). According to (34) for $m = 0, \dots, \kappa_j^{(a)} - 1$, we have

$$\begin{aligned} & (\check{C}_{n;i}^{(a)}(x)W(x)r_j^{(a)}(x))_{x=x_a}^{(m)} \\ &= \langle \check{P}_n^{[1]}(x), (\xi_i^{[a]})_x \rangle \eta_i^{(a)} \begin{bmatrix} ((x - x_a)^{\kappa_{\max(i,j)}^{(a)} - \kappa_i^{(a)}} w_{i,j}^{(a)}(x))_{x_a}^{(m)} \\ \vdots \\ ((x - x_a)^{\kappa_{\max(i,j)}^{(a)} - 1} w_{i,j}^{(a)}(x))_{x_a}^{(m)} \end{bmatrix}, \end{aligned}$$

and collecting all these equations in a matrix form we get (38). Finally, we notice that from (37) and (38) we deduce

$$\begin{aligned} \mathcal{J}_{\check{C}_n^{[1]}W}^{(j)}(x_a) &= \sum_{i=1}^{s_a} \langle \check{P}_n^{[1]}(x), (\xi_i^{[a]})_x \rangle \mathcal{W}_{i,j}^{(a)}, \\ \mathcal{J}_{\check{C}_n^{[1]}W}(x_a) &= \sum_{i=1}^{s_a} \langle \check{P}_n^{[1]}(x), (\xi_i^{[a]})_x \rangle \mathcal{W}_i^{(a)}. \end{aligned}$$

Now, using (36) we can write the second equation as

$$\begin{aligned} \mathcal{J}_{\check{C}_n^{[1]}W}(x_a) &= \sum_{i=1}^{s_a} \langle \check{P}_n^{[1]}(x), (\xi_i^{[a]})_x \rangle \mathcal{W}_i^{(a)} \\ &= \langle \check{P}_n^{[1]}(x), (\xi^{[a]})_x \rangle \mathcal{W}^{(a)}. \end{aligned}$$

A similar argument leads to the second relation in (39). □

Definition 32. For the Hankel masses, we also consider the matrices $\mathcal{T}_i^{(a)} \in \mathbb{C}^{p\kappa_i^{(a)} \times \alpha_a}$, $\mathcal{T}^{(a)} \in \mathbb{C}^{p\alpha_a \times \alpha_a}$ and $\mathcal{T} \in \mathbb{C}^{Np^2 \times Np}$ given by

$$\begin{aligned} \mathcal{T}_i^{(a)} &:= \mathcal{X}_i^{(a)} \eta_i^{(a)} \mathcal{W}_i^{(a)}, \\ \mathcal{T}^{(a)} &:= \begin{bmatrix} \mathcal{T}_1^{(a)} \\ \vdots \\ \mathcal{T}_{s_a}^{(a)} \end{bmatrix}, \quad \mathcal{T} := \text{diag}(\mathcal{T}^{(1)}, \dots, \mathcal{T}^{(q)}). \end{aligned}$$

2.4. Spectral Christoffel–Geronimus formulas

Proposition 31. *If $n \geq N$, the matrix coefficients of the connection matrix satisfy*

$$[\omega_{n,n-N}, \dots, \omega_{n,n-1}] = -(\mathcal{J}_{C_n^{[1]}} - \langle P_n^{[1]}(x), (\xi)_x \rangle \mathcal{W}) \begin{bmatrix} \mathcal{J}_{C_{n-N}^{[1]}} - \langle P_{n-N}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \\ \vdots \\ \mathcal{J}_{C_{n-1}^{[1]}} - \langle P_{n-1}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \end{bmatrix}^{-1}.$$

Proof. From the connection formula (24), for $n \geq N$

$$\check{C}_n^{[1]}(x)W(x) = \sum_{k=n-N}^{n-1} \omega_{n,k}C_k^{[1]}(x) + C_n^{[1]}(x),$$

and we conclude that

$$\mathcal{J}_{\check{C}_n^{[1]}W} = [\omega_{n,n-N}, \dots, \omega_{n,n-1}] \begin{bmatrix} \mathcal{J}_{C_{n-N}^{[1]}} \\ \vdots \\ \mathcal{J}_{C_{n-1}^{[1]}} \end{bmatrix} + \mathcal{J}_{C_n^{[1]}}.$$

Similarly, using Eq. (21), we get

$$\begin{aligned} &\langle \check{P}_n^{[1]}(x), (\xi)_x \rangle \mathcal{W} \\ &= [\omega_{n,n-N}, \dots, \omega_{n,n-1}] \begin{bmatrix} \langle P_{n-N}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \\ \vdots \\ \langle P_{n-1}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \end{bmatrix} + \langle P_n^{[1]}(x), (\xi)_x \rangle \mathcal{W}. \end{aligned} \tag{40}$$

Now, from (39) we deduce

$$\begin{aligned} &[\omega_{n,n-N}, \dots, \omega_{n,n-1}] \begin{bmatrix} \mathcal{J}_{C_{n-N}^{[1]}} \\ \vdots \\ \mathcal{J}_{C_{n-1}^{[1]}} \end{bmatrix} + \mathcal{J}_{C_n^{[1]}} \\ &= [\omega_{n,n-N}, \dots, \omega_{n,n-1}] \begin{bmatrix} \langle P_{n-N}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \\ \vdots \\ \langle P_{n-1}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \end{bmatrix} + \langle P_n^{[1]}(x), (\xi)_x \rangle \mathcal{W}, \end{aligned}$$

that is to say

$$\begin{aligned}
 & [\omega_{n,n-N}, \dots, \omega_{n,n-1}] \begin{bmatrix} \mathcal{J}_{C_{n-N}^{[1]}} - \langle P_{n-N}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \\ \vdots \\ \mathcal{J}_{C_{n-1}^{[1]}} - \langle P_{n-1}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \end{bmatrix} \\
 &= -(\mathcal{J}_{C_n^{[1]}} - \langle P_n^{[1]}(x), (\xi)_x \rangle \mathcal{W}). \quad \square
 \end{aligned}$$

Remark 7. In the next results, the jets of the Christoffel–Darboux kernels are considered with respect to the first variable x , and we treat the y -variable as a parameter.

Theorem 2 (Spectral Christoffel–Geronimus formulas). *When $n \geq N$, for monic Geronimus perturbations, with masses as described in (29), we have the following last quasideterminantal expressions for the perturbed biorthogonal matrix polynomials and its matrix norms:*

$$\begin{aligned}
 \check{P}_n^{[1]}(x) &= \Theta_* \begin{bmatrix} \mathcal{J}_{C_{n-N}^{[1]}} - \langle P_{n-N}^{[1]}(x), (\xi)_x \rangle \mathcal{W} & P_{n-N}^{[1]}(x) \\ \vdots & \vdots \\ \mathcal{J}_{C_n^{[1]}} - \langle P_n^{[1]}(x), (\xi)_x \rangle \mathcal{W} & P_n^{[1]}(x) \end{bmatrix}, \\
 \check{H}_n &= \Theta_* \begin{bmatrix} \mathcal{J}_{C_{n-N}^{[1]}} - \langle P_{n-N}^{[1]}(x), (\xi)_x \rangle \mathcal{W} & H_{n-N} \\ \mathcal{J}_{C_{n-N+1}^{[1]}} - \langle P_{n-N+1}^{[1]}(x), (\xi)_x \rangle \mathcal{W} & 0_p \\ \vdots & \vdots \\ \mathcal{J}_{C_n^{[1]}} - \langle P_n^{[1]}(x), (\xi)_x \rangle \mathcal{W} & 0_p \end{bmatrix}, \\
 (\check{P}_n^{[2]}(y))^T &= -\Theta_* \begin{bmatrix} \mathcal{J}_{C_{n-N}^{[1]}} - \langle P_{n-N}^{[1]}(x), (\xi)_x \rangle \mathcal{W} & H_{n-N} \\ \mathcal{J}_{C_{n-N+1}^{[1]}} - \langle P_{n-N+1}^{[1]}(x), (\xi)_x \rangle \mathcal{W} & 0_p \\ \vdots & \vdots \\ \mathcal{J}_{C_{n-1}^{[1]}} - \langle P_{n-1}^{[1]}(x), (\xi)_x \rangle \mathcal{W} & 0_p \\ W(y)(\mathcal{J}_{K_{n-1}^{(p)}}(y) - \langle K_{n-1}(x, y), (\xi)_x \rangle \mathcal{W}) + \mathcal{J}_V(y) & 0_p \end{bmatrix}.
 \end{aligned}$$

Proof. First, we consider the expressions for $\check{P}_n^{[1]}(x)$ and \check{H}_n . Using relation (21) we have

$$\check{P}_n^{[1]}(x) = P_n^{[1]}(x) + [\omega_{n,n-N}, \dots, \omega_{n,n-1}] \begin{bmatrix} P_{n-N}^{[1]}(x) \\ \vdots \\ P_{n-1}^{[1]}(x) \end{bmatrix},$$

from Proposition 31 we obtain

$$\begin{aligned} \check{P}_n^{[1]}(x) &= P_n^{[1]}(x) - (\mathcal{J}_{C_n^{[1]}} - \langle P_n^{[1]}(x), (\xi)_x \rangle \mathcal{W}) \\ &\quad \times \begin{bmatrix} \mathcal{J}_{C_{n-N}^{[1]}} - \langle P_{n-N}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \\ \vdots \\ \mathcal{J}_{C_{n-1}^{[1]}} - \langle P_{n-1}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \end{bmatrix}^{-1} \begin{bmatrix} P_{n-N}^{[1]}(x) \\ \vdots \\ P_{n-1}^{[1]}(x) \end{bmatrix}, \end{aligned}$$

and the result follows. To get the transformation for the H 's we proceed as follows. From (20) we deduce

$$\check{H}_n = \omega_{n,n-N} H_{n-N}. \tag{41}$$

But, according to Proposition 31, we have

$$\omega_{n,n-N} = -(\mathcal{J}_{C_n^{[1]}} - \langle P_n^{[1]}(x), (\xi)_x \rangle \mathcal{W}) \begin{bmatrix} \mathcal{J}_{C_{n-N}^{[1]}} - \langle P_{n-N}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \\ \vdots \\ \mathcal{J}_{C_{n-1}^{[1]}} - \langle P_{n-1}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \end{bmatrix}^{-1} \begin{bmatrix} I_p \\ 0_p \\ \vdots \\ 0_p \end{bmatrix}.$$

Hence,

$$\check{H}_n = -(\mathcal{J}_{C_n^{[1]}} - \langle P_n^{[1]}(x), (\xi)_x \rangle \mathcal{W}) \begin{bmatrix} \mathcal{J}_{C_{n-N}^{[1]}} - \langle P_{n-N}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \\ \vdots \\ \mathcal{J}_{C_{n-1}^{[1]}} - \langle P_{n-1}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \end{bmatrix}^{-1} \begin{bmatrix} H_{n-N} \\ 0_p \\ \vdots \\ 0_p \end{bmatrix}.$$

We now prove the result for $(\check{P}_n^{[2]}(y))^\top$. On one hand, according to Definition 12 we rewrite (28) as

$$\begin{aligned} \sum_{k=0}^{n-1} (\check{P}_k^{[2]}(y))^\top \check{H}_k^{-1} \check{C}_k^{[1]}(x) W(x) &= W(y) K_{n-1}^{(pc)}(x, y) \\ &\quad - [(\check{P}_n^{[2]}(y))^\top \check{H}_n^{-1}, \dots, (\check{P}_{n+N-1}^{[2]}(y))^\top \check{H}_{n+N-1}^{-1}] \\ &\quad \times \Omega[n] \begin{bmatrix} C_{n-N}^{[1]}(x) \\ \vdots \\ C_{n-1}^{[1]}(x) \end{bmatrix} + \mathcal{V}(x, y). \end{aligned}$$

Therefore, the corresponding spectral jets do satisfy

$$\sum_{k=0}^{n-1} (\check{P}_k^{[2]}(y))^\top \check{H}_k^{-1} \mathcal{J}_{\check{C}_k^{[1]}W} = W(y) \mathcal{J}_{K_{n-1}^{(pc)}}(y) - [(\check{P}_n^{[2]}(y))^\top \check{H}_n^{-1}, \dots, (\check{P}_{n+N-1}^{[2]}(y))^\top \check{H}_{n+N-1}^{-1}] \times \Omega[n] \begin{bmatrix} \mathcal{J}_{C_{n-N}^{[1]}} \\ \vdots \\ \mathcal{J}_{C_{n-1}^{[1]}} \end{bmatrix} + \mathcal{J}_\nu(y),$$

and, recalling (39), we conclude that

$$\sum_{k=0}^{n-1} (\check{P}_k^{[2]}(y))^\top \check{H}_k^{-1} \langle \check{P}_k^{[1]}(x), (\xi)_x \rangle \mathcal{W} = W(y) \mathcal{J}_{K_{n-1}^{(pc)}}(y) - [(\check{P}_n^{[2]}(y))^\top \check{H}_n^{-1}, \dots, (\check{P}_{n+N-1}^{[2]}(y))^\top \check{H}_{n+N-1}^{-1}] \times \Omega[n] \begin{bmatrix} \mathcal{J}_{C_{n-N}^{[1]}} \\ \vdots \\ \mathcal{J}_{C_{n-1}^{[1]}} \end{bmatrix} + \mathcal{J}_\nu(y). \tag{42}$$

On the other hand, from (27) we realize that

$$\sum_{k=0}^{n-1} (\check{P}_k^{[2]}(y))^\top \check{H}_k^{-1} \langle P_k^{[1]}(x), (\xi)_x \rangle \mathcal{W} = W(y) \langle K_{n-1}(x, y), (\xi)_x \rangle \mathcal{W} - [(\check{P}_n^{[2]}(y))^\top \check{H}_n^{-1}, \dots, (\check{P}_{n+N-1}^{[2]}(y))^\top \check{H}_{n+N-1}^{-1}] \Omega[n] \begin{bmatrix} \langle P_{n-N}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \\ \vdots \\ \langle P_{n-1}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \end{bmatrix},$$

which can be subtracted to (42) to get

$$W(y) (\mathcal{J}_{K_{n-1}^{(pc)}}(y) - \langle K_{n-1}(x, y), (\xi)_x \rangle \mathcal{W}) + \mathcal{J}_\nu(y) = [(\check{P}_n^{[2]}(y))^\top \check{H}_n^{-1}, \dots, (\check{P}_{n+N-1}^{[2]}(y))^\top \check{H}_{n+N-1}^{-1}] \times \Omega[n] \begin{bmatrix} \mathcal{J}_{C_{n-N}^{[1]}} - \langle P_{n-N}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \\ \vdots \\ \mathcal{J}_{C_{n-1}^{[1]}} - \langle P_{n-1}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \end{bmatrix}.$$

Hence, we obtain the formula

$$\begin{aligned}
 & [(\check{P}_n^{[2]}(y))^\top \check{H}_n^{-1}, \dots, (\check{P}_{n+N-1}^{[2]}(y))^\top \check{H}_{n+N-1}^{-1}] \Omega[n] \\
 &= (W(y)(\mathcal{J}_{K_{n-1}^{(pc)}}(y) - \langle K_{n-1}(x, y), (\xi)_x \rangle \mathcal{W}) + \mathcal{J}_V(y)) \\
 &\quad \times \begin{bmatrix} \mathcal{J}_{C_{n-N}^{[1]}} - \langle P_{n-N}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \\ \vdots \\ \mathcal{J}_{C_{n-1}^{[1]}} - \langle P_{n-1}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \end{bmatrix}^{-1}. \tag{43}
 \end{aligned}$$

Now, for $n \geq N$, from Definition 26 and the fact that $\omega_{n,n-N} = \check{H}_n(H_{n-N})^{-1}$, we get

$$\begin{aligned}
 (\check{P}_n^{[2]}(y))^\top &= (W(y)(\mathcal{J}_{K_{n-1}^{(pc)}}(y) - \langle K_{n-1}(x, y), (\xi)_x \rangle \mathcal{W}) \\
 &\quad + \mathcal{J}_V(y)) \begin{bmatrix} \mathcal{J}_{C_{n-N}^{[1]}} - \langle P_{n-N}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \\ \vdots \\ \mathcal{J}_{C_{n-1}^{[1]}} - \langle P_{n-1}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \end{bmatrix}^{-1} \begin{bmatrix} H_{n-N} \\ 0_p \\ \vdots \\ 0_p \end{bmatrix},
 \end{aligned}$$

and the result follows. □

2.5. Nonspectral Christoffel–Geronimus formulas

We now present an alternative *orthogonality relations* approach for the derivation of Christoffel-type formulas, that avoids the use of the second kind functions and of the spectral structure of the perturbing polynomial. A key feature of these results is that they hold even for perturbing matrix polynomials with singular leading coefficient.

Definition 33. For a given perturbed matrix of generalized kernels $\check{u}_{x,y} = u_{x,y}(W(y))^{-1} + v_{x,y}$, with $v_{x,y}W(y) = 0_p$, we define a semi-infinite block matrix

$$\begin{aligned}
 R &:= \langle P^{[1]}(x), \chi(y) \rangle_{\check{u}} \\
 &= \langle P^{[1]}(x), \chi(y) \rangle_{uW^{-1}} + \langle P^{[1]}(x), \chi(y) \rangle_v.
 \end{aligned}$$

Remark 8. Its blocks are $R_{n,l} = \langle P_n^{[1]}(x), I_p y^l \rangle_{\check{u}} \in \mathbb{C}^{p \times p}$. Observe that for a Geronimus perturbation of a Borel measure $d\mu(x, y)$, with general masses as in (29) we have

$$\begin{aligned}
 R_{n,l} &= \int P_n^{[1]}(x) d\mu(x, y) (W(y))^{-1} y^l \\
 &\quad + \sum_{a=1}^q \sum_{i=1}^{s_a} \sum_{m=0}^{\kappa_j^{(a)} - 1} \frac{1}{m!} \langle P_n^{[1]}(x), (\xi_{i,m}^{[a]})_x \rangle (l_j^{(a)}(y) y^l)_{x_a}^{(m)},
 \end{aligned}$$

that, when the masses are discrete and supported by the diagonal $y = x$, reduces to

$$R_{n,l} = \int P_n^{[1]}(x) d\mu(x, y) (W(y))^{-1} y^l + \sum_{a=1}^q \sum_{i=1}^{s_a} \sum_{m=0}^{\kappa_j^{(a)}-1} \frac{1}{m!} \left(P_n^{[1]}(x) x^l \xi_{i,m}^{[a]} l_j^{(a)}(x) \right)_{x_a}^{(m)}.$$

Proposition 32. *The following relations hold true:*

$$R = S_1 \check{G}, \tag{44}$$

$$\omega R = \check{H} (\check{S}_2)^{-\top}, \tag{45}$$

$$RW(\Lambda^\top) = H (S_2)^{-\top}. \tag{46}$$

Proof. (44) follows from Definition 33. Indeed,

$$\begin{aligned} R &= \langle P^{[1]}(x), \chi(y) \rangle_{\check{u}} \\ &= S_1 \langle \chi(x), \chi(y) \rangle_{\check{u}}. \end{aligned}$$

To deduce (45) we recall (16), (44), and the Gauss factorization of the perturbed matrix of moments

$$\begin{aligned} \omega R &= (\check{S}_1 (S_1)^{-1}) (S_1 \check{G}) \\ &= \check{S}_1 \check{G} \\ &= \check{S}_1 ((\check{S}_1)^{-1} \check{H} (\check{S}_2)^{-\top}). \end{aligned}$$

Finally, to get (46), we use (17) together with (45), which implies $\omega = \omega RW(\Lambda^\top) (S_2)^\top H^{-1}$, and as the resolvent it is unitriangular with a unique inverse matrix [14], we obtain the result. \square

From (45) it immediately follows that

Proposition 33. *The matrix R fulfills*

$$(\omega R)_{n,l} = \begin{cases} 0_p, & l \in \{0, \dots, n-1\}, \\ \check{H}_n, & n = l. \end{cases}$$

Proposition 34. *The matrix $\begin{bmatrix} R_{0,0} & \dots & R_{0,n-1} \\ \vdots & & \vdots \\ R_{n-1,0} & \dots & R_{n-1,n-1} \end{bmatrix}$ is nonsingular.*

Proof. From (44) we conclude for the corresponding truncations that $R_{[n]} = (S_1)_{[n]} \check{G}_{[n]}$ is nonsingular, as we are assuming, to ensure the orthogonality, that $\check{G}_{[n]}$ is nonsingular for all $n \in \{1, 2, \dots\}$. \square

Definition 34. Let us introduce the polynomials $r_{n,l}^K(z) \in \mathbb{C}^{p \times p}[z]$, $l \in \{0, \dots, n-1\}$, given by

$$\begin{aligned} r_{n,l}^K(z) &:= \langle W(z)K_{n-1}(x, z), I_p y^l \rangle_{\tilde{u}} - I_p z^l \\ &= \langle W(z)K_{n-1}(x, z), I_p y^l \rangle_{uW^{-1}} + \langle W(z)K_{n-1}(x, z), I_p y^l \rangle_v - I_p z^l. \end{aligned}$$

Proposition 35. For $l \in \{0, 1, \dots, n-1\}$ and $m = \min(n, N)$ we have

$$r_{n,l}^K(z) = [(\check{P}_n^{[2]}(z))^\top (\check{H}_n)^{-1}, \dots, (\check{P}_{n-1+N}^{[2]}(z))^\top (\check{H}_{n-1+N})^{-1}] \Omega[n] \begin{bmatrix} R_{n-m,l} \\ \vdots \\ R_{n-1,l} \end{bmatrix}.$$

Proof. It follows from (27), Definition 33, and (14). □

Definition 35. For $n \geq N$, given the matrix

$$\begin{bmatrix} R_{n-N,0} & \dots & R_{n-N,n-1} \\ \vdots & & \vdots \\ R_{n-1,0} & \dots & R_{n-1,n-1} \end{bmatrix} \in \mathbb{C}^{Np \times np},$$

we construct a submatrix of it by selecting Np columns among all the np columns. For that aim, we use indexes (i, a) labeling the columns, where i runs through $\{0, \dots, n-1\}$ and indicates the block, and $a \in \{1, \dots, p\}$ denotes the corresponding column in that block; i.e. (i, a) is an index selecting the a th column of the i -block. Given a set of N different couples $I = \{(i_r, a_r)\}_{r=1}^N$, with a lexicographic ordering, we define the corresponding square submatrix $R_n^\square := [\mathbf{c}_{(i_1, a_1)}, \dots, \mathbf{c}_{(i_{Np}, a_{Np})}]$. Here $\mathbf{c}_{(i_r, a_r)}$ denotes the a_r th column of the matrix

$$\begin{bmatrix} R_{n-N, i_r} \\ \vdots \\ R_{n-1, i_r} \end{bmatrix}.$$

The set of indexes I is said to be poised if R_n^\square is nonsingular. We also use the notation where $r_n^\square := [\tilde{\mathbf{c}}_{(i_1, a_1)}, \dots, \tilde{\mathbf{c}}_{(i_{Np}, a_{Np})}]$. Here $\tilde{\mathbf{c}}_{(i_r, a_r)}$ denotes the a_r th column of the matrix R_{n, i_r} . Given a poised set of indexes we define $(r_n^K(y))^\square$ as the matrix built up by taking from the matrices $r_{n, i_r}^K(y)$ the columns a_r .

Lemma 7. For $n \geq N$, there exists at least a poised set.

Proof. For $n \geq N$, we consider the rectangular block matrix

$$\begin{bmatrix} R_{n-N,0} & \dots & R_{n-N,n-1} \\ \vdots & & \vdots \\ R_{n-1,0} & \dots & R_{n-1,n-1} \end{bmatrix} \in \mathbb{C}^{Np \times np}.$$

As the truncation $R_{[n]}$ is nonsingular, this matrix is full rank, i.e. all its Np rows are linearly independent. Thus, there must be Np independent columns and the desired result follows. \square

Lemma 8. *Whenever the leading coefficient A_N of the perturbing polynomial $W(y)$ is nonsingular, we can decompose any monomial $I_p y^l$ as*

$$I_p y^l = \alpha_l(y)(W(y))^\top + \beta_l(y),$$

where $\alpha_l(y), \beta_l(y) = \beta_{l,0} + \dots + \beta_{l,N-1}y^{N-1} \in \mathbb{C}^{p \times p}[y]$, with $\deg \alpha_l(y) \leq l - N$.

Proposition 36. *Let us assume that the matrix polynomial $W(y) = A_N y^N + \dots + A_0$ has a nonsingular leading coefficient and $n \geq N$. Then, the set $\{0, 1, \dots, N - 1\}$ is poised.*

Proof. From Proposition 33 we deduce

$$[\omega_{n,n-N}, \dots, \omega_{n,n-1}] \begin{bmatrix} R_{n-N,l} \\ \vdots \\ R_{n-1,l} \end{bmatrix} = -R_{n,l},$$

for $l \in \{0, 1, \dots, n - 1\}$. In particular, the resolvent vector $[\omega_{n,n-N}, \dots, \omega_{n,n-1}]$ is a solution of the linear system

$$[\omega_{n,n-N}, \dots, \omega_{n,n-1}] \begin{bmatrix} R_{n-N,0} & \dots & R_{n-N,N-1} \\ \vdots & & \vdots \\ R_{n-1,0} & \dots & R_{n-1,N-1} \end{bmatrix} = -[R_{n,0}, \dots, R_{n,N-1}]. \quad (47)$$

We will show now that this is the unique solution to this linear system. Let us proceed by contradiction and assume that there is another solution, say $[\tilde{\omega}_{n,n-N}, \dots, \tilde{\omega}_{n,n-1}]$. Consider then the monic matrix polynomial

$$\tilde{P}_n(x) = P_n^{[1]}(x) + \tilde{\omega}_{n,n-1}P_{n-1}^{[1]}(x) + \dots + \tilde{\omega}_{n,n-N}P_{n-N}^{[1]}(x).$$

Because $[\tilde{\omega}_{n,n-N}, \dots, \tilde{\omega}_{n,n-1}]$ solves (47) we know that

$$\langle \tilde{P}_n(x), I_p y^l \rangle_{\tilde{u}} = 0_p, \quad l \in \{0, \dots, N - 1\}.$$

Lemma 8 implies the following relations for $\deg \alpha_l(y) < m$:

$$\begin{aligned} \langle P_m^{[1]}(x), I_p y^l \rangle_{\tilde{u}} &= \langle P_m^{[1]}(x), \alpha_l(y) \rangle_{\tilde{u}W} + \langle P_m^{[1]}(x), \beta_l(y) \rangle_{\tilde{u}} \\ &= \langle P_m^{[1]}(x), \alpha_l(y) \rangle_{\tilde{u}} + \langle P_m^{[1]}(x), \beta_l(y) \rangle_{\tilde{u}} \\ &= \langle P_m^{[1]}(x), \beta_l(y) \rangle_{\tilde{u}}. \end{aligned}$$

But $\deg \alpha_l(y) \leq l - N$, so that the previous equation will hold at least for $l - N < m$; i.e. $l < m + N$. Consequently, for $l \in \{0, \dots, n - 1\}$, we find

$$\begin{aligned} \langle \tilde{P}_n(x), I_p y^l \rangle_{\tilde{u}} &= \langle P_n^{[1]}(x), I_p y^l \rangle_{\tilde{u}} + \tilde{\omega}_{n,n-1} \langle P_{n-1}^{[1]}(x), I_p y^l \rangle_{\tilde{u}} + \dots \\ &\quad + \tilde{\omega}_{n,n-N} \langle P_{n-N}^{[1]}(y), I_p y^l \rangle_{\tilde{u}} \\ &= \langle P_n^{[1]}(x), \beta_l(y) \rangle_{\tilde{u}} + \tilde{\omega}_{n,n-1} \langle P_{n-1}^{[1]}(x), \beta_l(y) \rangle_{\tilde{u}} + \dots \\ &\quad + \tilde{\omega}_{n,n-N} \langle P_{n-N}^{[1]}(x), \beta_l(y) \rangle_{\tilde{u}} \\ &= \sum_{k=0}^{N-1} (R_{n,k} + \tilde{\omega}_{n,n-1} R_{n-1,k} + \dots + \tilde{\omega}_{n,n-N} R_{n-N,k}) (\beta_{l,k})^\top \\ &= 0_p. \end{aligned}$$

Therefore, from the uniqueness of the biorthogonal families, we deduce $\tilde{P}_n(x) = \check{P}_n^{[1]}(x)$, and, recalling (21), there is a unique solution of (47). Thus,

$$\begin{bmatrix} R_{n-N,0} & \dots & R_{n-N,n-1} \\ \vdots & & \vdots \\ R_{n-1,0} & \dots & R_{n-1,n-1} \end{bmatrix}$$

is nonsingular, and $I = \{0, \dots, N - 1\}$ is a poised set. □

Proposition 37. *For $n \geq N$, given a poised set, which always exists, we have*

$$[\omega_{n,n-N}, \dots, \omega_{n,n-1}] = -r_n^\square (R_n^\square)^{-1}.$$

Proof. It follows from Proposition 33. □

Theorem 3 (Non-spectral Christoffel–Geronimus formulas). *Given a matrix Geronimus transformation the corresponding perturbed polynomials, $\{\check{P}_n^{[1]}(x)\}_{n=0}^\infty$ and $\{\check{P}_n^{[2]}(y)\}_{n=0}^\infty$, and matrix norms $\{\check{H}_n\}_{n=0}^\infty$ can be expressed as follows. For $n \geq N$,*

$$\check{P}_n^{[1]}(x) = \Theta_* \left[\begin{array}{c|c} R_n^\square & \begin{array}{c} P_{n-N}^{[1]}(x) \\ \vdots \\ P_{n-1}^{[1]}(x) \end{array} \\ \hline r_n^\square & P_n^{[1]}(x) \end{array} \right], \quad (\check{P}_n^{[2]}(y))^\top A_N = -\Theta_* \left[\begin{array}{c|c} R_n^\square & \begin{array}{c} H_{n-N} \\ 0_p \\ \vdots \\ 0_p \end{array} \\ \hline (r_n^K(y))^\square & 0_p \end{array} \right],$$

and two alternative expressions

$$\check{H}_n = \Theta_* \left[\begin{array}{c|c} R_{n-N,n} & \\ \hline R_n^\square & \vdots \\ \hline r_n^\square & R_{n,n} \end{array} \right] = \Theta_* \left[\begin{array}{c|c} H_{n-N} & \\ \hline R_n^\square & 0_p \\ \hline r_n^\square & 0_p \end{array} \right].$$

Proof. For $m = \min(n, N)$, from the connection formula (18) we have

$$\check{P}_n^{[1]}(x) = [\omega_{n,n-m}, \dots, \omega_{n,n-1}] \begin{bmatrix} P_{n-m}^{[1]}(x) \\ \vdots \\ P_{n-1}^{[1]}(x) \end{bmatrix} + P_n^{[1]}(x),$$

and from Proposition 33 we deduce

$$\check{H}_n = [\omega_{n,n-m}, \dots, \omega_{n,n-1}] \begin{bmatrix} R_{n-m,n} \\ \vdots \\ R_{n-1,n} \end{bmatrix} + R_{n,n},$$

and use (41). Then, recalling Proposition 37 we obtain the desired formulas for $\check{P}_n^{[1]}(x)$ and \check{H}_n .

For $n \geq N$, we have

$$r_{n,l}^K(y) = [(\check{P}_n^{[2]}(y))^\top (\check{H}_n)^{-1}, \dots, (\check{P}_{n-1+N}^{[2]}(y))^\top (\check{H}_{n-1+N})^{-1}] \Omega[n] \begin{bmatrix} R_{n-N,l} \\ \vdots \\ R_{n-1,l} \end{bmatrix},$$

so that

$$(r_n^K(y))^\square (R_n^\square)^{-1} = [(\check{P}_n^{[2]}(y))^\top (\check{H}_n)^{-1}, \dots, (\check{P}_{n-1+N}^{[2]}(y))^\top (\check{H}_{n-1+N})^{-1}] \Omega[n].$$

In particular, recalling (20), we deduce that

$$(\check{P}_n^{[2]}(y))^\top A_N = (r_n^K(y))^\square (R_n^\square)^{-1} \begin{bmatrix} H_{n-N} \\ 0_p \\ \vdots \\ 0_p \end{bmatrix}. \quad \square$$

2.6. Spectral versus nonspectral

Definition 36. We introduce the truncation given by taking only the first N columns of a given semi-infinite matrix

$$R^{(N)} := \begin{bmatrix} R_{0,0} & R_{0,1} & \dots & R_{0,N-1} \\ R_{1,0} & R_{0,1} & \dots & R_{1,N-1} \\ \vdots & \vdots & & \vdots \end{bmatrix}.$$

Then, we can connect the spectral methods and the nonspectral techniques as follows

Proposition 38. *The following relation takes place*

$$\mathcal{J}_{C^{[1]}} - \langle P^{[1]}(x), (\xi)_x \rangle \mathcal{W} = -R^{(N)} \mathcal{B}\mathcal{Q}.$$

Proof. From (24) we deduce that

$$\check{C}^{[1]}(x)W(x) - \check{H}(\check{S}_2)^{-\top} \begin{bmatrix} \mathcal{B}(\chi(x))_{[N]} \\ 0_p \\ \vdots \end{bmatrix} = \omega C^{[1]}(x).$$

Taking the corresponding root spectral jets, we obtain

$$\mathcal{J}_{\check{C}^{[1]}W} - \check{H}(\check{S}_2)^{-\top} \begin{bmatrix} \mathcal{B}\mathcal{Q} \\ 0_p \\ \vdots \end{bmatrix} = \omega \mathcal{J}_{C^{[1]}},$$

that, together with (39), gives

$$\omega(\mathcal{J}_{C^{[1]}} - \langle P^{[1]}(x), (\xi)_x \rangle \mathcal{W}) = -\check{H}(\check{S}_2)^{-\top} \begin{bmatrix} \mathcal{B}\mathcal{Q} \\ 0_p \\ \vdots \end{bmatrix}.$$

Now, relation (45) implies

$$\omega(\mathcal{J}_{C^{[1]}} - \langle P^{[1]}(x), (\xi)_x \rangle \mathcal{W} + R^{(N)} \mathcal{B}\mathcal{Q}) = 0.$$

But, given that ω is a lower unitriangular matrix, and therefore with an inverse, see [14], the unique solution to $\omega X = 0$, where X is a semi-infinite matrix, is $X = 0$.

□

We now discuss an important fact, which ensures that the spectral Christoffel–Geronimus formulas presented in previous sections make sense.

Corollary 1. *If the leading coefficient A_N is nonsingular and $n \geq N$, then*

$$\begin{bmatrix} \mathcal{J}_{C_{n-N}^{[1]}} - \langle P_{n-N}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \\ \vdots \\ \mathcal{J}_{C_{n-1}^{[1]}} - \langle P_{n-1}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \end{bmatrix}$$

is nonsingular.

Proof. From Proposition 38 one deduces the following formula:

$$\begin{bmatrix} \mathcal{J}_{C_{n-N}^{[1]}} - \langle P_{n-N}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \\ \vdots \\ \mathcal{J}_{C_{n-1}^{[1]}} - \langle P_{n-1}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \end{bmatrix} = - \begin{bmatrix} R_{n-N,0} & \dots & R_{n-N,N-1} \\ \vdots & & \vdots \\ R_{n-1,0} & \dots & R_{n-1,N-1} \end{bmatrix} \mathcal{BQ}. \quad (48)$$

Now, Proposition 36 and Lemma 2 lead to the result. □

We stress at this point that (48) connects the spectral and the nonspectral methods. Moreover, when we border with a further block row we obtain

$$\begin{bmatrix} \mathcal{J}_{C_{n-N}^{[1]}} - \langle P_{n-N}^{[1]}(x), (\xi)_x \rangle \mathcal{W} \\ \vdots \\ \mathcal{J}_{C_n^{[1]}} - \langle P_n^{[1]}(x), (\xi)_x \rangle \mathcal{W} \end{bmatrix} = - \begin{bmatrix} R_{n-N,0} & \dots & R_{n-N,N-1} \\ \vdots & & \vdots \\ R_{n,0} & \dots & R_{n,N-1} \end{bmatrix} \mathcal{BQ}.$$

2.7. Applications

2.7.1. Unimodular Christoffel perturbations and nonspectral techniques

The spectral methods apply to those Geronimus transformations with a perturbing polynomial $W(y)$ having a nonsingular leading coefficient A_N . This was also the case for the techniques developed in [2] for matrix Christoffel transformations, where the perturbing polynomial had a nonsingular leading coefficient. However, we have shown that despite we can extend the use of the spectral techniques to the study of matrix Geronimus transformations, we also have a nonspectral approach applicable even for singular leading coefficients. For example, some cases that have appeared several times in the literature — see [21] — are unimodular perturbations and, consequently, with $W(y)$ having a singular leading coefficient. In this case, we have that $(W(y))^{-1}$ is a matrix polynomial, and we can consider the Geronimus transformation associated with the matrix polynomial $(W(y))^{-1}$ — as the spectrum

is empty $\sigma(W(y)) = \emptyset$, no masses appear — as a Christoffel transformation with perturbing matrix polynomial $W(y)$ of the original matrix of generalized kernels

$$\check{u}_{x,y} = u_{x,y}((W(y))^{-1})^{-1} = u_{x,y}W(y). \tag{49}$$

We can apply Theorem 3 with

$$R = \langle P_n^{[1]}(x), \chi(y) \rangle_{uW}, \quad R_{n,l} = \langle P_n^{[1]}(x), I_p y^l \rangle_{uW} \in \mathbb{C}^{p \times p}.$$

For example, when the matrix of generalized kernels is a matrix of measures μ , we can write

$$R_{n,l} = \int P_n^{[1]}(x) d\mu(x, y) W(y) y^l.$$

Here $W(x)$ is a Christoffel perturbation and $\deg((W(x))^{-1})$ gives you the number of original orthogonal polynomials required for the Christoffel-type formula. Theorem 3 can be nicely applied to get $\check{P}_n^{[1]}(x)$ and \check{H}_n . However, it only gives Christoffel–Geronimus formulas for $(\check{P}_n^{[2]}(y))^\top A_N$ and given that A_N is singular, we only partially recover $\check{P}_n^{[2]}(y)$. This problem disappears whenever we have symmetric generalized kernels $u_{x,y} = (u_{y,x})^\top$, see Remark 3, as then $P_n^{[1]}(x) = P_n^{[2]}(x) =: P_n(x)$ and biorthogonality collapses to orthogonality of $\{P_n(x)\}_{n=0}^\infty$. From (49), we need to require

$$u_{x,y}W(y) = (W(x))^\top (u_{y,x})^\top,$$

that when the initial matrix of kernels is itself symmetric $u_{x,y} = (u_{y,x})^\top$ reads $u_{x,y}W(y) = (W(x))^\top u_{x,y}$. Now, if we are dealing with Hankel matrices of generalized kernels $u_{x,y} = u_{x,x}$ we find $u_{x,x}W(x) = (W(x))^\top u_{x,x}$, that for the scalar case reads $u_{x,x} = u_0 I_p$ with u_0 a generalized function we need $W(x)$ to be a symmetric matrix polynomial. For this scenario, if $\{p_n(x)\}_{n=0}^\infty$ denotes the set of monic orthogonal polynomials associated with u_0 , we have $R_{n,l} = \langle u_0, p_n(x)W(x)x^l \rangle$.

For example, if we take $p = 2$, with the unimodular perturbation given by

$$W(x) = \begin{bmatrix} (A_2)_{1,1}x^2 + (A_1)_{1,1}x + (A_0)_{1,1} & (A_2)_{1,2}x^2 + (A_1)_{1,2}x + (A_0)_{1,2} \\ (A_2)_{1,2}x^2 + (A_1)_{1,2}x + (A_0)_{1,2} & (A_2)_{2,2}x^2 + (A_1)_{2,2}x + (A_0)_{2,2} \end{bmatrix}$$

we have, that the inverse is the following matrix polynomial:

$$(W(x))^{-1} = \frac{1}{\det W(x)} \times \begin{bmatrix} (A_2)_{2,2}x^2 + (A_1)_{2,2}x + (A_0)_{2,2} & -(A_2)_{1,2}x^2 - (A_1)_{1,2}x - (A_0)_{1,2} \\ -(A_2)_{1,2}x^2 - (A_1)_{1,2}x - (A_0)_{1,2} & (A_2)_{1,1}x^2 + (A_1)_{1,1}x + (A_0)_{1,1} \end{bmatrix},$$

where $\det W(x)$ is a constant, and the inverse has also degree 2. Therefore, for $n \in \{2, 3, \dots\}$, we have the following expressions for the perturbed matrix orthogonal

polynomials

$$\check{P}_n(x) = \Theta_* \begin{bmatrix} \langle u_0, p_{n-2}(x)x^k(A_2x^2 + A_1x + A_0) \rangle \\ \langle u_0, p_{n-1}(x)x^k(A_2x^2 + A_1x) \rangle \\ \langle u_0, p_n(x)x^k A_2x^2 \rangle \\ \langle u_0, p_{n-2}(x)x^l(A_2x^2 + A_1x + A_0) \rangle & p_{n-2}(x)I_p \\ \langle u_0, p_{n-1}(x)x^l(A_2x^2 + A_1x + A_0) \rangle & p_{n-1}(x)I_p \\ \langle u_0, p_n(x)x^l(A_2x^2 + A_1x) \rangle & p_n(x)I_p \end{bmatrix},$$

and the corresponding matrix norms or quasi-tau matrices are

$$\check{H}_n = \Theta_* \begin{bmatrix} \langle u_0, p_{n-2}(x)x^k(A_2x^2 + A_1x + A_0) \rangle & \langle u_0, p_{n-2}(x)x^l(A_2x^2 + A_1x + A_0) \rangle \\ \langle u_0, p_{n-1}(x)x^k(A_2x^2 + A_1x) \rangle & \langle u_0, p_{n-1}(x)x^l(A_2x^2 + A_1x + A_0) \rangle \\ \langle u_0, p_n(x)x^k A_2x^2 \rangle & \langle u_0, p_n(x)x^l(A_2x^2 + A_1x) \rangle \\ \langle u_0, p_{n-2}(x)x^n(A_2x^2 + A_1x + A_0) \rangle \\ \langle u_0, p_{n-1}(x)x^n(A_2x^2 + A_1x + A_0) \rangle \\ \langle u_0, p_n(x)x^n(A_2x^2 + A_1x + A_0) \rangle \end{bmatrix}.$$

Here the natural numbers k and l satisfy $0 \leq k < l \leq n - 1$ and are among those (we know that they do exist) that fulfil

$$\det \begin{bmatrix} \langle u_0, p_{n-2}(x)x^k(A_2x^2 + A_1x + A_0) \rangle & \langle u_0, p_{n-2}(x)x^l(A_2x^2 + A_1x + A_0) \rangle \\ \langle u_0, p_{n-1}(x)x^k(A_2x^2 + A_1x) \rangle & \langle u_0, p_{n-1}(x)x^l(A_2x^2 + A_1x + A_0) \rangle \end{bmatrix} \neq 0.$$

Observe that the case of size $p = 2$ unimodular matrix polynomials is particularly simple, because the degree of the perturbation and its inverse coincide. However, for bigger sizes this is not the case. For a better understanding, let us recall that unimodular matrices always factorize in terms of elementary matrix polynomials and elementary matrices, which are of the following form:

- (i) Elementary matrix polynomials: $e_{i,j}(x) = I_p + E_{i,j}p(x)$ with $i \neq j$ and $E_{i,j}$ the matrix with a 1 at the (i, j) entry and zero elsewhere, and $p(x) \in \mathbb{C}[x]$.
- (ii) Elementary matrices:
 - (a) $I_p + (c - 1)E_{i,i}$ with $c \in \mathbb{C}$.
 - (b) $\eta^{(i,j)} = I_p - E_{i,i} - E_{j,j} + E_{j,i} + E_{i,j}$: the identity matrix with the i th and j th rows interchanged.

The inverses of these matrices are elementary again

$$\begin{aligned} (e_{i,j}(x))^{-1} &= I_p - p(x)E_{i,j}, \\ (I_p + (c - 1)E_{i,i})^{-1} &= I_p + (c^{-1} - 1)E_{i,i}, \\ (\eta^{(i,j)})^{-1} &= \eta^{(i,j)}, \end{aligned}$$

and the inverse of a general unimodular matrix polynomial can be computed immediately once its factorization in terms of elementary matrices is given. However, the degree of the matrix polynomial and its inverse requires a separate analysis.

If our perturbation $W(x) = I_p + p(x)E_{i,j}$ is an elementary matrix polynomial, with $\deg p(x) = N$, then we have that $(W(x))^{-1} = I_p - p(x)E_{i,j}$ and $\deg W(x) = \deg((W(x))^{-1}) = N$. If we assume a departing matrix of generalized kernels $u_{x,y}$, for $n \geq N$, the first family of perturbed polynomials will be

$$\check{P}_n^{[1]}(x) = \Theta_* \begin{bmatrix} \langle P_{n-N}^{[1]}(x), y^{k_1}(I_p + p(y)E_{i,j}) \rangle_u & \dots & & & \\ & \vdots & & & \\ \langle P_n^{[1]}(x), y^{k_1}(I_p + p(y)E_{i,j}) \rangle_u & \dots & & & \\ & & \langle P_{n-N}^{[1]}(x), y^{k_N}(I_p + p(y)E_{i,j}) \rangle_u & P_{n-N}^{[1]}(x) & \\ & & \vdots & \vdots & \\ \langle P_n^{[1]}(x), y^{k_N}(I_p + p(y)E_{i,j}) \rangle_u & & & P_n^{[1]}(x) & \end{bmatrix}.$$

Here, the sequence of different integers $\{k_1, \dots, k_N\} \subset \{1, \dots, n-1\}$ is such that

$$\det \begin{bmatrix} \langle P_{n-N}^{[1]}(x), y^{k_1}(I_p + p(y)E_{i,j}) \rangle_u & \dots & \langle P_{n-N}^{[1]}(x), y^{k_N}(I_p + p(y)E_{i,j}) \rangle_u \\ \vdots & & \vdots \\ \langle P_{n-1}^{[1]}(x), y^{k_1}(I_p + p(y)E_{i,j}) \rangle_u & \dots & \langle P_{n-1}^{[1]}(x), y^{k_N}(I_p + p(y)E_{i,j}) \rangle_u \end{bmatrix} \neq 0.$$

A bit more complex situation appears when we have the product of different elementary matrix polynomials, for example

$$W(x) = (I_p + p_{i_1,j_1}^{(1)}(x)E_{i_1,j_1})(I_p + p_{i_2,j_2}^{(2)}(x)E_{i_2,j_2}),$$

which has two possible forms depending on whether $j_1 \neq i_2$ or $j_1 = i_2$

$$W(x) = \begin{cases} I_p + p_{i_1,j_1}^{(1)}(x)E_{i_1,j_1} + p_{i_2,j_2}^{(2)}(x)E_{i_2,j_2}, & j_1 \neq i_2, \\ I_p + p_{i_1,j_1}^{(1)}(x)E_{i_1,j_1} + p_{j_2,j_2}^{(2)}(x)E_{j_2,j_2} + p_{i_1,j_1}^{(1)}(x)p_{j_1,j_2}^{(2)}(x)E_{i_1,j_2}, & j_1 = i_2, \end{cases}$$

so that

$$\deg(W(x)) = \begin{cases} (1 - \delta_{i_1,i_2} \delta_{j_1,j_2}) \max(\deg(p_{i_1,j_1}^{(1)}(x)), \deg(p_{i_2,j_2}^{(2)}(x))) \\ \quad + \delta_{i_1,i_2} \delta_{j_1,j_2} \deg(p_{i_1,j_1}^{(1)}(x) + p_{i_2,j_2}^{(2)}(x)), & j_1 \neq i_2, \\ \deg(p_{i_1,j_1}^{(1)}(x) + p_{j_1,j_2}^{(2)}(x)), & j_1 = i_2. \end{cases}$$

For the inverse, we find

$$(W(x))^{-1} = \begin{cases} I_p - p_{i_1,j_1}^{(1)}(x)E_{i_1,j_1} - p_{i_2,j_2}^{(2)}(x)E_{i_2,j_2}, & j_2 \neq i_1, \\ I_p - p_{i_1,j_1}^{(1)}(x)E_{i_1,j_1} - p_{i_2,i_1}^{(2)}(x)E_{i_2,i_1} \\ \quad + p_{i_1,j_1}^{(1)}(x)p_{i_2,i_1}^{(2)}(x)E_{i_2,j_1}, & j_2 = i_1, \end{cases}$$

and

$$\deg((W(x))^{-1}) = \begin{cases} (1 - \delta_{i_1, i_2} \delta_{j_1, j_2}) \max(\deg(p_{i_1, j_1}^{(1)}(x)), \deg(p_{i_2, j_2}^{(2)}(x))) \\ \quad + \delta_{i_1, i_2} \delta_{j_1, j_2} \deg(p_{i_1, j_1}^{(1)}(x) + p_{i_2, j_2}^{(2)}), & j_2 \neq i_1, \\ \deg(p_{i_1, j_1}^{(1)}(x)) + \deg(p_{i_2, i_1}^{(2)}(x)), & j_2 = i_1. \end{cases}$$

Thus, if either $j_1 \neq i_2$ and $j_2 \neq i_1$, or when $j_1 = i_2$ and $j_2 = i_1$, the degrees $W(x)$ and $(W(x))^{-1}$ coincide, for $j_1 = i_2$ and $j_2 \neq i_1$ we find $\deg W(x) > \deg((W(x))^{-1})$ and when $j_1 \neq i_2$ and $j_2 = i_1$ we have $\deg W(x) < \deg((W(x))^{-1})$. Consequently, the degrees of unimodular matrix polynomials can be bigger than, equal to or smaller than the degrees of its inverses.

We will be interested in unimodular perturbations $W(x)$ that factorize in terms of K elementary polynomial factors $\{e_{i_m, j_m}(x)\}_{m=1}^K$ and L exchange factors $\{\eta^{(l_n, q_n)}\}_{n=1}^L$. We will use the following notation for elementary polynomials and elementary matrices:

$$(i, j)_{p_{i,j}(x)} := E_{i,j} p_{i,j}(x) \quad [l, q] := \eta_{l,q},$$

suited to take products among them, according to the product table

$$\begin{aligned} (i_1, j_1)_{p_{i_1, j_1}} (i_2, j_2)_{p_{i_2, j_2}} &= \delta_{j_1, i_2} (i_1, j_2)_{p_{i_1, j_1} p_{i_2, j_2}}, \\ [l, q](i, j)_{p_{i,j}} &= (1 - \delta_{l,i})(1 - \delta_{q,i})(i, j)_{p_{i,j}} + \delta_{l,i}(q, j)_{p_{i,j}} + \delta_{q,i}(l, j)_{p_{i,j}}, \\ (i, j)_{p_{i,j}} [l, q] &= (1 - \delta_{l,j})(1 - \delta_{q,j})(i, j)_{p_{i,j}} + \delta_{l,j} j(i,)_{p_{i,j}} + \delta_{q,j}(i, l)_{p_{i,j}}. \end{aligned}$$

Bearing this in mind, we denote all the possible permutations of a vector with K entries, having i out of these equal to 1 and the rest equal to zero, by $\sigma_i^K = \{\sigma_{i,j}^K\}_{j=1}^{|\sigma_i^K|}$ with $\sigma_{i,j}^K = ((\sigma_{i,j}^K)_1, \dots, (\sigma_{i,j}^K)_K) \in (\mathbb{Z}_2)^K$ where $(\sigma_{i,j}^K)_r \in \mathbb{Z}_2 := \{1, 0\}$ and $|\sigma_i^K| = \binom{K}{i}$, we can rewrite a given unimodular perturbation as a sum. Actually, any unimodular polynomial that factorizes in terms of K elementary polynomials $e_{i,j}(x)$ and L elementary matrices $\eta^{(l,q)}$, in a given order, can be expanded into a sum of 2^K terms

$$\begin{aligned} W(x) &= e_{i_1, j_1}(x) \cdots e_{i_r, j_r}(x) \eta^{(l_1, q_1)} \cdots \eta^{(l_t, q_t)} e_{i_{r+1}, j_{r+1}}(x) \cdots \eta^{(l_L, q_L)} \cdots e_{i_K, j_K}(x) \\ &= \sum_{i=0}^K \sum_{j=1}^{|\sigma_i^K|} (i_1, j_1)_{p_{i_1, j_1}}^{(\sigma_{i,j}^K)_1} \cdots (i_r, j_r)_{p_{i_r, j_r}}^{(\sigma_{i,j}^K)_r} [l_1, q_1] \\ &\quad \cdots [l_t, q_t] (i_{r+1}, j_{r+1})_{p_{i_{r+1}, j_{r+1}}}^{(\sigma_{i,j}^K)_{r+1}} \cdots [l_L, q_L] \cdots (i_K, j_K)_{p_{i_K, j_K}}^{(\sigma_{i,j}^K)_K}, \end{aligned}$$

where $(i, j)_{p_{i,j}}^0 = \mathbb{I}_p$. Notice that although in the factorization of W we have assumed that it starts and ends with elementary polynomials, the result would still be valid if it started and/or ended with an interchange elementary matrix η . We notationally simplify these types of expressions by considering the sequences of couples of natural

numbers $\{i_1, j_1\}, \{i_2, j_2\}, \dots, \{i_k, j_k\}$, where $\{n, m\}$ stands either for $(n, m)_{p_m, n}$ or $[m, n]$, and identifying paths. We say that two couples of naturals $\{k, l\}$ and $\{n, m\}$ are linked if $l = n$. When we deal with a couple $[n, m]$ the order is not of the natural numbers which is not relevant, for example (k, l) and $[l, m]$ are linked as well as (k, l) and $[m, l]$ are linked. A path of length l is a subset of I of the form

$$\{\{a_1, a_2\}, \{a_2, a_3\}, \{a_3, a_4\}, \dots, \{a_{l-1}, a_l\}, \{a_l, a_{l+1}\}\}_l.$$

The order of the sequence is respected for the construction of each path. Thus, the element (a_i, a_{i+1}) , as an element of the sequence I , is previous to the element (a_{i+1}, a_{i+2}) in the sequence. A path is proper if it does not belong to a longer path. Out of the 2^K terms that appear only paths remain. In order to know the degree of the unimodular polynomial one must check the factors of the proper paths, and look for the maximum degree involved in those factors. For a better understanding let us work out a couple of significant examples. These examples deal with non-symmetric matrices and, therefore, we have complete Christoffel-type expressions for $\check{P}_n^{[1]}(x)$ and \check{H}_n , but also the mentioned penalty for $P_n^{[2]}(x)$. First, let us consider a polynomial with $K = 5, L = 0$ and $p = 6$,

$$W(x) = e_{1,2}(x)e_{2,3}(x)e_{3,6}(x)e_{4,3}(x)e_{3,5}(x)$$

in terms of sequences of couples the paths for this unimodular polynomial has the following structure:

$$\begin{aligned} & \{\emptyset\}_{i=5}, \\ & \{\emptyset\}_{i=4}, \\ & \underline{\{(1, 2), (2, 3), (3, 6)\}_{i=3}}, \underline{\{(1, 2), (2, 3), (3, 5)\}_{i=3}}, \\ & \underline{\{(4, 3), (3, 5)\}_{i=2}}, \{(2, 3), (3, 5)\}_{i=2}, \{(2, 3), (3, 6)\}_{i=2}, \{(1, 2)(2, 3)\}_{i=2}, \\ & \{(1, 2)\}_{i=1}, \{(2, 3)\}_{i=1}, \{(3, 6)\}_{i=1}, \{(4, 3)\}_{i=1}, \{(3, 5)\}_{i=1}, \\ & \{I_6\}_{i=0}, \end{aligned}$$

where $\{I_6\}_{i=0}$ indicates that the product not involving couples produces the identity matrix (in general will be a product of interchanging matrices) and we have underlined the proper paths. Thus

$$\begin{aligned} W(x) &= e_{1,2}(x)e_{2,3}(x)e_{3,6}(x)e_{4,3}(x)e_{3,5}(x) \\ &= (1, 6)_{p_{1,2}p_{2,3}p_{3,6}} + (1, 5)_{p_{1,2}p_{2,3}p_{3,5}} + (4, 5)_{p_{4,3}p_{3,5}} + (2, 5)_{p_{2,3}p_{3,5}} \\ &\quad + (2, 6)_{p_{2,3}p_{3,6}} + (1, 3)_{p_{1,2}p_{2,3}} \\ &\quad + (1, 2)_{p_{1,2}} + (2, 3)_{p_{2,3}} + (3, 6)_{p_{3,6}} + (4, 3)_{p_{4,3}} + (3, 5)_{p_{3,5}} + I_5 \end{aligned}$$

$$= \begin{bmatrix} 1 & p_{1,2}(x) & p_{1,2}(x)p_{2,3}(x) & 0 & p_{1,2}(x)p_{2,3}(x)p_{3,5}(x) & p_{1,2}(x)p_{2,3}(x)p_{3,6}(x) \\ 0 & 1 & p_{2,3}(x) & 0 & p_{2,3}(x)p_{3,5}(x) & p_{2,3}(x)p_{3,6}(x) \\ 0 & 0 & 1 & 0 & p_{3,5}(x) & p_{3,6}(x) \\ 0 & 0 & p_{4,3}(x) & 1 & p_{4,3}(x)p_{3,5}(x) & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Its inverse is

$$(W(x))^{-1} = (e_{3,5}(x))^{-1}(e_{4,3}(x))^{-1}(e_{3,6}(x))^{-1}(x)(e_{2,3}(x))^{-1}(e_{1,2}(x))^{-1},$$

and the paths are

$$\begin{aligned} & \{\emptyset\}_{i=5}, \\ & \{\emptyset\}_{i=4}, \\ & \{\emptyset\}_{i=3}, \\ & \underline{\{(4, 3), (3, 6)\}_{i=2}}, \\ & \underline{\{(3, 5)\}_{i=1}}, \{(4, 3)\}_{i=1}, \{(3, 6)\}_{i=1}, \underline{\{(2, 3)\}_{i=1}}, \underline{\{(1, 2)\}_{i=1}}, \\ & \{I_6\}_{i=0}. \end{aligned}$$

Thus,

$$\begin{aligned} (W(x))^{-1} &= (4, 6)_{p_{4,3}p_{3,6}} + (3, 5)_{-p_{3,5}} + (4, 3)_{-p_{4,3}} \\ &+ (3, 6)_{-p_{3,6}} + (2, 3)_{-p_{2,3}} + (1, 2)_{-p_{1,2}} + I_6 \\ &= \begin{bmatrix} 1 & -p_{1,2}(x) & 0 & 0 & 0 & 0 \\ 0 & 1 & -p_{2,3}(x) & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & -p_{3,5}(x) & -p_{3,6}(x) \\ 0 & 0 & -p_{4,3}(x) & 1 & 0 & p_{4,3}(x)p_{3,6}(x) \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}. \end{aligned}$$

Then, looking at the proper paths, we find

$$\begin{aligned} \deg W(x) &= \max(\deg p_{1,2}(x) + \deg p_{2,3}(x) + \deg p_{3,6}(x), \deg p_{1,2}(x) \\ &+ \deg p_{2,3}(x) + \deg p_{3,5}(x), \deg p_{4,3}(x) + \deg p_{3,5}(x)), \\ \deg((W(x))^{-1}) &= \max(\deg p_{1,2}(x), \deg p_{2,3}(x), \deg p_{3,6}(x) \\ &+ \deg p_{4,3}(x), \deg p_{3,5}(x)). \end{aligned}$$

For example, if we assume that

$$\begin{aligned} \deg p_{1,2}(x) &= 2, & \deg p_{2,3}(x) &= 1, & \deg p_{3,6}(x) &= 2, \\ \deg p_{4,3}(x) &= 1, & \deg p_{3,5}(x) &= 3, \end{aligned}$$

we get for the corresponding unimodular matrix polynomial and its inverse

$$\deg(W(x)) = 6, \quad \deg((W(x))^{-1}) = 3,$$

so that, for example, the first family of perturbed biorthogonal polynomials, for $n \geq 3$, is

$$\check{P}_n^{[1]}(x) = \Theta_* \begin{bmatrix} \langle P_{n-3}^{[1]}(x), y^{k_1} W(y) \rangle_u & \langle P_{n-3}^{[1]}(x), x^{k_2} W(y) \rangle_u & \langle P_{n-3}^{[1]}(x), y^{k_3} W(y) \rangle_u & P_{n-3}^{[1]}(x) \\ \langle P_{n-2}^{[1]}(x), y^{k_1} W(y) \rangle_u & \langle P_{n-2}^{[1]}(x), y^{k_2} W(y) \rangle_u & \langle P_{n-2}^{[1]}(x), y^{k_3} W(y) \rangle_u & P_{n-2}^{[1]}(x) \\ \langle P_{n-1}^{[1]}(x), y^{k_1} W(y) \rangle_u & \langle P_{n-1}^{[1]}(x), y^{k_2} W(y) \rangle_u & \langle P_{n-1}^{[1]}(x), y^{k_3} W(y) \rangle_u & P_{n-1}^{[1]}(x) \\ \langle P_n^{[1]}(x), y^{k_1} W(y) \rangle_u & \langle P_n^{[1]}(x), y^{k_2} W(y) \rangle_u & \langle P_n^{[1]}(x), y^{k_3} W(y) \rangle_u & P_n^{[1]}(x) \end{bmatrix}. \tag{50}$$

Here, the sequence of different integers $\{k_1, k_2, k_3\} \subset \{1, \dots, n-1\}$ is such that

$$\det \begin{bmatrix} \langle P_{n-3}^{[1]}(x), y^{k_1} W(y) \rangle_u & \langle P_{n-3}^{[1]}(x), y^{k_2} W(y) \rangle_u & \langle P_{n-3}^{[1]}(x), y^{k_3} W(y) \rangle_u \\ \langle P_{n-2}^{[1]}(x), y^{k_1} W(y) \rangle_u & \langle P_{n-2}^{[1]}(x), y^{k_2} W(y) \rangle_u & \langle P_{n-2}^{[1]}(x), y^{k_3} W(y) \rangle_u \\ \langle P_{n-1}^{[1]}(x), y^{k_1} W(y) \rangle_u & \langle P_{n-1}^{[1]}(x), y^{k_2} W(y) \rangle_u & \langle P_{n-1}^{[1]}(x), y^{k_3} W(y) \rangle_u \end{bmatrix} \neq 0.$$

Let us now work out a polynomial with $K = L = 4$ and $p = 5$. The unimodular matrix polynomial we consider is

$$W(x) = e_{2,1}(x)\eta^{(1,4)}\eta^{(5,4)}e_{5,1}(x)\eta^{(3,2)}e_{2,3}(x)\eta^{(3,1)}e_{1,5}(x).$$

The paths are

$$\{\emptyset\}_{i=4},$$

$$\{\emptyset\}_{i=3},$$

$$\underline{\{(2, 1), [1, 4], [5, 4], (5, 1), [3, 2], [3, 1]\}_{i=2}, \{[1, 4], [5, 4], [3, 2], (2, 3), [3, 1], (1, 5)\}_{i=2}},$$

$$\{(2, 1), [1, 4], [5, 4], [3, 2], [3, 1]\}_{i=1}, \{[1, 4], [5, 4], (5, 1), [3, 2], [3, 1]\}_{i=1},$$

$$\{[1, 4], [5, 4], [3, 2], (2, 3), [3, 1]\}_{i=1}, \{[1, 4], [5, 4], [3, 2], [3, 1], (1, 5)\}_{i=1},$$

$$\{[1, 4], [5, 4], [3, 2], [3, 1]\}_{i=0},$$

so that

$$W(x) = (2, 3)_{p_{2,1}p_{5,1}} + (3, 5)_{p_{2,3}p_{1,5}} + (2, 5)_{p_{2,1}} + (1, 3)_{p_{5,1}} + (3, 1)_{p_{2,3}} + (2, 5)_{p_{1,5}} + [1, 4][5, 4][3, 2][3, 1] \quad (51)$$

$$= \begin{bmatrix} 0 & 0 & p_{5,1}(x) & 0 & 1 \\ 1 & 0 & p_{2,1}(x)p_{5,1}(x) & 0 & p_{2,1}(x) + p_{1,5}(x) \\ p_{2,3}(x) & 1 & 0 & 0 & p_{2,3}(x)p_{1,5}(x) \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}. \quad (52)$$

The inverse matrix is

$$(W(x))^{-1} = (e_{1,5}(x))^{-1}\eta^{(3,1)}(e_{2,3}(x))^{-1}\eta^{(3,2)}(e_{5,1}(x))^{-1}\eta^{(5,4)}\eta^{(1,4)}(e_{2,1}(x))^{-1},$$

with paths given by

$$\{\emptyset\}_{i=4},$$

$$\{\emptyset\}_{i=3},$$

$$\underline{\{(1, 5), [3, 1], [3, 2], (5, 1), [5, 4], [1, 4]\}_{i=2}}, \underline{\{[3, 1], (2, 3), [3, 2], [5, 4], [1, 4], (2, 1)\}_{i=2}},$$

$$\{[3, 1], [3, 2], [5, 4], [1, 4], (2, 1)\}_{i=1}, \{[3, 1], [3, 2], (5, 1), [5, 4], [1, 4]\}_{i=1},$$

$$\{[3, 1], (2, 3), [3, 2], [5, 4], [1, 4]\}_{i=1}, \{(1, 5), [3, 1], [3, 2], [5, 4], [1, 4]\}_{i=1},$$

$$\{[3, 1], [3, 2], [5, 4], [1, 4]\}_{i=0},$$

and consequently,

$$(W(x))^{-1} = (1, 4)_{p_{1,5}p_{5,1}} + (2, 1)_{p_{2,3}p_{2,1}} + (1, 1)_{-p_{2,1}} + (5, 4)_{-p_{5,1}} + (2, 2)_{-p_{2,3}} + (1, 1)_{-p_{1,5}} + [3, 1][3, 2][5, 4][1, 4] \\ = \begin{bmatrix} -p_{2,1}(x) - p_{1,5}(x) & 1 & 0 & p_{1,5}(x)p_{5,1}(x) & 0 \\ p_{2,3}(x)p_{2,1}(x) & -p_{2,3}(x) & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & -p_{5,1}(x) & 0 \end{bmatrix}.$$

Proper paths, which we have underlined, give the degrees of the polynomials

$$\deg W(x) = \max(\deg p_{2,1}(x) + \deg p_{5,1}(x), \deg p_{1,5}(x) + \deg p_{2,3}(x)),$$

$$\deg((W(x))^{-1}) = \max(\deg p_{1,5}(x) + \deg p_{5,1}(x), \deg p_{2,3}(x) + \deg p_{2,1}(x)).$$

For example, if we assume that

$$\deg p_{2,1}(x) = 2, \quad \deg p_{5,1}(x) = 1, \quad \deg p_{1,5}(x) = 2, \quad \deg p_{2,3}(x) = 1,$$

we find $\deg W(x) = \deg((W(x))^{-1}) = 3$ and formula (50) is applicable for $W(x)$ as given in (51).

If we seek for symmetric unimodular polynomials of the form

$$W(x) = V(x)(V(x))^T,$$

where $V(x)$ is a unimodular matrix polynomial. For example, we put $p = 4$, and consider

$$V(x) = \begin{bmatrix} 1 & p_{1,2}(x)p_{3,2}(x) & p_{1,2}(x) & 0 \\ 0 & p_{3,2}(x) & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & p_{4,3}(x) & 1 \end{bmatrix},$$

in such a way the perturbing symmetric unimodular matrix polynomial is

$$W(x) = \begin{bmatrix} 1 + (p_{1,2}(x))^2(p_{3,2}(x))^2 & p_{1,2}(x)(p_{3,2}(x))^2 + p_{1,2}(x) \\ p_{1,2}(x)(p_{3,2}(x))^2 + p_{1,2}(x) & 1 + (p_{3,2}(x))^2 \\ p_{1,2}(x)p_{3,2}(x) & p_{3,2}(x) \\ p_{1,2}(x)p_{4,3}(x) & p_{4,3}(x) \\ p_{1,2}(x)p_{3,2}(x) & p_{1,2}(x)p_{4,3}(x) \\ p_{3,2}(x) & p_{4,3}(x) \\ 1 & 0 \\ 0 & 1 + (p_{4,3}(x))^2 \end{bmatrix}.$$

Let us assume that

$$\deg p_{1,2}(x) = 3, \quad \deg p_{3,2}(x) = 1, \quad \deg p_{4,3}(x) = 1,$$

then

$$\deg W(x) = 8, \quad \deg((W(x))^{-1}) = 4.$$

Now, we take a scalar matrix of linear functionals $u = u_0 I_p$, with $u_0 \in (\mathbb{R}[x])'$ positive definite, and assume that the polynomials $p_{1,2}(x), p_{2,3}(x), p_{3,4}(x) \in \mathbb{R}[x]$. Then, we obtain matrix orthogonal polynomials $\{P_n(x)\}_{n=0}^\infty$ for the matrix of linear functionals $W(x)u_0$, which in terms of the sequence of scalar orthogonal polynomials

$\{p_n(x)\}_{n=0}^\infty$ of the linear functional u_0 are, for $n \geq 4$,

$$P_n(x) = \Theta_* \begin{bmatrix} \langle u_0, p_{n-4}(x)x^{k_1}W(x) \rangle & \langle u_0, p_{n-4}(x)x^{k_2}W(x) \rangle & \langle u_0, p_{n-4}(x)x^{k_3}W(x) \rangle \\ \langle u_0, p_{n-3}(x)x^{k_1}W(x) \rangle & \langle u_0, p_{n-3}(x)x^{k_2}W(x) \rangle & \langle u_0, p_{n-3}(x)x^{k_3}W(x) \rangle \\ \langle u_0, p_{n-2}(x)x^{k_1}W(x) \rangle & \langle u_0, p_{n-2}(x)x^{k_2}W(x) \rangle & \langle u_0, p_{n-2}(x)x^{k_3}W(x) \rangle \\ \langle u_0, p_{n-1}(x)x^{k_1}W(x) \rangle & \langle u_0, p_{n-1}(x)x^{k_2}W(x) \rangle & \langle u_0, p_{n-1}(x)x^{k_3}W(x) \rangle \\ \langle u_0, p_n(x)x^{k_1}W(x) \rangle & \langle u_0, p_n(x)x^{k_2}W(x) \rangle & \langle u_0, p_n(x)x^{k_3}W(x) \rangle \end{bmatrix} \cdot \begin{bmatrix} \langle u_0, p_{n-4}(x)x^{k_4}W(x) \rangle & p_{n-4}(x)I_p \\ \langle u_0, p_{n-3}(x)x^{k_4}W(x) \rangle & p_{n-3}(x)I_p \\ \langle u_0, p_{n-2}(x)x^{k_4}W(x) \rangle & p_{n-2}(x)I_p \\ \langle u_0, p_{n-1}(x)x^{k_4}W(x) \rangle & p_{n-1}(x)I_p \\ \langle u_0, p_n(x)x^{k_4}W(x) \rangle & p_n(x)I_p \end{bmatrix}.$$

The set $\{k_1, k_2, k_3, k_4\} \subset \{1, \dots, n-1\}$ is such that

$$\det \begin{bmatrix} \langle u_0, p_{n-4}(x)x^{k_1}W(x) \rangle & \langle u_0, p_{n-4}(x)x^{k_2}W(x) \rangle \\ \langle u_0, p_{n-3}(x)x^{k_1}W(x) \rangle & \langle u_0, p_{n-3}(x)x^{k_2}W(x) \rangle \\ \langle u_0, p_{n-2}(x)x^{k_1}W(x) \rangle & \langle u_0, p_{n-2}(x)x^{k_2}W(x) \rangle \\ \langle u_0, p_{n-1}(x)x^{k_1}W(x) \rangle & \langle u_0, p_{n-1}(x)x^{k_2}W(x) \rangle \\ \langle u_0, p_{n-4}(x)x^{k_3}W(x) \rangle & \langle u_0, p_{n-4}(x)x^{k_4}W(x) \rangle \\ \langle u_0, p_{n-3}(x)x^{k_3}W(x) \rangle & \langle u_0, p_{n-3}(x)x^{k_4}W(x) \rangle \\ \langle u_0, p_{n-2}(x)x^{k_3}W(x) \rangle & \langle u_0, p_{n-2}(x)x^{k_4}W(x) \rangle \\ \langle u_0, p_{n-1}(x)x^{k_3}W(x) \rangle & \langle u_0, p_{n-1}(x)x^{k_4}W(x) \rangle \end{bmatrix} \neq 0.$$

2.7.2. Degree one matrix Geronimus transformations

We consider a degree one perturbing polynomial of the form

$$W(x) = xI_p - A,$$

and assume, for the sake of simplicity, that all ξ are taken zero, i.e. there are no masses. Observe that in this case a Jordan pair (X, J) is such that $A = XJX^{-1}$, and Lemma 1 implies that the root spectral jet of a polynomial $P(x) = \sum_k P_k x^k \in \mathbb{C}^{p \times p}[x]$ is $\mathcal{J}_P = P(A)X$, where we understand a right evaluation, i.e. $P(A) := \sum_k P_k A^k$. An similar argument, for $\sigma(A) \cap \text{supp}_y(u) = \emptyset$, yields

$$\mathcal{J}_{C_n^{[1]}} = \langle P^{[1]}(x), (A - I_p y)^{-1} X \rangle_u,$$

expressed in terms of the resolvent $(A - I_p y)^{-1}$ of A . Formally, it can be written as

$$\mathcal{J}_{C_n^{[1]}} = C_n^{[1]}(A)X,$$

where we again understand a right evaluation in the Taylor series of the Cauchy transform. Moreover, we also need the root spectral jet of the mixed Christoffel–Darboux kernel

$$\mathcal{J}_{K_{n-1}^{(pc)}}(y) = \sum_{k=0}^{n-1} (P_k^{[2]}(y))^\top (H_k)^{-1} C_k^{[1]}(A) X =: K_{n-1}^{(pc)}(A, y) X,$$

that for a Hankel generalized kernel $u_{x,y}$, using the Christoffel–Darboux formula for mixed kernels, reads

$$\begin{aligned} \mathcal{J}_{K_{n-1}^{(pc)}}(y) &= ((P_{n-1}^{[2]}(y))^\top (H_{n-1})^{-1} C_n^{[1]}(A) \\ &\quad - (P_n^{[2]}(y))^\top (H_{n-1})^{-1} C_{n-1}^{[1]}(A) + I_p)(A - I_p y)^{-1} X. \end{aligned}$$

We also have $\mathcal{V}(x, y) = I_p$ so that $\mathcal{J}_{\mathcal{V}} = X$.

Thus, for $n \geq 1$ we have

$$\begin{aligned} \check{P}_n^{[1]}(x) &= \Theta_* \begin{bmatrix} C_{n-1}^{[1]}(A) X & P_{n-1}^{[1]}(x) \\ C_n^{[1]}(A) X & P_n^{[1]}(x) \end{bmatrix} \\ &= P_n^{[1]}(x) - C_n^{[1]}(A) (C_{n-1}^{[1]}(A))^{-1} P_{n-1}^{[1]}(x), \\ \check{H}_n &= \Theta_* \begin{bmatrix} C_{n-1}^{[1]}(A) X & H_{n-1} \\ C_n^{[1]}(A) X & 0_p \end{bmatrix} \\ &= -C_n^{[1]}(A) (C_{n-1}^{[1]}(A))^{-1} H_{n-1}, \\ (\check{P}_n^{[2]}(y))^\top &= \Theta_* \begin{bmatrix} C_{n-1}^{[1]}(A) X & H_{n-1} \\ (I_p y - A) (K_{n-1}^{(pc)}(A, y) + I_p) X & 0_p \end{bmatrix} \\ &= ((I_p y - A) K_{n-1}^{(pc)}(A, y) + I_p) (C_{n-1}^{[1]}(A))^{-1} H_{n-1}. \end{aligned}$$

For a Hankel matrix of bivariate generalized functionals, i.e. with a Hankel Gram matrix so that the Christoffel–Darboux formula holds, we have

$$\begin{aligned} (\check{P}_n^{[2]}(y))^\top &= -(I_p y - A) ((P_{n-1}^{[2]}(y))^\top (H_{n-1})^{-1} C_n^{[1]}(A) \\ &\quad - (P_n^{[2]}(y))^\top (H_{n-1})^{-1} C_{n-1}^{[1]}(A)) (I_p y - A)^{-1} H_{n-1}. \end{aligned}$$

Appendix A. Schur Complements and Quasideterminants

We first notice that the Schur complement was not introduced by Schur but by Haynsworth in 1968 in [44, 45]. Haynsworth coined that named because the Schur determinant formula given in what today is known as Schur lemma in [65]. For an ample overview on Schur complement and many of its applications see [72]. The most easy examples of quasideterminants are Schur complements. Gel’fand and collaborators have made many essential contributions to the subject, see [27] for an excellent survey on the subject. Olver’s on a paper on multivariate interpola-

tion, see [58], discusses an alternative interesting approach to the subject. In the late 1920 Richardson [61, 62], and Heyting [46] studied possible extensions of the determinant notion to division rings. Heyting defined the *designant* of a matrix with noncommutative entries, which for 2×2 matrices was the Schur complement, and generalized to larger dimensions by induction. Let us stress that both Richardson's and Heyting's *quasideterminants* were generically rational functions of the matrix coefficients. In 1931, Ore [59] gave a polynomial proposal, the Ore's determinant. A definitive impulse to the modern theory was given by the Gel'fand's school [22, 23, 28–31]. Quasideterminants defined over free division rings were early noticed that are not an analog of the commutative determinant but rather of ratio determinants. An essential aspect for quasideterminants is the *heredity principle*, quasideterminants of quasideterminants are quasideterminants; there is no analog of such a principle for determinants. Many of the properties of determinants extend to this case, see the cited papers and also [49] for quasi-minors expansions. Already in the early 1990 the Gelf'and school [29] noticed the role quasideterminants for some integrable systems, see also [60] for some recent work in this direction regarding non-Abelian Toda and Painlevé II equations. Nimmo and his collaborators, the Glasgow school, have studied the relation of quasideterminants and integrable systems, in particular we can mention the papers [36–38, 50]. All this paved the route, using the connection with orthogonal polynomials *à la Cholesky*, to the appearance of quasideterminants in the multivariate orthogonality context. Later, in 2006 Olver applied quasideterminants to multivariate interpolation [58].

A.1. Schur complements

Given $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ in block form the Schur complement with respect to A (if $\det A \neq 0$) is

$$\text{SC} \left(\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right) \equiv M/A := D - CA^{-1}B.$$

The Schur complement with respect to D (if $\det D \neq 0$) is

$$\text{SC}_D \left(\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right) \equiv M/D := A - BD^{-1}C.$$

Observe that we have the block Gauss factorization

$$\begin{aligned} \begin{pmatrix} A & B \\ C & D \end{pmatrix} &= \begin{pmatrix} \mathbb{I} & 0 \\ CA^{-1} & \mathbb{I} \end{pmatrix} \begin{pmatrix} A & 0 \\ 0 & M/A \end{pmatrix} \begin{pmatrix} \mathbb{I} & A^{-1}B \\ 0 & \mathbb{I} \end{pmatrix} \\ &= \begin{pmatrix} \mathbb{I} & BD^{-1} \\ 0 & \mathbb{I} \end{pmatrix} \begin{pmatrix} M/D & 0 \\ 0 & D \end{pmatrix} \begin{pmatrix} \mathbb{I} & 0 \\ D^{-1}C & \mathbb{I} \end{pmatrix} \end{aligned}$$

implies the Schur determinant formula $\det M = \det(A) \det(M/A)$. This is in fact the Schur lemma in a disguise form, in fact Schur lemma in [65] assumes that

$[A, C] = 0$ so that $\det M = \det(AD - BC)$. In terms of the Schur complements we have the following well-known expressions for the inverse matrices

$$\begin{aligned}
 M^{-1} &= \begin{pmatrix} \mathbb{I} & -A^{-1}B \\ 0 & \mathbb{I} \end{pmatrix} \begin{pmatrix} A^{-1} & 0 \\ 0 & (M/A)^{-1} \end{pmatrix} \begin{pmatrix} \mathbb{I} & 0 \\ -CA^{-1} & \mathbb{I} \end{pmatrix} \\
 &= \begin{pmatrix} A^{-1} + A^{-1}B(M/A)^{-1}CA^{-1} & -A^{-1}B(M/A)^{-1} \\ -(M/A)^{-1}CA^{-1} & (M/A)^{-1} \end{pmatrix} \\
 &= \begin{pmatrix} \mathbb{I} & 0 \\ -D^{-1}C & \mathbb{I} \end{pmatrix} \begin{pmatrix} (M/D)^{-1} & 0 \\ 0 & D^{-1} \end{pmatrix} \begin{pmatrix} \mathbb{I} & -BD^{-1} \\ 0 & \mathbb{I} \end{pmatrix} \\
 &= \begin{pmatrix} (M/D)^{-1} & -(M/D)^{-1}BD^{-1} \\ -D^{-1}B(M/D)^{-1} & D^{-1} + D^{-1}(M/D)^{-1}BD^{-1} \end{pmatrix}. \tag{53}
 \end{aligned}$$

A.2. Quasideterminants and the heredity principle

Given any partitioned matrix where $A_{i,j} \in \mathbb{R}^{m_i \times m_j}$ for $i, j \in \{1, \dots, k-1\}$, and $A_{k,k} \in \mathbb{R}^{\kappa_1 \times \kappa_2}$, $A_{i,k} \in \mathbb{R}^{m_i \times \kappa_2}$ and $A_{k,j} \in \mathbb{R}^{\kappa_1 \times m_j}$, we are going to define its quasideterminant *à la Olver*. We start with $k = 2$, so that $A = \begin{pmatrix} A_{1,1} & A_{1,2} \\ A_{2,1} & A_{2,2} \end{pmatrix}$, in this case the first quasideterminant is different to that of the Gel'fand school where $\Theta_1(A) = |A|_{2,2} = \begin{vmatrix} A_{1,1} & A_{1,2} \\ A_{2,1} & A_{2,2} \end{vmatrix}$. There is another quasideterminant $\Theta_2(A) = A/A_{22} = |A|_{1,1} = \begin{vmatrix} A_{1,1} & A_{1,2} \\ A_{2,1} & A_{2,2} \end{vmatrix}$, the other Schur complement, and we need $A_{2,2}$ to be an invertible square matrix. Other quasideterminants that can be considered for regular square blocks are $\begin{vmatrix} A_{1,1} & A_{1,2} \\ A_{2,1} & A_{2,2} \end{vmatrix}$ and $\begin{vmatrix} A_{1,1} & A_{1,2} \\ A_{2,1} & A_{2,2} \end{vmatrix}$.

Following [58] we remark that quasideterminantal reduction is a commutative operation. This is the heredity principle formulated by Gel'fand and Retakh [27, 31]: quasideterminants of quasideterminants are quasideterminants. Let us illustrate this by reproducing a nice example discussed in [58]. We consider the matrix and take the quasideterminant with respect to the first diagonal block, which we define as the Schur complement indicated by the non-dashed lines, to get a matrix with blocks with subindexes involving 2 and 3 but not 1. Notice also that we are allowed to take blocks of different sizes we have taken the quasideterminant with respect to a bigger block, composed of two rows and columns of basic blocks. This is the Olver's generalization of Gel'fand's *et al.* construction. Now, we take the quasideterminant given by the Schur complement as indicated by the dashed lines, to get

$$\Theta_2(\Theta_1(A)) = \begin{vmatrix} A_{2,2} - A_{2,1}A_{1,1}^{-1}A_{1,2} & A_{2,3} - A_{2,1}A_{1,1}^{-1}A_{1,3} \\ A_{3,2} - A_{3,1}A_{1,1}^{-1}A_{1,2} & \boxed{A_{3,3} - A_{3,1}A_{1,1}^{-1}A_{1,3}} \end{vmatrix} \tag{54}$$

$$\begin{aligned}
 &= A_{3,3} - A_{3,1}A_{1,1}^{-1}A_{1,3} - (A_{3,2} - A_{3,1}A_{1,1}^{-1}A_{1,2})(A_{2,2} \\
 &\quad - A_{2,1}A_{1,1}^{-1}A_{1,2})^{-1}(A_{2,3} - A_{2,1}A_{1,1}^{-1}A_{1,3}). \tag{55}
 \end{aligned}$$

We are ready to compute, for the very same matrix

$$A = \left(\begin{array}{cc|c} A_{1,1} & A_{1,2} & A_{1,3} \\ A_{2,1} & A_{2,2} & A_{2,3} \\ \hline A_{3,1} & A_{3,2} & A_{3,3} \end{array} \right), \tag{56}$$

the quasideterminant associated to the first two diagonal blocks, that we label as $\{1, 2\}$; i.e. the Schur complement indicated by the non-dashed lines in (56), to get

$$\begin{aligned} \Theta_{\{1,2\}}(A) &= \left| \begin{array}{ccc} A_{1,1} & A_{1,2} & A_{1,3} \\ A_{2,1} & A_{2,2} & A_{2,3} \\ A_{1,3} & A_{2,3} & \boxed{A_{3,3}} \end{array} \right| \\ &= A_{3,3} - (A_{3,1} \quad A_{3,2}) \begin{pmatrix} A_{1,1} & A_{1,2} \\ A_{2,1} & A_{2,2} \end{pmatrix}^{-1} \begin{pmatrix} A_{1,3} \\ A_{2,3} \end{pmatrix}. \end{aligned}$$

But recalling (53)

$$\begin{aligned} &\begin{pmatrix} A_{1,1} & A_{1,2} \\ A_{2,1} & A_{2,2} \end{pmatrix}^{-1} \\ &= \begin{pmatrix} A_{1,1}^{-1} + A_{1,1}^{-1}A_{1,2}(A_{2,2} - A_{2,1}A_{1,1}^{-1}A_{1,2})^{-1}A_{2,1}A_{1,1}^{-1} & -A_{1,1}^{-1}A_{1,2}(A_{2,2} - A_{2,1}A_{1,1}^{-1}A_{1,2})^{-1} \\ -(A_{2,2} - A_{2,1}A_{1,1}^{-1}A_{1,2})^{-1}A_{2,1}A_{1,1}^{-1} & (A_{2,2} - A_{2,1}A_{1,1}^{-1}A_{1,2})^{-1} \end{pmatrix} \end{aligned}$$

we get

$$\begin{aligned} \Theta_{\{1,2\}}(A) &= A_{3,3} - A_{3,1}A_{1,1}^{-1}A_{1,3} + A_{3,1}A_{1,1}^{-1}A_{1,2} \\ &\quad \times (A_{2,2} - A_{2,1}A_{1,1}^{-1}A_{1,2})^{-1}A_{2,1}A_{1,1}^{-1}A_{1,3} \\ &\quad - A_{3,2}(A_{2,2} - A_{2,1}A_{1,1}^{-1}A_{1,2})^{-1}A_{2,1}A_{1,1}^{-1}A_{1,3} \\ &\quad - A_{3,1}A_{1,1}^{-1}A_{1,2}(A_{2,2} - A_{2,1}A_{1,1}^{-1}A_{1,2})^{-1}A_{2,3} \\ &\quad + A_{3,2}(A_{2,2} - A_{2,1}A_{1,1}^{-1}A_{1,2})^{-1}A_{2,3} \end{aligned}$$

which is identical to (54), so that

$$\Theta_2(\Theta_1(A)) = \Theta_{\{1,2\}}(A).$$

Given any set $I = \{i_1, \dots, i_m\} \subset \{1, \dots, k\}$ the heredity principle allows us to define the quasideterminant^b

$$\Theta_I(A) = \Theta_{i_1}(\Theta_{i_2}(\dots \Theta_{i_m}(A) \dots))$$

^bIn [58], it is defined as the Schur complement with respect to a big block built up by the blocks determined by the indices I .

and the ℓ th quasideterminant is

$$\Theta^{(\ell)}(A) = \Theta_{\{1, \dots, \ell-1, \ell+1, \dots, k\}}(A) = |A|_{\ell, \ell} = \begin{vmatrix} A_{1,1} & A_{1,2} & \dots & A_{1,\ell} & \dots & A_{1,k} \\ A_{2,1} & A_{2,2} & \dots & A_{2,\ell} & \dots & A_{2,k} \\ \vdots & \vdots & & \vdots & & \vdots \\ A_{\ell,1} & A_{\ell,2} & \dots & \boxed{A_{\ell,\ell}} & \dots & A_{\ell,k} \\ \vdots & \vdots & & \vdots & & \vdots \\ A_{k,1} & A_{k,2} & \dots & A_{k,\ell} & \dots & A_{k,k} \end{vmatrix}.$$

The last quasideterminant is denoted by

$$\Theta_*(A) = \Theta^{(k)}(A) = |A|_{k,k} = \begin{vmatrix} A_{1,1} & A_{1,2} & \dots & A_{1,k} \\ A_{2,1} & A_{2,2} & \dots & A_{2,k} \\ \vdots & \vdots & & \vdots \\ A_{k,1} & A_{k,2} & \dots & \boxed{A_{k,k}} \end{vmatrix}.$$

Acknowledgments

We thank the comments and suggestions by the referees. They have contributed to improve the presentation of the manuscript.

G. Ariznabarreta and M. Mañas thank financial support from the Spanish “Ministerio de Economía y Competitividad” research project [MTM2015-65888-C4-3-P], *Ortogonalidad, teoría de la aproximación y aplicaciones en física matemática*.

G. Ariznabarreta thank financial support from the Universidad Complutense de Madrid Program “Ayudas para Becas y Contratos Complutenses Predoctorales en España 2011”.

J. C. García-Ardila and F. Marcellán thank financial support from the Spanish “Ministerio de Economía y Competitividad” research project [MTM2015-65888-C4-2-P], *Ortogonalidad, teoría de la aproximación y aplicaciones en física matemática*.

References

- [1] M. Adler and P. van Moerbeke, Generalized orthogonal polynomials, discrete KP and Riemann–Hilbert problems, *Comm. Math. Phys.* **207** (1999) 589–620.
- [2] C. Álvarez Fernández, G. Ariznabarreta, J. C. García-Ardila, M. Mañas and F. Marcellán, Christoffel transformations for matrix orthogonal polynomials in the real line and the non-Abelian 2D Toda lattice hierarchy, *Int. Math. Res. Not.* **2017** (2017) 1285–1341.
- [3] C. Álvarez-Fernández, M. Mañas and U. Fidalgo Prieto, The multicomponent 2D Toda hierarchy: generalized matrix orthogonal polynomials, multiple orthogonal polynomials and Riemann–Hilbert problems, *Inverse Problems* **26** (2010) 055009.

- [4] R. Álvarez-Nodarse, A. J. Durán and A. Martínez de los Ríos, *Orthogonal matrix polynomials satisfying second-order difference equation*, *J. Approx. Theory* **169** (2013) 40–55.
- [5] A. I. Aptekarev and E. M. Nikishin, The scattering problem for a discrete Sturm–Liouville operator, *Math. USSR-Sb.* **49** (1984) 325–355.
- [6] G. Ariznabarreta, J. C. García-Ardila, M. Mañas and F. Marcellán, Non-Abelian integrable hierarchies: Matrix biorthogonal polynomials and perturbations, *J. Phys. A: Math. Theor.* **51** (2018) 205204.
- [7] G. Ariznabarreta and M. Mañas, Matrix orthogonal Laurent polynomials on the unit circle and Toda type integrable systems, *Adv. Math.* **264** (2014) 396–463.
- [8] G. Ariznabarreta and M. Mañas, Multivariate orthogonal polynomials and integrable systems, *Adv. Math.* **302** (2016) 628–739.
- [9] M. I. Bueno and F. Marcellán, Darboux transformation and perturbation of linear functionals, *Linear Algebra Appl.* **384** (2004) 215–242.
- [10] M. J. Cantero, F. Marcellán, L. Moral and L. Velázquez, Darboux transformations for CMV matrices, *Adv. Math.* **298** (2016) 122–206.
- [11] M. M. Castro and F. A. Grünbaum, Orthogonal matrix polynomials satisfying first-order differential equations: A collection of instructive examples, *J. Nonlinear Math. Phys.* **4** (2005) 63–76.
- [12] T. S. Chihara, *An Introduction to Orthogonal Polynomials*, in Mathematics and its Applications Series, Vol. 13 (Gordon and Breach Science Publishers, New York, 1978).
- [13] E. B. Christoffel, Über die Gaussische quadratur und eine verallgemeinerung derselben, *J. Reine Angew. Math.* **55** (1858) 61–82 (in German).
- [14] R. Cooke, *Infinite Matrices and Sequences Spaces* (MacMillan, London, 1950). Reprinted in Dover Books on Mathematics (Dover Publications, 2014).
- [15] D. Damanik, A. Pushnitski and B. Simon, The analytic theory of matrix orthogonal polynomials, *Surveys Approx. Theory* **4** (2008) 1–85.
- [16] M. Derevyagin, J. C. García-Ardila and F. Marcellán, Multiple Geronimus transformations, *Linear Algebra Appl.* **454** (2014) 158–183.
- [17] M. Derevyagin and F. Marcellán, A note on the Geronimus transformation and Sobolev orthogonal polynomials, *Numer. Algorithms* **67** (2014) 271–287.
- [18] A. J. Durán, On orthogonal polynomials with respect to a definite positive matrix of measures, *Canad. J. Math.* **47** (1995) 88–112.
- [19] A. J. Durán, Markov’s theorem for orthogonal matrix polynomials, *Canad. J. Math.* **48** (1996) 1180–1195.
- [20] A. J. Durán, Matrix inner product having a matrix symmetric second-order differential operator, *Rocky Mountain J. Math.* **27** (1997) 585–600.
- [21] A. J. Durán and F. A. Grünbaum, A survey on orthogonal matrix polynomials satisfying second-order differential equations, *J. Comput. Appl. Math.* **178** (2005) 169–190.
- [22] P. Etinghof, I. M. Gel’fand, S. Gel’fand and V. S. Retakh, Factorization of differential operators, quasideterminants, and nonabelian Toda field equations, *Math. Res. Lett.* **4** (1997) 413–425.
- [23] P. Etinghof, I. M. Gel’fand, S. Gel’fand and V. S. Retakh, Nonabelian integrable systems, quasideterminants, and Marchenko lemma, *Math. Res. Lett.* **5** (1998) 1–12.
- [24] P. A. Fuhrmann, Orthogonal matrix polynomials and system theory, *Rend. Sem. Mat. Univ. Politec. Torino* (1987), Special Issue, 68–124.
- [25] W. Gautschi, An algorithmic implementation of the generalized Christoffel theorem in *Numerical Integration*, ed. G. Hämmerlin, International Series of Numerical Mathematics, Vol. 57 (Birkhäuser, Basel, 1982), pp. 89–106.

- [26] W. Gautschi, *Orthogonal Polynomials Computation and Approximation*, Numerical Mathematics and Scientific Computation (Oxford University Press, Oxford, 2004).
- [27] I. M. Gel'fand, S. Gel'fand, V. S. Retakh and R. Wilson, Quasideterminants, *Adv. Math.* **193** (2005) 56–141.
- [28] I. M. Gel'fand, D. Krob, A. Lascoux, B. Leclerc, V. S. Retakh and J.-Y. Thibon, Noncommutative symmetric functions, *Adv. Math.* **112** (1995) 218–348.
- [29] I. M. Gel'fand and V. S. Retakh, Determinants of matrices over noncommutative rings, *Funct. Anal. Appl.* **25** (1991) 91–102.
- [30] I. M. Gel'fand and V. S. Retakh, Theory of noncommutative determinants, and characteristic functions of graphs, *Funct. Anal. Appl.* **26** (1992) 231–246.
- [31] I. M. Gel'fand and V. S. Retakh, quasideterminants. I, *Selecta Math. (New Series)* **3** (1997) 517–546.
- [32] I. M. Gel'fand and G. E. Shilov, *Generalized Functions. Vol. I: Properties and Operations* (Academic Press, New York, 1964). Reprinted in the AMS Chelsea Publishing (American Mathematical Society, Providence, RI, 2016).
- [33] I. M. Gel'fand and G. E. Shilov, *Generalized Functions. Vol. II: Spaces of Fundamental Solutions and Generalized Functions* (Academic Press, New York, 1968). Reprinted in the AMS Chelsea Publishing (American Mathematical Society, Providence, RI, 2016).
- [34] J. S. Geronimo, Scattering theory and matrix orthogonal polynomials on the real line, *Circuits Syst. Signal Process* **1** (1982) 471–495.
- [35] J. Geronimus, On polynomials orthogonal with regard to a given sequence of numbers and a theorem by W. Hahn, *Izv. Akad. Nauk SSSR* **4** (1940) 215–228 (in Russian).
- [36] C. R. Gilson and S. R. Macfarlane, Dromion solutions of noncommutative Davey–Stewartson equations, *J. Phys. A: Math. Theor.* **42** (2009) 235202.
- [37] C. R. Gilson, J. C. Nimmo and Y. Ohta, Quasideterminant solutions of a non-Abelian Hirota–Miwa equation, *J. Phys. A: Math. Theor.* **40** (2007) 12607–12617.
- [38] C. R. Gilson, J. C. Nimmo and C. M. Sooman, On a direct approach to quasideterminant solutions of a noncommutative modified KP equation, *J. Phys. A: Math. Theor.* **41** (2008) 085202.
- [39] I. Gohberg, P. Lancaster and L. Rodman, *Matrix Polynomials*, Computer Science and Applied Mathematics (Academic Press, [Harcourt Brace Jovanovich Publisher], New York, 1982).
- [40] L. Golinskii, On the scientific legacy of Ya.L. Geronimus (to the hundredth anniversary), in *Self-Similar Systems*, eds. V. B. Priezhev and V. P. Spiridonov (Publishing Department, Joint Institute for Nuclear Research, Moscow Region, Dubna, 1999), pp. 273–281.
- [41] F. A. Grünbaum, I. Pacharoni and J. Tirao, Matrix valued orthogonal polynomials of the Jacobi type, *Indag. Math.* **14** (2003) 353–366.
- [42] F. A. Grünbaum, I. Pacharoni and J. Tirao, Matrix valued orthogonal polynomials of the Jacobi type: The role of group representation theory, *Ann. Inst. Fourier* **55** (2005) 2051–2068.
- [43] W. Hahn, Über die Jacobischen polynome und zwei verwandte polynomklassen, *Math. Z.* **39** (1935) 634–638 (in German).
- [44] E. V. Haynsworth, On the Schur complement, *Basel Mathematical Notes*, University of Basel (1968), Vol. 20, 17 pp.
- [45] E. V. Haynsworth, Reduction of a matrix using properties of the Schur complement, *Linear Algebra Appl.* **3** (1970) 23–29.

- [46] A. Heyting, Die theorie der linearen Gleichungen in einer zahlenspezies mit nichtkommutativer multiplikation, *Math. Ann.* **98** (1928) 465–490.
- [47] L. Hörmander, *The Analysis of Partial Differential Operators I, Distribution Theory and Fourier Analysis*, 2nd edn. (Springer, New York, 1990).
- [48] M. G. Krein, The fundamental propositions of the theory of representations of Hermitian operators with deficiency index (m, m) , *Ukrain. Math. J.* **1** (1949) 3–66 (in Russian).
- [49] D. Krob and B. Leclerc, Minor identities for quasideterminants and quantum determinants, *Comm. Math. Phys.* **169** (1995) 1–23.
- [50] C. X. Li and J. J. C. Nimmo, Darboux transformations for a twisted derivation and quasideterminant solution to the super KdV equation, *Proc. Roy. Soc. A* **466** (2010) 2471–2493.
- [51] F. Marcellán and Y. Xu, On Sobolev orthogonal polynomials, *Expo. Math.* **33** (2015) 308–352.
- [52] A. S. Markus *Introduction to the Spectral Theory of Polynomials Operator Pencil*, Translated from the Russian by H. H. McFaden. Translation edited by Ben Silver. With an appendix by M. V. Keldysh. Translations of Mathematical Monographs, Vol. 71 (American Mathematical Society, Providence, RI, 1988).
- [53] P. Maroni, Sur quelques espaces de distributions qui sont des formes linéaires sur l'espace vectoriel des polynômes, in *Orthogonal Polynomials and Their Applications*, eds. C. Brezinski *et al.* Lecture Notes in Mathematics, Vol. 1171 (Springer-Verlag, 1985), pp. 184–194 (in French).
- [54] P. Maroni, Le calcul des formes linéaires et les polynômes orthogonaux semiclassiques, in *Orthogonal Polynomials and their Applications*, ed. M. Alfaro *et al.* Lecture Notes in Mathematics, Vol. 1329 (Springer-Verlag, 1988), pp. 279–290 (in French).
- [55] P. Maroni, Sur la suite de polynômes orthogonaux associée á la forme $u = \delta_c + \lambda(x - c)^{-1}L$, *Period. Math. Hungar.* **21** (1990) 223–248 (in French).
- [56] L. Miranian, Matrix valued orthogonal polynomials on the real line: Some extensions of the classical theory, *J. Phys. A: Math. Gen.* **38** (2005) 5731–5749.
- [57] A. F. Nikiforov, S. K. Suslov and V. B. Uvarov, *Classical Orthogonal Polynomials of a Discrete Variable* (Springer, Berlin, 1991).
- [58] P. J. Olver, On multivariate interpolation, *Stud. Appl. Math.* **116** (2006) 201–240.
- [59] O. Ore, Linear equations in non-commutative fields, *Ann. Math.* **32** (1931) 463–477.
- [60] V. Retakh and V. Rubtsov, Noncommutative toda chains, Hankel quasideterminants and Painlevé II equation, *J. Phys. A: Math. Theor.* **43** (2010) 505204.
- [61] A. R. Richardson, Hypercomplex determinants, *Messenger Math.* **55** (1926) 145–152.
- [62] A. R. Richardson, Simultaneous linear equations over a division algebra, *Proc. London Math. Soc.* **28** (1928) 395–420.
- [63] L. Rodman, Orthogonal matrix polynomials, in *Orthogonal Polynomials*, ed. P. Nevai, NATO Advanced Science Institute Series C. Mathematical and Physical Sciences, Vol. 294 (Kluwer, Dordrecht, 1990). pp. 345–362.
- [64] L. Rowen, *Ring Theory*, Vol. 1 (Academic Press, San Diego CA, 1988).
- [65] I. Schur, Über Potenzreihen, die im Innern des Einheitskreises beschränkt sind [I], *J. Reine Angew. Math.* **147** (1917) 205–232.
- [66] L. Schwartz, Théorie des noyaux, in *Proc. Int. Congr. Mathematicians* (Cambridge, MA, 1950), Vol. 1 (American Mathematical Society, Providence, RI, 1952) pp. 220–230.
- [67] L. Schwartz, *Théorie des Distributions* (Hermann, Paris, 1978).

- [68] B. Simon, The Christoffel–Darboux kernel, in *Perspectives in Partial Differential Equations, Harmonic Analysis and Applications*, eds. D. Mitrea and M. Mitrea, Proceedings of Symposia in Pure Mathematics, Vol. 79 (2008) 295–346.
- [69] A. Sinap and W. Van Assche, Polynomial interpolation and Gaussian quadrature for matrix-valued functions, *Linear Algebra Appl.* **207** (1994) 71–114.
- [70] A. Sinap and W. Van Assche, Orthogonal matrix polynomials and applications, *J. Computat. Appl. Math.* **66** (1996) 27–52.
- [71] G. Szegő, *Orthogonal Polynomials*, 4th edn., American Mathematical Society Colloquium Publication Series, Vol. 23 (American Mathematical Society, Providence RI, 1975).
- [72] F. Zhang (ed.), *The Schur Complement and its Applications* (Springer, New York, 2005).
- [73] A. Zhedanov, Rational spectral transformations and orthogonal polynomials, *J. Comput. Appl. Math.* **85** (1997) 67–86.