

Click here to view linked References

Order cones: A tool for deriving k -dimensional faces of cones of subfamilies of monotone games*

P. García-Segador

National Statistics Institute

Paseo de la Castellana, 183

28071 Madrid (Spain)

pedro.garcia.segador@ine.es

ORCID: 0000-0001-5079-4900

P. Miranda[†]

Complutense University of Madrid

Plaza de Ciencias, 3

28040 Madrid (Spain)

pmiranda@mat.ucm.es

ORCID: 0000-0002-4931-3636

Abstract

In this paper we introduce the concept of order cone. This concept is inspired by the concept of order polytopes, a well-known object coming from Combinatorics. Similarly to order polytopes, order cones are a special type of polyhedral cones whose geometrical structure depends on the properties of a partially ordered set (brief poset). This allows to study these properties in terms of the subjacent poset, a problem that is usually simpler to solve. From the point of view of applicability, it can be seen that many cones appearing in the literature of monotone TU-games are order cones. Especially, it can be seen that the cones of monotone games with restricted cooperation are order cones, no matter the structure of the set of feasible coalitions.

Keywords: Monotone games, restricted cooperation, order polytope, cone.

1 Introduction

Consider a finite set of n players $N = \{1, 2, \dots, n\}$. We will denote subsets of N by capital letters A, B, \dots and by $\mathcal{P}(N)$ the set of parts of N . A **game** v is a function $v : \mathcal{P}(N) \rightarrow \mathbb{R}$ satisfying $v(\emptyset) = 0$. The value $v(A)$ represents the minimal worth coalition A can obtain if all players in A agree to cooperate, no matter what players outside A might do.

In general, several additional conditions can be imposed on function v . One of the most natural conditions is monotonicity in v , i.e. $v(A) \leq v(B)$ if $A \subset B$. This means that if players add to a coalition, the corresponding worth increases. We will denote by $\mathcal{MG}(N)$ the set of all monotone games on N . Other popular conditions are additivity, supermodularity, and many others (see (Grabisch, 2016)).

On the other hand, it could be the case that some coalitions fail to form. Thus, v cannot be defined on some of the elements of $\mathcal{P}(N)$ and we have a subset $\mathcal{FC}(N)$ of $\mathcal{P}(N)$ containing all *feasible coalitions*. By a similar argument, coalitions with a fixed value may be left outside $\mathcal{FC}(N)$. From now on, we will not include \emptyset in $\mathcal{FC}(N)$. Usually, $\mathcal{FC}(N)$ has a concrete structure.

*This paper has been supported by the Ministry of Economy and Competitiveness of Spain under Grant PGC2018-095194-B-100 and by the Interdisciplinary Mathematical Institute of Complutense University.

[†]Corresponding author

1
2
3 If we fix an order on $\mathcal{FC}(N)$ (for example, the usual order on subsets of a set), a game v can be
4 identified to the point of $\mathbb{R}^{|\mathcal{FC}(N)|}$ given by $\mathbf{v} := (v(A))_{\{A:A \in \mathcal{FC}(N)\}}$. With some abuse of notation, we
5 will denote by v both the game and the corresponding point. Then, the set of games on N satisfying
6 a given condition (monotonicity, supermodularity, ...) and/or such that the set of feasible coalitions
7 is $\mathcal{FC}(N)$ can be seen as a set in $\mathbb{R}^{|\mathcal{FC}(N)|}$. In many cases, this set is usually a convex polyhedron.
8 Hence, it can be given in terms of its vertices and extremal rays. Following this line, many papers
9 have been devoted to solve the problem of obtaining different geometrical aspects of these polyhedra
10 for particular cases (see e.g. (Grabisch and Kroupa, 2019; Shapley, 1971)).

11
12 Continuing this line, in this paper, we introduce the concept of order cone. Order cones are
13 defined in terms of a poset and its structure relays on the structure of the corresponding poset.
14 Besides, we will show that order cones are deeply related to order polytopes. As many results are
15 known for order polytopes, it is possible to translate such properties to order cones.
16

17
18 As it will become clear below, order cones are a class of cones including the cones of monotone
19 games with restricted cooperation, no matter which the set $\mathcal{FC}(N)$ is. Thus, order cones allow to
20 study this set of cones in a general way. For example, we will characterize the set of extremal rays
21 of the cone $\mathcal{MG}(N)$, a problem that to our knowledge has not been solved yet (Grabisch, 2016).
22

23 Interestingly enough, order cones can be applied to other situations different to monotone games
24 with restricted cooperation. As an example dealing with such a case, we study the cone of monotone
25 k -symmetric games. This also adds more insight about the relationship between order cones and
26 order polytopes.
27

28 The rest of the paper goes as follows: In next section we introduce the basic concepts and results
29 about cones and order polytopes. Next, we define order cones and study some of its geometrical
30 properties. We then apply these results for some special cases of monotone games with restricted
31 cooperation. We finish with the conclusions and open problems.
32

33
34

35 2 Basic results

36
37 In order to be self-contained and fix the notation, let us start introducing some concepts and results
38 that will be needed throughout the paper.

39 A **cone** is a non-empty subset \mathcal{C} of \mathbb{R}^n such that if $\mathbf{x} \in \mathcal{C}$, then $\alpha \mathbf{x} \in \mathcal{C}$ for all $\alpha \geq 0$. Note that $\mathbf{0}$
40 is in any cone. Additionally, we say that the cone is **convex** if it is a convex set of \mathbb{R}^n ; equivalently,
41 a cone is convex if for any $\mathbf{x}, \mathbf{y} \in \mathcal{C}$, it follows
42

$$43 \mathbf{x} + \mathbf{y} \in \mathcal{C}.$$

44
45 Given a set \mathcal{S} , we define its **conic hull (or conic extension)** as the smallest cone containing \mathcal{S} .
46

47 A convex cone \mathcal{C} is **polyhedral** if additionally it is a polyhedron. This means that it can be
48 written as
49

$$50 \mathcal{C} := \{\mathbf{x} : A\mathbf{x} \leq \mathbf{0}\}, \tag{1}$$

51
52 for some matrix $A \in \mathcal{M}_{m \times n}$ of binding conditions. Two polyhedral cones are **affinely isomorphic**
53 if there is a bijective affine map from one cone onto the other. Given a polyhedral cone \mathcal{C} and
54 $\mathbf{x} \in \mathcal{C}$, $\mathbf{x} \neq \mathbf{0}$, the set $\{\alpha \mathbf{x} : \alpha \geq 0\}$ is called a **ray**. In general we will identify a ray with the point
55 \mathbf{x} . Notice also that for polyhedral cones, all rays pass through $\mathbf{0}$. Point \mathbf{x} defines an **extremal ray**
56 if $\mathbf{x} \in \mathcal{C}$ and there are $n - 1$ binding conditions for \mathbf{x} that are linearly independent. Equivalently, \mathbf{x}
57 cannot be written as a convex combination of two linearly independent points of \mathcal{C} .
58
59
60
61
62
63
64
65

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

It is well-known that a convex polyhedron only has a finite set of vertices and a finite set of extremal rays. The following result is well-known for convex polyhedra:

Theorem 1. *Let \mathcal{P} be a convex polyhedron on \mathbb{R}^n . Let us denote by $\mathbf{x}_1, \dots, \mathbf{x}_r$ the vertices of \mathcal{P} and by $\mathbf{v}_1, \dots, \mathbf{v}_s$ the vectors defining extremal rays. Then, for any $\mathbf{x} \in \mathcal{P}$, there exists $\alpha_1, \dots, \alpha_r$ such that $\alpha_1 + \dots + \alpha_r = 1, \alpha_i \geq 0, i = 1, \dots, r$, and β_1, \dots, β_s such that $\beta_i \geq 0, i = 1, \dots, s$, satisfying that*

$$\mathbf{x} = \sum_{i=1}^r \alpha_i \mathbf{x}_i + \sum_{j=1}^s \beta_j \mathbf{v}_j.$$

Given a polyhedral cone, if $\mathbf{x} \in \mathcal{C}, \mathbf{x} \neq \mathbf{0}$, it follows that \mathbf{x} cannot be a vertex of \mathcal{C} . Thus, for a polyhedral cone, the only possible vertex is $\mathbf{0}$. Thus, for the particular case of polyhedral cones, Theorem 1 writes as follows.

Corollary 1. *For a polyhedral cone \mathcal{C} whose extremal rays are defined by $\mathbf{v}_1, \dots, \mathbf{v}_s$, any $\mathbf{x} \in \mathcal{C}$ can be written as*

$$\mathbf{x} = \sum_{j=1}^s \beta_j \mathbf{v}_j, \quad \beta_j \geq 0, j = 1, \dots, s.$$

Consequently, in order to determine the polyhedral cone it suffices to obtain the extremal rays.

We will say that a cone is **pointed** if $\mathbf{0}$ is a vertex. The following result characterizes pointed cones.

Theorem 2. *For a polyhedral cone \mathcal{C} the following statements are equivalent:*

- \mathcal{C} is pointed.
- \mathcal{C} contains no line.
- $\mathcal{C} \cap (-\mathcal{C}) = \mathbf{0}$.

Finally, in this paper we will deal with the problem of obtaining the faces of order cones. Remember that given a polyhedron $\mathcal{P} \subseteq \mathbb{R}^n$, a non-empty subset $\mathcal{F} \subseteq \mathcal{P}$ is a **face** if there exist $\mathbf{v} \in \mathbb{R}^n, c \in \mathbb{R}$ such that

$$\mathbf{v}^t \mathbf{x} \leq c, \forall \mathbf{x} \in \mathcal{P}, \quad \mathbf{v}^t \mathbf{x} = c, \forall \mathbf{x} \in \mathcal{F}.$$

We denote this face as $\mathcal{F}_{\mathbf{v},c}$. The **dimension** of a face is the dimension of the smallest affine space containing the face. A common way to obtain faces is turning into equalities some of the inequalities of (1) defining \mathcal{P} .

Theorem 3. *(Cook et al., 1988) Let $A \in \mathcal{M}_{m \times n}$. Then any non-empty face of $\mathcal{P} = \{\mathbf{x} : A\mathbf{x} \leq \mathbf{b}\}$ corresponds to the set of solutions to*

$$\begin{aligned} \sum_j a_{ij} x_j &= b_i \text{ for all } i \in I \\ \sum_j a_{ij} x_j &\leq b_i \text{ for all } i \notin I, \end{aligned}$$

for some set $I \subseteq \{1, \dots, m\}$.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

The set of faces with the inclusion relation determines a lattice known as the **face lattice** of the polyhedron.

Let us now recall the basic results about order polytopes. Consider a poset (P, \preceq) , or P for short, with p elements. Elements of P are denoted x, y and so on. We will say that x is **covered** by y , denoted $x < y$, if $x \preceq y$ and there is no $z \in P \setminus \{x, y\}$ such that $x < z < y$. A subset $F \subseteq P$ is a *filter* if $x \in F$ and $x < y$ implies $y \in F$. We will denote by $\mathcal{F}(P)$ the set of filters of P . It is well-known that $(\mathcal{F}(P), \subseteq)$ is a distributive lattice (Davey and Priestley, 2002). Posets are usually represented through *Hasse diagrams*. A poset is connected if the corresponding Hasse diagram is a connected graph.

For any poset, it is possible to define a polytope on \mathbb{R}^p , called the order polytope of P .

Definition 1. (Stanley, 1986) Given a poset (P, \preceq) , we associate to P a polytope $\mathcal{O}(P)$ over \mathbb{R}^p , called the **order polytope** of P , formed by the p -uples f of real numbers indexed by the elements of P satisfying

- $0 \leq f(x) \leq 1$ for every element x in P , and
- $f(x) \leq f(y)$ whenever $x \preceq y$ in P .

Thus, the polytope $\mathcal{O}(P)$ consists in the order-preserving functions from P to $[0, 1]$. Note that we obtain an equivalent definition if the second condition turns into

$$f(x) \leq f(y) \text{ whenever } x < y.$$

The main advantage of order polytopes is that they allow to study the properties of the polytope in terms of the subjacent poset P . For example, if the poset is a chain, it can be shown that the corresponding order polytope is a **simplex**, i.e. a generalization of a triangle in the p -dimensional space.

Order polytopes has a tight relation with the set of capacities (see Definition 4 below). Indeed, it can be seen (Combarro and Miranda, 2010) that the set of fuzzy measures over a finite referential N of n elements, seen as a subset of $\mathbb{R}^{2^n - 2}$, is the order polytope with respect to the set $\mathcal{P}(N) \setminus \{\emptyset, N\}$ with the inclusion order. Other families of normalized measures are order polytopes, too, as for example the set of k -symmetric measures when the partition of indifference is known (Combarro and Miranda, 2010).

The following facts related to order polytopes are well-known and are discussed in (Stanley, 1986).

Proposition 1. Given a finite poset P , the vertices of $\mathcal{O}(P)$ are the characteristic functions v_F of filters F of P , i.e.

$$v_F(x) := \begin{cases} 1 & \text{if } x \in F \\ 0 & \text{otherwise} \end{cases}$$

Consequently, $\mathcal{O}(P)$ is a 0/1-polytope. Moreover,

$$\mathcal{O}(P) = \text{Conv} (v_F : F \subseteq P \text{ filter}),$$

and these points are in convex position, i.e., $v_F \notin \text{Conv} (v_{F'} : F \neq F' \subseteq P \text{ filter})$.

Next result characterizes whether two vertices are adjacent in $\mathcal{O}(P)$.

Theorem 4. Let P be a finite poset and consider two filters $F_1, F_2 \in \mathcal{F}(P)$. Then the vertices v_{F_1} and v_{F_2} are adjacent to each other if and only if $F_1 \subset F_2$ and $F_2 \setminus F_1$ is a connected subposet of P .

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

For obtaining the k -dimensional faces of an order polytope, additionally to the general methods presented for general polyhedrons, we can derive another way using the order structure of P . For this, we need to consider the poset

$$\hat{P} := \perp \oplus P \oplus \top,$$

where we have added to P a minimum \perp and a maximum \top . Then, $\mathcal{O}(P)$ is equivalent to the polytope given by

- $0 = f(\perp), f(\top) = 1$.
- $f(x) \leq f(y)$ whenever $x \preceq y$ in \hat{P} .

Now, note that turning an inequality of Definition 1 makes $f(x) = f(y)$ for some x, y such that $x \prec y$. Therefore, we can associate faces to partitions $\{B_1, \dots, B_k\}$ of \hat{P} in a way such that the face is the set of functions f such that $f(x) = f(y)$ for all x, y in the same block. However, not any partition defines a face. A partition $\{B_1, \dots, B_k\}$ is *connected* if B_i is connected as a subset of \hat{P} . Defining $B_i \prec B_j$ if there exists $x \in B_i, y \in B_j$ such that $x \preceq_P y$, we say that the partition is *compatible* if \prec is antisymmetric. Finally, the partition is *closed* if for $i \neq j$, there exists $g \in \mathcal{O}(P)$ constant in each block such that $g(B_i) \neq g(B_j)$. Now, the following holds.

Theorem 5. *A closed partition of \hat{P} defines a face of $\mathcal{O}(P)$ if and only if it is compatible and connected.*

This result is especially useful for high-dimensional faces as for example facets, as it is easy to check if these conditions on the partition hold. For faces of small dimension, we can solve the problem in another way. Note that any face can be defined equivalently as the convex hull of the vertices in the face. Hence, a face can be associated to its vertices. However, not every set of vertices defines a face. Thus, it suffices to obtain a condition for a subset of vertices to define a face. On the other hand, in order polytopes vertices are related to filters of P . If we focus on the set of filters defining a face, the following characterization arises.

Theorem 6. *(Friedl, 2017) Let $L \subseteq \mathcal{F}(P)$. Then, L determines a face if and only if L is an embedded lattice of $\mathcal{F}(P)$, i.e. for any two filters $F, F' \in \mathcal{F}(P)$*

$$J \cup J', J \cap J' \in L \Leftrightarrow J, J' \in L.$$

3 Order cones

Let us now turn to the concept of order cones. The idea is to remove the condition $f(a) \leq 1$ from Definition 1. Thus, the resulting set is no longer bounded. This is what we will call an order cone. Formally,

Definition 2. *Let P be a finite poset with p elements. The **order cone** $\mathcal{C}(P)$ is formed by the p -tuples f of real numbers indexed by the elements of P satisfying*

- i) $0 \leq f(x)$ for every $x \in P$,
- ii) $f(x) \leq f(y)$ whenever $x \preceq y$ in P .

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Besides, $\alpha \mathbf{w}_1, (1 - \alpha) \mathbf{w}_2 \in \mathcal{C}(P)$. Remark that \mathbf{w}_1 and \mathbf{w}_2 are linearly independent because there exists a coordinate i such that $v_i = 1$. Consequently, \mathbf{v} does not define an extremal ray, a contradiction.

Next, let us now show that \mathbf{v} is adjacent to $\mathbf{0}$. Otherwise, the segment $[\mathbf{0}, \mathbf{v}]$ is not an edge of $\mathcal{O}(P)$. Consequently, $\frac{1}{2}\mathbf{v}$ can be written as

$$\frac{1}{2}\mathbf{v} = \alpha \mathbf{y}_1 + (1 - \alpha) \mathbf{y}_2,$$

where $\mathbf{y}_1, \mathbf{y}_2 \in \mathcal{O}(P)$ such that they are outside $[\mathbf{0}, \mathbf{v}]$. Thus,

$$\mathbf{v} = 2\alpha \mathbf{y}_1 + 2(1 - \alpha) \mathbf{y}_2.$$

Finally, $2\alpha \mathbf{y}_1, 2(1 - \alpha) \mathbf{y}_2 \in \mathcal{C}(P)$, so we conclude that \mathbf{v} does not define an extremal ray, which is a contradiction.

Now, \mathbf{v} is related to a filter $F \subseteq P$. On the other hand, $\mathbf{0}$ is related to the empty filter. As \mathbf{v} is adjacent to $\mathbf{0}$, we can apply Theorem 4 to conclude that $F = F \setminus \emptyset$ is a connected filter of P .

Let us now prove the reverse. Consider \mathbf{v} an adjacent vertex to $\mathbf{0}$ in $\mathcal{O}(P)$ and assume that \mathbf{v} does not define an extremal ray. Then, there exists $\mathbf{w}_1, \mathbf{w}_2 \in \mathcal{C}(P)$ and not proportional to \mathbf{v} such that

$$\mathbf{v} = \mathbf{w}_1 + \mathbf{w}_2 = \frac{1}{2}2\mathbf{w}_1 + \frac{1}{2}2\mathbf{w}_2 = \frac{1}{2}\mathbf{w}'_1 + \frac{1}{2}\mathbf{w}'_2.$$

Now, for $\epsilon > 0$ small enough, we have

$$\epsilon \mathbf{v} = \frac{1}{2}\epsilon \mathbf{w}'_1 + \frac{1}{2}\epsilon \mathbf{w}'_2,$$

and $\epsilon \mathbf{w}'_1 \leq \mathbf{1}, \epsilon \mathbf{w}'_2 \leq \mathbf{1}$. Hence, $\epsilon \mathbf{w}'_1, \epsilon \mathbf{w}'_2 \in \mathcal{O}(P)$ by Lemma 3, and hence $[\mathbf{0}, \mathbf{v}]$ is not an edge of $\mathcal{O}(P)$, in contradiction with \mathbf{v} adjacent to $\mathbf{0}$. □

Let us now turn to the problem of obtaining the faces of $\mathcal{C}(P)$. As explained in Theorem 3, faces arise when inequalities turn into equalities. Let us consider the inequality $f(x) \leq f(y)$ for $x \preceq y$ and assume this inequality is turned into an equality. This means that x and y identified to each other; let us call z this new element. In terms of posets, this translate into transforming P into another poset (P', \preceq') defined as $P' := P \setminus \{x, y\} \cup \{z\}$ and \preceq' given by:

$$\begin{cases} a \preceq' b \Leftrightarrow a \preceq b & \text{if } a, b \neq z \\ z \preceq' b \Leftrightarrow x \preceq b \\ a \preceq' z \Leftrightarrow a \preceq y \end{cases}$$

Similar conclusions arise when $0 \leq f(x)$ turns into an equality. Moreover, if \mathcal{F} is the face obtained by turning inequalities into equalities, the projection

$$\pi : \begin{array}{ccc} \mathcal{F} & \rightarrow & \mathcal{C}(P') \\ (f(a), \dots, f(x), f(y), \dots, f(b)) & \leftrightarrow & (f(a), \dots, f(x), f(y), \dots, f(b)) \end{array}$$

is a bijective affine map. Consequently, the following holds.

Lemma 4. *The faces of an order cone are affinely isomorphic to order cones.*

Compare this result with the corresponding result for order polytopes (Stanley, 1986).

1

2

3 **Lemma 5.** For an order cone $\mathcal{C}(P)$, the vertex $\mathbf{0}$ is in all non-empty faces. Consequently, all faces
4 can be written as $\mathcal{F}_{\mathbf{v},0}$.

5

6 *Proof.* It suffices to show that for a non-empty face $\mathcal{F}_{\mathbf{v},c}$, it is $c = 0$. First, $\mathbf{v}^t \mathbf{0} \leq c$, so that $c \geq 0$.

7 Suppose $c > 0$. As $\mathcal{F}_{\mathbf{v},c}$ is non-empty, there exist $\mathbf{x} \in \mathcal{C}(P)$ such that $\mathbf{v}^t \mathbf{x} = c$. But then,
8 $\mathbf{v}^t 2\mathbf{x} = 2c > c$, a contradiction. Thus, $c = 0$ and $\mathbf{0} \in \mathcal{F}_{\mathbf{v},0}$. \square

9

10 With this in mind, Theorem 7 can be extended to characterize all the faces of the order cone, not
11 only the extremal rays.

12

13 **Theorem 8.** Let P be a finite poset and $\mathcal{C}(P)$ and $\mathcal{O}(P)$ the corresponding order cone and order
14 polytope, respectively. For a pair $(\mathbf{v}, 0)$, the set $\mathcal{F}'_{\mathbf{v},0} = \mathcal{C}(P) \cap \{\mathbf{x} : \mathbf{v}^t \mathbf{x} = 0\}$ is a face of $\mathcal{C}(P)$ if and
15 only if $\mathcal{F}_{\mathbf{v},0} = \mathcal{O}(P) \cap \{\mathbf{x} : \mathbf{v}^t \mathbf{x} = 0\}$ is a face of $\mathcal{O}(P)$. Moreover, $\dim(\mathcal{F}'_{\mathbf{v},0}) = \dim(\mathcal{F}_{\mathbf{v},0})$.

16

17 *Proof.* Let $\mathcal{F}_{\mathbf{v},0}$ be a face of $\mathcal{O}(P)$ containing $\mathbf{0}$ and let us show that it determines a face on $\mathcal{C}(P)$.
18 First, let us show that $\mathbf{v}^t \mathbf{x} \leq 0, \forall \mathbf{x} \in \mathcal{C}(P)$. Otherwise, there exists $\mathbf{x}_0 \in \mathcal{C}(P)$ such that $\mathbf{v}^t \mathbf{x}_0 > 0$.
19 But then $\mathbf{v}^t \epsilon \mathbf{x}_0 > 0, \forall \epsilon > 0$. As ϵ can be taken small enough so that $\epsilon \mathbf{x}_0 \leq \mathbf{1}$, it follows by Lemma 3
20 that $\epsilon \mathbf{x}_0 \in \mathcal{O}(P)$ and as $\mathbf{v}^t \epsilon \mathbf{x}_0 > 0$, and we get a contradiction. Hence, the pair $(\mathbf{v}, 0)$ determines a
21 face $\mathcal{F}'_{\mathbf{v},0}$ of $\mathcal{C}(P)$.

22 Consider now a face $\mathcal{F}'_{\mathbf{v},0}$ of $\mathcal{C}(P)$. Hence, $\mathbf{v}^t \mathbf{x} \leq 0, \forall \mathbf{x} \in \mathcal{C}(P)$. But then, $\mathbf{v}^t \mathbf{x} \leq 0, \forall \mathbf{x} \in \mathcal{O}(P)$
23 and as $\mathbf{0} \in \mathcal{F}'_{\mathbf{v},0}$, this determines a face of $\mathcal{O}(P)$.

24 Let us now see that for each pair $(\mathbf{v}, 0)$, $\dim(\mathcal{F}'_{\mathbf{v},0}) = \dim(\mathcal{F}_{\mathbf{v},0})$. First, as $\mathcal{F}_{\mathbf{v},0} \subseteq \mathcal{F}'_{\mathbf{v},0}$, we have
25 $\dim(\mathcal{F}_{\mathbf{v},0}) \leq \dim(\mathcal{F}'_{\mathbf{v},0})$.

26 On the other hand, let k be the dimension of $\mathcal{F}'_{\mathbf{v},0}$. This implies that there are k vectors $\mathbf{v}_1, \dots, \mathbf{v}_k$
27 linearly independent in $\mathcal{F}'_{\mathbf{v},0}$. But now, we can find $\epsilon > 0$ small enough such that $\epsilon \mathbf{v}_1 \leq \mathbf{1}, \dots, \epsilon \mathbf{v}_k \leq \mathbf{1}$.
28 Thus, $\epsilon \mathbf{v}_1, \dots, \epsilon \mathbf{v}_k \in \mathcal{F}_{\mathbf{v},0}$ and hence, $\dim(\mathcal{F}_{\mathbf{v},0}) \geq \dim(\mathcal{F}'_{\mathbf{v},0})$. \square

29

30 As a consequence, we can adapt Theorem 6 for order cones as follows.

31

32 **Theorem 9.** Let $L \subseteq \mathcal{F}(P)$. Then, L determines a face of $\mathcal{C}(P)$ if and only if L is an embedded
33 lattice of $\mathcal{F}(P)$ containing the empty filter.

34

35 **Remark 1.** From Theorem 8, in order to find faces of an order cone, we need to look for faces of
36 the corresponding order polytope containing $\mathbf{0}$. As previously explained in Theorem 3, if we consider
37 the expression of $\mathcal{O}(P)$ as a polyhedron, faces arise turning inequalities into equalities. Vertices in
38 the face are the vertices of the polyhedron satisfying these equalities. If we consider \hat{P} , vertex $\mathbf{0}$
39 corresponds to function

40

$$41 \quad f(x) = \begin{cases} 0 & x \neq \top \\ 1 & x = \top \end{cases}$$

42

43 Consequently, $\mathbf{0}$ satisfies $f(x) = f(y)$ when $y \neq \top$. Thus, we look for the faces where the
44 inequalities turned into equalities do not depend on \top .

45 In terms of Theorem 5, we have to look for partitions defining faces containing $\mathbf{0}$. Note that each
46 block B_i defines a subset of P such that all elements in B_i attain the same value for all points in the
47 face. Therefore, faces containing $\mathbf{0}$ mean that there is a block containing only \top .

48

49 **Example 1.** Consider the polytope given in Figure 1 left.

50 In this case, we have three elements and both the order polytope and the cone order cone can be
51 depicted in \mathbb{R}^3 , with the first coordinate corresponding to 1, the second one to 2 and the third to 12,
52 see Figure 2. The cone $\mathcal{C}(P)$ is given by 3-dimensional vectors f satisfying

53

54

55

56




Figure 1: Example of poset P (left), his extension \hat{P} (center) and his filter lattice (right).

$$0 \leq f(1), 0 \leq f(2), f(1) \leq f(12), f(2) \leq f(12).$$

Let us then explain the previous results for this poset. First, let us start obtaining the vectors defining extremal rays. According to Theorem 7, it suffices to obtain the non-empty filters that are connected subsets of P . Non-empty filters of P are:

$$\{\{12\}, \{1, 12\}, \{2, 12\}, \{1, 2, 12\}\}.$$

All of them are connected subsets of P . Hence, we have 4 extremal rays, whose respective vectors are

$$(0, 0, 1), (1, 0, 1), (0, 1, 1), (1, 1, 1).$$




Figure 2: Order polytope $\mathcal{O}(P)$ (left) and order cone $\mathcal{C}(P)$ (right).

Let us now deal with the facets. For this, consider the poset $\hat{P} = \perp \oplus P \oplus \top$ (see Figure 1 center). According to Theorems 5 and 8, the facets are given by considering one of the following equalities:

$$f(\perp) = f(1), f(\perp) = f(2), f(1) = f(1, 2), f(2) = f(1, 2), f(1, 2) = f(\top).$$

This translates into transforming poset \hat{P} in a new poset where the elements in the equality identify to each other (see Lemma 4). The posets for the previous equalities are given in Figure 3.

Note that the facets containing $\mathbf{0}$ are those whose defining equality does not involve \top , as $\mathbf{0}$ satisfies any other equality. In our case, they correspond to the first four cases. Thus, we have four




Figure 3: Subposets when turning an inequality into an equality.

facets containing $\mathbf{0}$ and all of them are simplices (indeed triangles) because the corresponding polytope is a chain.

For the 1-dimensional faces, we have to consider two equalities. However, we have to be careful with the selected equalities because they might imply other equalities. For example, if we consider $f(\perp) = f(1), f(1) = f(1, 2)$, this also implies $f(\perp) = f(2)$, and hence we obtain a point instead of an edge. In our case, the edges containing $\mathbf{0}$ are given by the pairs of equalities defining an edge and not involving \top . There are four pairs in these conditions that are

$$\{f(\perp) = f(1), f(\perp) = f(2)\}, \{f(\perp) = f(1), f(2) = f(1, 2)\},$$

$$\{f(\perp) = f(2), f(1) = f(1, 2)\}, \{f(1) = f(1, 2), f(2) = f(1, 2)\}.$$

Alternatively, we could use the characterization given in Theorem 9. In this case, we have to consider the filter lattice (see Figure 1 right).

Hence, edges are given by pairs of filters defining a sublattice and involving the empty filter. Thus, the possible choices are the following pairs:

$$\{\{\emptyset\}, \{12\}\}, \{\{\emptyset\}, \{1, 12\}\}, \{\{\emptyset\}, \{2, 12\}\}, \{\{\emptyset\}, \{1, 2, 12\}\}.$$

Thus, the extremal rays of $\mathcal{C}(P)$ are given by vectors $(0, 0, 1), (1, 0, 1), (0, 1, 1), (1, 1, 1)$.

For 2-dimensional faces, we have to consider all possible sublattices of height 2 and involving the filter. These sublattices are:

$$\{\{\emptyset\}, \{1, 12\}, \{1, 2, 12\}\}, \{\{\emptyset\}, \{2, 12\}, \{1, 2, 12\}\}, \{\{\emptyset\}, \{12\}, \{1, 12\}\}, \{\{\emptyset\}, \{12\}, \{2, 12\}\}.$$

Hence, the 2-dimensional faces for $\mathcal{C}(P)$ are defined by vectors

$$\{(1, 0, 1), (1, 1, 1)\}, \{(0, 1, 1), (1, 1, 1)\}, \{(0, 0, 1), (1, 0, 1)\}, \{(0, 0, 1), (0, 1, 1)\}.$$

Notice that we cannot consider

$$\{\{\emptyset\}, \{1, 12\}, \{2, 12\}, \{1, 2, 12\}\}, \{\{\emptyset\}, \{12\}, \{1, 12\}, \{2, 12\}\},$$

because they are not embedded sublattices.

4 Application to Game Theory

In this section, we show that some well-known cones appearing in the field of monotone games can be seen as order cones. Hence, all the results developed in the previous section can be applied to these cones. The first example deals with the general case of monotone games when all coalitions are feasible. We next extend this to the case where $\mathcal{FC}(N) \subset \mathcal{P}(N) \setminus \{\emptyset\}$. As an example of applicability for subfamilies of monotone games satisfying a property on v but not on the set of feasible coalitions, we also treat the case of k -symmetric monotone games.

4.1 The cone of general monotone games

Consider monotone games when all coalitions are feasible, i.e. the set $\mathcal{MG}(N)$. We consider $\mathcal{MG}(N)$ as a subset of $\mathbb{R}^{2^n - 1}$ (we have removed the coordinate for \emptyset because its value is fixed). This set is given by all games satisfying $v(A) \leq v(B)$ whenever $A \subset B$. Thus, a game $v \in \mathcal{MG}(N)$ is characterized by the following conditions:

- $0 \leq v(A)$.
- $v(A) \leq v(B)$ if $A \subseteq B$.

Then, $\mathcal{MG}(N) = \mathcal{C}(\mathcal{P}(N) \setminus \{\emptyset\})$, where the order relation \prec on $\mathcal{P}(N) \setminus \{\emptyset\}$ is given by $A \prec B$ if and only if $A \subset B$. For example, for $|N| = 3$, this poset is given in Figure 4. However, little else is known about $\mathcal{MG}(N)$; for example, the set of extremal rays is not known and this question appears in (Grabisch, 2016) as an open problem. We will study this set at the light of the results of the previous section. Let us first deal with the extremal rays.

Corollary 2. *The vectors defining an extremal ray of $\mathcal{MG}(N)$ are defined by non-empty filters of $\mathcal{P}(N) \setminus \{\emptyset\}$.*

Proof. Following Theorem 7, we need to find the filters of $\mathcal{P}(N) \setminus \{\emptyset\}$ that are connected. But in this case, all filters except the empty filter, corresponding to vertex $\mathbf{0}$, contain N . Hence, all of them are connected. \square

For obtaining the number of extremal rays, note that any filter in a poset is characterized in terms of its minimal elements and that these minimal elements are an antichain of the poset. For the boolean poset $\mathcal{P}(N)$, the number of antichains is known as the Dedekind numbers, D_n . The first values of D_n are given in Table 1.

For $\mathcal{MG}(N)$, we have to remove the antichain $\{\emptyset\}$ because the poset defining the order cone is $\mathcal{P}(N) \setminus \{\emptyset\}$. Besides, the empty antichain corresponds to $\mathbf{0}$ and thus, it should be removed, too. Hence, the number of extremal rays of $\mathcal{MG}(N)$ is $D_n - 2$.

Example 2. *Let us compute the extremal rays of the order cone $\mathcal{MG}(N)$ where $N = \{1, 2, 3\}$. Note that $\mathcal{C}(P)$ is a cone in \mathbb{R}^7 . Then, considering $P = B_3 \setminus \{\emptyset\}$, it suffices to compute the filters of P .*

A list with these filters is:

$$\begin{aligned} \mathfrak{CF}(P) = & \{\emptyset, \{123\}, \{12, 123\}, \{13, 123\}, \{23, 123\}, \{12, 13, 123\}, \{12, 23, 123\}, \{13, 23, 123\}, \\ & \{12, 13, 23, 123\}, \{1, 12, 13, 123\}, \{1, 12, 13, 23, 123\}, \{2, 12, 23, 123\}, \{2, 12, 13, 23, 123\}, \{3, 13, 23, 123\}, \\ & \{3, 12, 13, 23, 123\}, \{1, 2, 12, 13, 23, 123\}, \{1, 3, 12, 13, 23, 123\}, \{2, 3, 12, 13, 23, 123\}, \{1, 2, 3, 12, 13, 23, 123\}\}. \end{aligned}$$

Removing \emptyset , we have a total of 18 extremal rays. Note that $D_3 = 20$.

n	$M(n)$
0	2
1	3
2	6
3	20
4	168
5	7 581
6	7 828 354
7	2 414 682 040 998
8	56 130 437 228 687 557 907 788

Table 1: First Dedekind numbers.




Figure 4: Boolean poset $P = B_3 \setminus \{\emptyset\}$.

Similarly, we can apply Theorem 8 to obtain all k -dimensional faces of the cone $\mathcal{MG}(N)$.

Corollary 3. *The non-empty faces of $\mathcal{MG}(N)$ are given by the non-empty faces of $\mathcal{O}(\mathcal{P}(N) \setminus \{\emptyset\})$ containing vertex $\mathbf{0}$.*

However, we will see that in this case we can do better.

Definition 3. *Let \mathcal{P} be a convex polytope and \mathbf{x} be a point outside the affine space generated by \mathcal{P} , denoted $\text{aff}(\mathcal{P})$. Point \mathbf{x} is called **apex**. We define a **pyramid** with base \mathcal{P} and apex \mathbf{x} , denoted by $\text{pyr}(\mathcal{P}, \mathbf{x})$, as the polytope whose vertices are the ones of \mathcal{P} and \mathbf{x} .*

Note that for a pyramid $\text{pyr}(\mathcal{P}, \mathbf{x})$, any vertex of \mathcal{P} is adjacent to \mathbf{x} . Moreover, there is a simple way to find faces containing \mathbf{x} for a pyramid that we write below.

Proposition 2. *For a pyramid of apex \mathbf{x} and base \mathcal{P} , the k -dimensional faces containing \mathbf{x} are given by the $(k - 1)$ -dimensional faces of \mathcal{P} .*

From now on, in order to simplify the notation, we will assume that the last coordinate in vector \mathbf{v} corresponds to the value $v(N)$.

Proposition 3. *Consider the poset $\mathcal{P}(N) \setminus \{\emptyset\}$ with the relation order $A \prec B \Leftrightarrow A \subset B$. Then, the order polytope $\mathcal{O}(\mathcal{P}(N) \setminus \{\emptyset\})$ is a pyramid with apex $\mathbf{0}$ and base $\{(\mathbf{x}, 1) : \mathbf{x} \in \mathcal{O}(\mathcal{P}(N) \setminus \{\emptyset, N\})\}$.*

Proof. Note that for any non-empty filter F , it follows that $N \in F$. Then, the characteristic function of any non-empty filter v_F satisfies $v_F(N) = 1$. Hence, any vertex of $\mathcal{O}(\mathcal{P}(N) \setminus \{\emptyset\})$ except $\mathbf{0}$ is in the hyperplane $v(N) = 1$. Consequently, $\mathcal{O}(\mathcal{P} \setminus \{\emptyset\})$ is a pyramid with apex $\mathbf{0}$. Finally, the points

1
2
3 \mathbf{v} of $\mathcal{O}(\mathcal{P}(N)\setminus\{\emptyset\})$ in the hyperplane $v(N) = 1$ satisfy $v(A) \leq v(B)$ if $A \subseteq B$. Thus, these points
4 can be associated to the order polytope $\mathcal{O}(\mathcal{P}(N)\setminus\{\emptyset, N\})$, where the order relation \preceq is given by
5 $A \preceq B \Leftrightarrow A \subseteq B$. □

6
7 This allows us to study the k -dimensional faces of $\mathcal{MG}(X)$ from a different point of view that
8 the one of Theorems 9 and 8. In particular, as apex \mathbf{x} is adjacent to every vertex in the base \mathcal{P} ,
9 edges are given by segments $[\mathbf{x}, \mathbf{y}]$ with \mathbf{y} a vertex of \mathcal{P} , thus recovering the result of Corollary 2. In
10 general, applying Proposition 2, the following holds.

11
12
13 **Corollary 4.** *The k -dimensional faces of $\mathcal{MG}(N)$ are given by the $(k - 1)$ -dimensional faces of*
14 *$\mathcal{O}(\mathcal{P}(N)\setminus\{\emptyset, N\})$.*

15
16 The order polytope $\mathcal{O}(\mathcal{P}(N)\setminus\{\emptyset, N\})$ is a well-known polytope corresponding to the set of capac-
17 ities or fuzzy measures.

18
19
20 **Definition 4.** *A capacity on X is a map $\mu : \mathcal{P}(N) \rightarrow \mathbb{R}$, satisfying*

- 21
22 *i) $\mu(\emptyset) = 0, \mu(N) = 1$ (normalization).*
23
24 *ii) $\mu(A) \leq \mu(B), \forall A \subseteq B$ (monotonicity).*

25
26
27 This notion was proposed by Choquet (Choquet, 1953) and independently by Sugeno under
28 the name of fuzzy measure (Sugeno, 1974). These measures are also called “non-additive mesasures”
29 (Denneberg, 1994). From the point of view of Game Theory, capacities are just normalized monotone
30 games. Capacities constitute an extension of a probability distribution, where additivity is turned
31 into monotonicity and they have been applied in many different fields, as for example Decision Making
32 (see (Grabisch, 2016) and references therein). The set of capacities on a referential set N is denoted
33 by $\mathcal{FM}(N)$ and it can be seen (Combarro and Miranda, 2010) that

$$\mathcal{FM}(N) = \mathcal{O}(\mathcal{P}(N)\setminus\{\emptyset, N\}).$$

34
35
36
37
38
39 It is worth-noting that the geometrical structure (apart the dimension) of $\mathcal{O}(\mathcal{P}(N)\setminus\{\emptyset, N\})$ is
40 quite different from the geometrical structure of $\mathcal{O}(\mathcal{P}(N)\setminus\{\emptyset\})$. For example, in $\mathcal{O}(\mathcal{P}(N)\setminus\{\emptyset\})$ all
41 vertices are adjacent to $\mathbf{0}$, while this is not the case for $\mathcal{O}(\mathcal{P}(N)\setminus\{\emptyset, N\})$ (see (Combarro and Miranda,
42 2010)).

43
44 For this order polytope, many results are known, as for example whether two vertices are adjacent
45 or the centroid (Combarro and Miranda, 2010, 2008). Applying Corollary 4, we conclude that 2-
46 dimensional faces of $\mathcal{MG}(N)$ are given by an edge of $\mathcal{FM}(N) = \mathcal{O}(\mathcal{P}(N)\setminus\{\emptyset, N\})$. On the other
47 hand, an edge in $\mathcal{O}(\mathcal{P}(N)\setminus\{\emptyset, N\})$ is given by two adjacent vertices $\mathbf{v}_{F_1}, \mathbf{v}_{F_2}$. Another characterization
48 specific for $\mathcal{O}(\mathcal{P}(N)\setminus\{\emptyset, N\})$ is given in (Combarro and Miranda, 2008). Moreover, as both F_1, F_2
49 are adjacent to $\mathbf{0}$, the following holds.

50
51
52
53 **Corollary 5.** *Any 2-dimensional face of $\mathcal{MG}(N)$ are defined in terms of 2-dimensional simplices*
54 *given by $\{\mathbf{0}, \mathbf{v}_{F_1}, \mathbf{v}_{F_2}\}$ where $F_2 \setminus F_1$ is a connected subposet of $\mathcal{P}(N)\setminus\{\emptyset, N\}$.*

55
56 **Example 3.** *Continuing with the previous example, the previous discussion allows to derive the 2-*
57 *dimensional faces of $\mathcal{MG}(N)$, as by Corollary 4 they can be given in terms of edges of $\mathcal{O}(\mathcal{P}(N)\setminus\{\emptyset, N\})$.*
58 *The filters of $\mathcal{P}(N)\setminus\{\emptyset, N\}$ are:*

$$\mathcal{F}(P) = \{\emptyset, \{12\}, \{13\}, \{23\}, \{12, 13\}, \{12, 23\}, \{13, 23\},$$

1
2
3 $\{12, 13, 23\}, \{1, 12, 13\}, \{1, 12, 13, 23\}, \{2, 12, 23\}, \{2, 12, 13, 23\}, \{3, 13, 23\},$
4 $\{3, 12, 13, 23\}, \{1, 2, 12, 13, 23\}, \{1, 3, 12, 13, 23\}, \{2, 3, 12, 13, 23\}, \{1, 2, 3, 12, 13, 23\}$.
5

6 *Now, we have to search for pairs of adjacent vertices in $\mathcal{O}(\mathcal{P}(N) \setminus \{\emptyset, N\})$, for example using*
7 *Theorem 4. It is easy but tedious to show that there are 76 pairs in these conditions.*
8
9

10 4.2 The cone of games with restricted cooperation

11 Let us now treat the problem when we face a situation of restricted cooperation. Then, several
12 coalitions are not allowed and we have a set $\mathcal{FC}(N) \subset \mathcal{P}(N) \setminus \{\emptyset\}$ of feasible coalitions. Many papers
13 have been devoted to this subject, usually imposing an algebraic structure on $\mathcal{FC}(N)$ (see e.g. (Faigle,
14 1989; Pulido and Sánchez-Soriano, 2006; Katsev and Yanovskaya, 2013; Grabisch, 2011)). From the
15 point of view of polyhedra, if a coalition is not feasible, this implies that this subset is removed from
16 $\mathcal{FC}(N)$. We will denote by $\mathcal{MG}_{\mathcal{FC}(N)}(N)$ the set of all monotone games whose feasible coalitions are
17 $\mathcal{FC}(N)$. Thus, a game $v \in \mathcal{MG}_{\mathcal{FC}(N)}(N)$ is characterized by the following conditions:
18
19

- 20 • $0 \leq v(A), A \in \mathcal{FC}(N)$.
- 21 • $v(A) \leq v(B)$ if $A \subseteq B, A, B \in \mathcal{FC}(N)$.

22 Then, $\mathcal{MG}_{\mathcal{FC}(N)}(N) = \mathcal{C}(\mathcal{FC}(N))$, where the order relation \prec on $\mathcal{FC}(N)$ is given by $A \prec B$ if
23 and only if $A \subset B$.

24 Assume first that $N \in \mathcal{FC}(N)$. This is the usual situation, as most of the solution concepts on
25 Game Theory assume that all players agree to form the grand coalition (see e.g. (Grabisch, 2013)).
26 In this case, the following holds.
27

28 **Corollary 6.** *If $N \in \mathcal{FC}(N)$, then the set of extremal rays of $\mathcal{MG}_{\mathcal{FC}(N)}(N)$ is given by*

$$29 \{v_F : \emptyset \neq F, F \text{ filter of } \mathcal{FC}(N)\}.$$

30 *Proof.* Applying Theorem 7, the set of extremal rays is given by the set of vertices v_F of $\mathcal{O}(\mathcal{FC}(N))$
31 such that F is a connected filter in $\mathcal{FC}(N)$. As $N \in \mathcal{FC}(N)$, it follows that all filters are connected
32 subposets of $\mathcal{FC}(N)$, so that we have as many extremal rays as vertices in $\mathcal{O}(\mathcal{FC}(N))$ different from
33 vertex $\mathbf{0}$. And this value is given by the number of filters minus one (for the empty filter corresponding to
34 vertex $\mathbf{0}$). \square
35

36 Indeed, we can translate in this case the results obtained for $\mathcal{MG}(N)$. Assuming the last coordi-
37 nate corresponds to subset N , the following holds.
38

39 **Proposition 4.** *Assume $N \in \mathcal{FC}(N)$ and consider the poset $\mathcal{FC}(N)$ with the relation order $A \prec$
40 $B \Leftrightarrow A \subset B$. Then, the order polytope $\mathcal{O}(\mathcal{FC}(N))$ is a pyramid with apex $\mathbf{0}$ and base $\{(\mathbf{x}, 1) : \mathbf{x} \in$
41 $\mathcal{O}(\mathcal{FC}(N) \setminus \{N\})\}$.
42*

43 *Proof.* It is a straightforward translation of the proof of Proposition 3. \square
44

45 This implies that have two possibilities for studying $\mathcal{MG}_{\mathcal{FC}(N)}(N)$. First, we can apply the general
46 results for any order cones developed in Section 3. Alternatively, we can apply Proposition 2 and
47 derive the results from the structure of the order polytope $\mathcal{O}(\mathcal{FC}(N) \setminus \{N\})$ just as it has been done
48 for $\mathcal{MG}(N)$. In this last case, the following holds.
49




Figure 5: Hasse diagram of the poset of a game with restricted cooperation.

Corollary 7. *The k -dimensional faces of $\mathcal{MG}(\mathcal{FC}(N))$ are given by the $(k - 1)$ -dimensional faces of $\mathcal{O}(\mathcal{FC}(N) \setminus \{N\})$.*

Example 4. *Suppose a situation with four players, and assume that the only feasible coalitions are $\mathcal{FC}(N) = \{12, 23, 34, 1234\}$. The corresponding Hasse diagram is given in Figure 5.*

For this example, the non-empty filters of $\mathcal{FC}(N)$ are:

$$F_1 = \{1234\}, F_2 = \{12, 1234\}, F_3 = \{23, 1234\}, F_4 = \{34, 1234\}, F_5 = \{12, 23, 1234\},$$

$$F_6 = \{12, 34, 1234\}, F_7 = \{23, 34, 1234\}, F_8 = \{12, 23, 34, 1234\}.$$

Thus, we have 8 extremal rays. For example, the extremal ray corresponding to F_5 is given by vector $\mathbf{v} = (1, 1, 0, 1)$, where the third coordinate corresponds to subset $\{34\}$.

For k -dimensional faces, it just suffice to note that $\mathcal{FC}(N) \setminus \{1234\}$ is an antichain. Then, $\mathcal{O}(\mathcal{FC}(N) \setminus \{N\})$ is a cube. For example, for finding 2-dimensional faces, we have to consider pairs of adjacent vertices of the cube $\mathcal{O}(\mathcal{FC}(N) \setminus \{N\})$ (there are 12 pairs). Similarly, for 3-dimensional faces we have to consider 2-dimensional faces of the cube (six cases), and there is just one 4-dimensional face.

Now, assume $N \notin \mathcal{FC}(N)$. This situation is more tricky and needs to study each case applying Theorems 7 and 8. For example, in this situation it could happen that some vertices are not adjacent to $\mathbf{0}$ and thus, they do not define an extremal ray. Moreover, the 2-dimensional faces are not defined necessarily via 2-dimensional simplices.

As examples for this case, we study two situations. Assume $\mathcal{FC}(N) \cup \{\emptyset\}$ is a poset with a top element \top and thus, we can extend all the results that we have obtained when $N \in \mathcal{FC}(N)$.

Proposition 5. *Consider the poset with top element $\mathcal{FC}(N) \cup \{\emptyset\}$ with the relation order $A \prec B \Leftrightarrow A \subset B$ and top element \top . Then, the order polytope $\mathcal{O}(\mathcal{FC}(N) \setminus \{\emptyset\})$ is a pyramid with apex $\mathbf{0}$ and base $\{(\mathbf{x}, 1) : \mathbf{x} \in \mathcal{O}(\mathcal{FC}(N) \setminus \{\top\})\}$.*

Corollary 8. *The k -dimensional faces of $\mathcal{MG}(\mathcal{FC}(N))$ are given by the $(k - 1)$ -dimensional faces of $\mathcal{O}(\mathcal{FC}(N) \setminus \{\top\})$.*

Suppose as a second example that $\mathcal{FC}(N)$ is a union of connected posets

$$\mathcal{FC}(N) = P_1 \cup \dots \cup P_r, \quad P_i \text{ connected.}$$


In this case, the only connected filters are the connected filters $F_i \subseteq P_i$. Then, we have:

Proposition 6. *If $\mathcal{FC}(N) = P_1 \cup \dots \cup P_r$, where P_i is a connected poset, $i = 1, \dots, r$, then the extremal rays of $\mathcal{MG}(\mathcal{FC}(N))$ are given by \mathbf{v}_{F_i} where F_i is a non-empty connected filter of P_i .*

1
2 For example, if $|P_i| = 1 \forall i$, then $\mathcal{FC}(N)$ is an antichain and the only connected filters are the
3 singletons. Thus, there are just r extremal rays for $\mathcal{MG}(\mathcal{FC}(N))$. Indeed, note that the corresponding
4 order polytope is the r -dimensional cube and thus the vertices adjacent to $\mathbf{0}$ are $\mathbf{e}_i, i = 1, \dots, r$.

5
6 In general, we have to study the properties of the corresponding poset.

7
8 **Example 5.** Assume again a 4-players game and let us consider the coalitions given in Figure 6 left.
9 We have in this case a 4-dimensional cone order.



10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27 Figure 6: Hasse diagram of the poset P of a game with restricted cooperation (left) and his extension
28 \hat{P} .

29
30 Fixing the order for coordinates 12, 13, 34, 123, the vertices of the corresponding order polytope
31 are given in Table 2.

Filter	\emptyset	123	34	34, 123	12, 123
Vertex	(0,0,0,0)	(0,0,0,1)	(0,0,1,0)	(0,0,1,1)	(1,0,0,1)
Filter	13, 123	12, 12, 123	12, 34, 123	13, 34, 123	12, 13, 34, 123
Vertex	(0,1,0,1)	(1,1,0,1)	(1,0,1,1)	(0,1,1,1)	(1,1,1,1)

32
33
34
35
36
37
38
39
40
41 Table 2: Filters and vertices of poset of Figure 6.

42
43
44
45 Vertices defining an extremal ray are those whose corresponding filter is connected. The five
46 vertices in these conditions are written in boldface.

47
48 In order to obtain the facets of this order cone, we look for facets of the corresponding order
49 polytope containing $\mathbf{0}$ (Theorem 8). For this, we consider $\perp \oplus P \oplus \top$ (see Figure 6 right). As we are
50 looking for facets, we just turn an inequality not involving \top into an equality. Then, the facets are
51 given in Table 3.

52
53 Another way to look for extremal rays is Theorem 6. For this, we need to build the lattice of
54 filters, that is given in Figure 7.

55
56 Then, the extremal rays are given by filters that together with \emptyset form an embedded sublattice.
57 These filters are

$$\{123\}, \{34\}, \{12, 123\}, \{13, 123\}, \{12, 13, 123\}.$$

Restriction	$f(\perp) = f(12)$	$f(\perp) = f(13)$	$f(\perp) = f(34)$	$f(12) = f(123)$	$f(13) = f(123)$
Vertices	(0,0,0,0)	(0,0,0,0)	(0,0,0,0)	(0,0,0,0)	(0,0,0,0)
	(0,0,0,1)	(0,0,0,1)	(0,0,0,1)	(1,0,0,1)	(0,1,0,0)
	(0,0,1,0)	(0,0,1,0)	(1,0,0,1)	(1,0,1,1)	(0,1,1,1)
	(0,0,1,1)	(0,0,1,1)	(0,1,0,1)	(1,1,0,1)	(1,1,0,1)
	(0,1,0,1)	(0,1,0,1)	(1,1,0,1)	(1,1,1,1)	(1,1,1,1)
	(0,1,1,1)	(1,0,1,1)			

Table 3: Facets of the order cone of poset of Figure 6.




Figure 7: Lattice of filters.

4.3 The cone of k -symmetric measures

As explained before, order cones can be applied to more general situations than games with restricted cooperation. In this subsection we will apply it to k -symmetric monotone games. We have chosen this case because the set of k -symmetric capacities with respect to a fixed partition is an order polytope (Combarro and Miranda, 2010).

The concept of k -symmetry appears in the theory of capacities as an attempt to reduce the complexity (Miranda et al., 2002). The subjacent idea is that several players could act exactly in the same way, so that we do not need to care about which players in these conditions are in a coalition and we just need to know how many players are inside it. The key concept of k -symmetric monotone game is *subset of indifference*. Basically, a subset of indifference is a group of indistinguishable elements in terms of game v . Mathematically, this translates into

$$v(B_1 \cup C) = v(B_2 \cup C), \forall C \subseteq X \setminus A, B_1 B_2 \subset A, |B_1| = |B_2|.$$

This allows us to write a coalition in terms of the number of players inside each subset of indifference.

Lemma 6. (Miranda et al., 2002) *If $\{A_1, \dots, A_k\}$ is a partition of indifference for N , then any $C \subseteq N$ can be identified with a k -dimensional vector (c_1, \dots, c_k) with $c_i := |C \cap A_i|$.*

Then, each coalition writes (c_1, \dots, c_k) with $c_i = 0, \dots, |A_i|$. For a given game v , it can be seen that it is always possible to partitionate N in several subsets of indifference. Several partitions are possible, but it can be proved (Miranda et al., 2002) that there is an only one being the coarsest.

Definition 5. *We say that a game is k -symmetric with respect to the partition A_1, \dots, A_k if this is the coarsest partition of N in subsets of indifference.*

We denote by $\mathcal{MG}^k(A_1, \dots, A_k)$ the set of monotone games v such that A_1, \dots, A_k are subsets of indifference for v (but not necessarily k -symmetric; for example, any symmetric monotone game, in which all players are indifferent, belongs to $\mathcal{MG}^k(A_1, \dots, A_k)$). Then, $v \in \mathcal{MG}^k(A_1, \dots, A_k)$ is characterized as follows:

- $v(0, \dots, 0) = 0$.
- $v(a_1, \dots, a_k) \leq v(b_1, \dots, b_k)$ if $a_i \leq b_i, i = 1, \dots, k$.

Consider then the poset

$$P = \{(c_1, \dots, c_k) : c_i = 0, \dots, |A_i|, i = 1, \dots, k\}$$

with the order relation $(c_1, \dots, c_k) \preceq (b_1, \dots, b_k)$ if and only if $c_i \leq b_i, i = 1, \dots, k$.

Then, it follows that $\mathcal{MG}^k(A_1, \dots, A_k) = \mathcal{C}(P \setminus \{(0, \dots, 0)\})$ and the results of Section 3 can be applied to obtain the geometrical aspects of this cone. Moreover, as $(|A_1|, \dots, |A_k|)$ is a top element in the poset, we can apply the results obtained for $\mathcal{MG}(N)$.

Corollary 9. *The vectors defining an extremal ray of $\mathcal{MG}^k(A_1, \dots, A_k)$ are defined by non-empty filters of $P \setminus \{(0, \dots, 0)\}$.*

Proposition 7. *Consider the poset $P = \{(c_1, \dots, c_k) : c_i = 0, \dots, |A_i|, i = 1, \dots, k\}$. Then, the order polytope $\mathcal{O}(P \setminus \{(0, \dots, 0)\})$ is a pyramid with base $\{(\mathbf{x}, 1) : \mathbf{x} \in \mathcal{O}(P \setminus \{(0, \dots, 0), (|A_1|, \dots, |A_k|)\})\}$ and apex $\mathbf{0}$.*

Corollary 10. *The k -dimensional faces of $\mathcal{MG}^k(A_1, \dots, A_k)$ are given by the $(k - 1)$ -dimensional faces of $\mathcal{O}(P \setminus \{(0, \dots, 0), (|A_1|, \dots, |A_k|)\})$.*

Let us study two particular cases.


Example 6. *For $\mathcal{MG}^1(N)$, the set of monotone symmetric games, the corresponding order polytope is a chain of n elements. Thus, we have n non-empty filters F_1, \dots, F_n , given by $F_i := \{i, \dots, n\}$ and $\mathbf{v}_{F_i} = (0, \dots, 1, \dots, 1)$. Therefore, we have n extremal rays.*

Besides, by Theorem 4, we conclude that all vertices are adjacent to each other. Hence, we have $\binom{n}{2}$ 2-dimensional faces and in general, the number of k -dimensional faces is $\binom{n}{k}$, for $k \geq 2$.

1
2
3 **Example 7.** For the 2-symmetric case $\mathcal{MG}^2(A_1, A_2)$, it has been proved in (García-Segador and
4 Miranda, 2020) that the order polytope $\mathcal{FM}^2(A_1, A_2)$ can be associated to a Young diagram (Bandlow,
5 2008) of shape $\lambda = (|A_2|, \dots, |A_2|)$.

6 Moreover, there is a correspondence between filters and staircase walks from $(0, 0)$ to (a_1, a_2) in
7 a $(|A_1| + 1) \times (|A_2| + 1)$ grid (see Figure 8). Cell (i, j) represents the subset (i, j) . In this sense,
8 the walk separates subsets with value 0 from subsets with value 1 (see (García-Segador and Miranda,
9 2020)). For example, the empty filter corresponds to the staircase walk going from $(0, 0)$ to $(a_1, 0)$
10 and then to (a_1, a_2) .
11
12

13
14 Figure 8: Staircase walk in a 4×4 grid and a staircase walk.



15
16
17
18
19
20
21
22
23
24
25
26
27 Then, the number of vertices of $\mathcal{FM}^2(A_1, A_2)$ is the number of possible staircase walks, that is
28 given by
29

$$\binom{a_1 + a_2 + 2}{a_1 + 1},$$

30
31
32
33
34 and by Corollary 9 the number of vertices determining an extremal ray is $\binom{a_1 + a_2 + 2}{a_1 + 1} - 1$.
35
36

37 5 Conclusions

38
39
40 In this paper we have introduced the concept of order cones. This concept is a natural extension of
41 order polytopes, a well-known object in Combinatorics with which order cones share many properties.
42 We have shown that all order cones are pointed, and we have derived some of their geometrical
43 properties. Namely, we have characterized its k -dimensional faces. In particular, we have obtained
44 a characterization of extremal ray in terms of the corresponding subjacent poset. The results in the
45 paper show that the geometrical structure of order cones can be derived from the order structure of
46 the subjacent poset, thus simplifying many results.
47
48

49 We feel that order cones could be a powerful tool to study different cones appearing in Game
50 Theory in a general way. As examples of applicability, in the second part of the paper, we have
51 applied these results to some special subfamilies of monotone games that satisfy the conditions of
52 order cone. We have shown that the results derived in the first part can be applied to the set of
53 monotone games with restricted cooperation, no matter the structure of the set of feasible coalitions.
54 Then, we have studied in the first place the set of monotone games when all coalitions are allowed.
55 For this case, we have shown that it is closely related to the order polytope of capacities. In a second
56 step, we have studied this set when a set of feasible coalitions arises. We have shown that the set of
57 monotone games with restricted cooperation always leads to an order cone whose structure relies on
58 the poset of feasible coalitions. And we have seen that roughly speaking, there are two possible cases:
59
60
61
62
63
64
65

1
2
3 the one with a top element (usually N) as a feasible coalition, that is very similar to the general
4 case, and the case where there are several maxima, that leads to a more complicated problem.

5 Finally, we have studied an example where an order cone arises if constraints are added to the
6 values of the game. This shows that order cones can be applied to situations different of monotone
7 games with restricted cooperation. More concretely, we have studied the set of k -symmetric monotone
8 games.
9

10 We also feel that the concept of order cone could be an interesting tool for studying several families
11 of monotone games just focusing on the subjacent poset. Note on the other hand that the order
12 relation is essential for order cones. This means that the definition fails if we remove monotonicity.
13 Studying a generalization dealing with this situation seems to be a complex problem that we intend
14 to study in the future.
15
16

17 18 19 **References**

20
21 Bandlow, J. 2008. An elementary proof of the hook formula. *The Electronic Journal of Combinatorics*,
22 (15). Research paper 45.

23
24 Choquet, G. 1953. Theory of capacities. *Annales de l'Institut Fourier*, (5):131–295.

25
26 Combarro, E. F. and Miranda, P. 2008. On the polytope of non-additive measures. *Fuzzy Sets and*
27 *Systems*, 159(16):2145–2162.

28
29 Combarro, E. F. and Miranda, P. 2010. Adjacency on the order polytope with applications to the
30 theory of fuzzy measures. *Fuzzy Sets and Systems*, 180:384–398.

31
32 Cook, W., Cunningham, W., Pulleyblank, W. and Schrijver, A. 1988. *Combinatorial Optimization*,
33 Wiley Series in Discrete Mathematics and Optimization. Wiley.

34
35 Davey, B. A. and Priestley, H. A. 2002. *Introduction to lattices and order*. Cambridge University
36 Press.

37
38 Denneberg, D. 1994. *Non-additive measures and integral*. Dordrecht (The Netherlands): Kluwer
39 Academic.

40
41 Faigle, U. 1989. Cores of games with restricted cooperation. *Zeitschrift für Operations Research*,
42 33(6):405–422.

43
44 Friedl, T. 2017. *Double posets and real invariant varieties: Two interactions between combinatorics*
45 *and geometry*. PhD thesis, Fachbereich Mathematik und Informatik der Freien Universität Berlin-
46 Tokyo Institute of Technology.

47
48 García-Segador, P. and Miranda, P. 2020. Applying young diagrams to 2-symmetric fuzzy measures
49 with an application to general fuzzy measures. *Fuzzy Sets and Systems*, 379:20–36.

50
51 Grabisch, M. 2011. Ensuring the boundedness of the core of games with restricted cooperation.
52 *Annals of Operations Research*, 191:137–154.

53
54 Grabisch, M. 2013. The core of games on ordered structures and graphs. *Annals of Operations*
55 *Research*, 204:33–64. doi: 10.1007/s10479-012-1265-4.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Grabisch, M. 2016. *Set functions, games and capacities in Decision Making*, volume 46 of *Theory and Decision Library*. Springer.

Grabisch, M. and Kroupa, T. 2019. The cone of supermodular games on finite distributive lattices. *Discrete Applied Mathematics*, 260:144–154.

Katsev, I. and Yanovskaya, E. 2013. The prenucleolus for games with restricted cooperation. *Mathematical Social Sciences*, 66(1):56–65.

Miranda, P., Grabisch, M. and Gil, P. 2002. p -symmetric fuzzy measures. *International Journal of Uncertainty, Fuzziness and Knowledge-Based Systems*, 10 (Suppl.):105–123.

Pulido, M. and Sánchez-Soriano, J. 2006. Characterization of the core in games with restricted cooperation. *European Journal of Operational Research*, 175(2):860–869.

Shapley, L. S. 1971. Cores of convex games. *International Journal of Game Theory*, 1:11–26.

Stanley, R. 1986. Two poset polytopes. *Discrete Comput. Geom.*, 1(1):9–23.

Sugeno, M. 1974. *Theory of fuzzy integrals and its applications*. PhD thesis, Tokyo Institute of Technology.